

UNIVERSITÀ DEGLI STUDI DI TRIESTE

DOTTORATO DI RICERCA IN FISICA - XV CICLO

ANNO ACCADEMICO 2001-2002

Dissipative Dynamics in Particle Physics

Raffaele Romano

Ph.D. Thesis



TUTORE: Dott. Fabio Benatti (Università di Trieste)

COORDINATORE: Prof.ssa Francesca Matteucci (Università di Trieste)

DISSIPATIVE DYNAMICS IN PARTICLE PHYSICS

Raffaele Romano

Contents

Introduction	3
1 Open Quantum Systems	7
1.1 States, observables and dynamical maps	7
1.2 Positivity and Complete Positivity	11
1.3 Reduced dynamics: the Generalized Master Equation	14
1.4 The Markovian approximation. The Weak Coupling Limit	19
1.5 Quantum Dynamical Semigroups	21
1.6 Factorized dynamics of quantum dynamical semigroups	25
1.7 The stochastic approach	27
1.8 Entropy and long time behavior in open quantum systems	30
2 Bidimensional Systems	33
2.1 Evolution of states	33
2.2 Positivity and Complete Positivity	36
2.3 Positivity and CP for entropy increasing evolutions	38
2.4 Perturbative expansion	39
3 Neutral B Mesons	45
3.1 The physics of B mesons	45
3.2 The gravitational bath	48
3.3 The dissipative evolution	49
3.4 Invariance properties	51
3.5 Observables	53
3.6 Decay rates for a single B meson	55
3.7 Asymmetries for a single B meson	56
3.8 Decay rates for entangled B meson	59
3.9 Asymmetries for entangled B meson	61
3.10 Discussion and conclusions	64

4	Neutrinos	67
4.1	The physics of neutrinos in vacuum	68
4.2	The model of interaction	70
4.3	The Markovian approximation	72
4.4	The generator of the dissipative evolution	75
4.5	An example of flavor changing dissipative evolution	79
4.6	Non homogeneous matter	81
4.7	The solar matter case	81
4.8	Discussion and conclusions	82
5	Neutron interferometry	85
5.1	The physical apparatus	86
5.2	The stochastic thermal bath	89
5.3	Three different stochastic backgrounds	92
5.3.1	White noise	93
5.3.2	Diagonal covariance matrix	93
5.3.3	Single component field correlation	94
5.4	Positivity and complete positivity	95
5.5	Noncontextuality: the Clauser-Horne-Shimony-Holt inequality	97
5.5.1	White noise	99
5.5.2	Diagonal covariance matrix	100
5.5.3	Single component field correlation	100
5.6	Discussion and conclusions	101
A	Dissipative parameters in the eigenstates basis	105
B	Expressions of the time evolutors	109
C	Neutral B mesons decays	111
C.1	Semileptonic decays	111
C.2	Decay in a CP eigenstate f	112
D	Dissipative parameters in the neutrino case	115
E	Frequencies of counts	117
	Bibliography	119

12 - *Vom Ziele der Wissenschaft* - Wie? Das letzte Ziel der Wissenschaft sei, dem Menschen möglichst viel Lust und möglichst wenig Unlust zu schaffen? Wie, wenn nun Lust und Unlust so mit einem Stricke zusammengeknüpft wären, dass, wer möglichst viel von der einen haben will, auch möglichst viel von der andern haben muss,-dass, wer das "Himmelhoch-Jauchzen" lernen will, sich auch für das "zum-Todebetrübt" bereit halten muss? Und so steht es vielleicht! Die Stoiker glaubten wenigstens, dass es so stehe, und waren consequent, als sie nach möglichst wenig Lust beehrten, um möglichst wenig Unlust vom Leben zu haben (wenn man den Spruch im Munde führte "Der Tugendhafte ist der Glücklichsste", so hatte man in ihm sowohl ein Aushängeschild der Schule für die grosse Masse, als auch eine casuistische Feinheit für die Feinen). Auch heute noch habt ihr die Wahl: entweder möglichst wenig Unlust, kurz Schmerzlosigkeit-und im Grunde dürften Socialisten und Politiker aller Parteien ihren Leuten ehrlicher Weise nicht mehr verheissen-oder möglichst viel Unlust als Preis für das Wachstum einer Fülle von feinen und bisher selten gekosteten Lüsten und Freuden! Entschliesst ihr euch für das Erstere, wollt ihr also die Schmerzhaftigkeit der Menschen herabdrücken und vermindern, nun, so müsst ihr auch ihre Fähigkeit zur Freude herabdrücken und vermindern. In der Tat kann man mit der Wissenschaft das eine wie das andere Ziel fördern! Vielleicht ist sie jetzt noch bekannter wegen ihrer Kraft, den Menschen um seine Freuden zu bringen, und ihn kälter, statuenhafter, stoischer zu machen. Aber sie könnte auch noch als die grosse Schmerzbringerin entdeckt werden!-Und dann würde vielleicht zugleich ihre Gegenkraft entdeckt sein, ihr ungeheures Vermögen, neue Sternenwelten der Freude aufleuchten zu lassen!

F. Nietzsche, *Die fröhliche Wissenschaft* ("la gaya scienza"), 1882

Introduction

The subject of this thesis is the study of dissipative dynamics and their properties in particle physics.

Physical systems can be considered closed when they are isolated from their environment; otherwise, due to interactions between them, there occur exchanges of energy, entropy and matter from system to environment and vice versa. Physical systems are termed open in such a case.

Typically, the physical systems we will be concerned with have few degrees of freedom in comparison to the very (infinitely) many of the environment and matter exchanges are neglected. Closed quantum systems evolve reversibly in time according to the unitary dynamics generated by the Schrödinger equation; in particular, there cannot be decoherence since Hilbert space vectors evolve in time into Hilbert space vectors.

On the contrary, the time-evolution of open quantum systems is quite complicated due to the coupling with the environment. It can be formally derived from the unitary dynamics of the closed system formed by the open system plus the environment, by eliminating (or summing over) the degrees of freedom of the latter.

The result is a so-called reduced dynamics for the open system alone that does not conserve its energy and entropy; the dynamics is thus dissipative and, moreover, does not transform pure states into themselves, rather it destroys quantum coherence.

The reduced dynamics generally contains memory effects, that is it depends not only on the initial conditions and the observation time, but on the whole history of the system in between. However, if the coupling with the environment is weak, it is possible to approximate this dynamics with semigroups of linear transformations in which the memory effects have disappeared, namely by Markovian semigroups which constitute the class of time-evolutions studied in this thesis [1, 2, 3].

Open quantum systems are met in many branches of physics; in particular, the approach based on semigroups of linear transformations has been applied in several contexts, ranging from phenomena of quantum relaxation in magnetic resonance [4], to quantum optics [5, 6, 7, 8], to foundational aspects of quantum mechanics [9, 13], to the study of non-standard dynamics in elementary particle physics [44, 49, 117, 118] and of interference phenomena with photons or neutrons [10, 11, 12].

Since reduced dynamics are decohering, the states of open systems must be described by density matrices; their eigenvalues represent probabilities, so that they have to be

positive. In order to have a consistent statistical interpretation of the formalism, the dynamical maps describing the time-evolution must preserve such positivity; technically speaking, they need to be positivity preserving or positive for short. However, when there is entanglement between the open system and its surroundings, positivity is no longer sufficient to guarantee the consistency of the physical description. A stronger request need to be enforced: the so-called complete positivity. Formally, complete positivity assures the positivity of states of composite systems in which there is a statistical but not a dynamical coupling between the system under study and a generic n -level system.

Due to its rather abstract definition, complete positivity is sometime considered as a mere mathematical property rather than a physically motivated request. Indeed, the literature about open quantum systems abounds in reduced dynamics that are not completely positive (and some examples will be discussed below) [125, 126, 27].

The physical systems and the discussions presented in this thesis provide some arguments in favor of complete positivity by showing explicitly the unavoidable problems that arise by considering dynamics that are not completely positive. Moreover, since the models that will be presented represent concrete systems amenable to actual experiments, they offer the possibility of direct tests of this property.

In Chapter 1, after introducing the basics of positivity and complete positivity for generic time-evolutions, we develop the formalism leading to the reduced dynamics of open quantum systems; in particular the weak coupling limit and the resulting general structure of semigroups.

In Chapter 2 we restrict our attention to open quantum systems described by two-dimensional Hilbert spaces, for the concrete physical systems we shall consider can be described in this framework. We introduce the coefficients that characterize general irreversible reduced dynamics fulfilling the Markov property; in order to guarantee either the positivity or the complete positivity request, they must satisfy some constraints, expressed by sets of inequalities.

In the next three chapters this formalism is applied to three specific contexts: the study of irreversible dynamics of neutral **B** mesons in a gravitational environment (Chapter 3), of a beam of neutrinos crossing a layer of matter (Chapter 4), and finally of neutrons in interferometric experiments in presence of stochastic magnetic fields (Chapter 5).

The analysis of a possible dissipative time-evolution for the neutral **B** mesons is suggested by general considerations about the behavior of quantum gravity (or, from a more fundamental level, string theory) at the Planck scale. The approach is phenomenological, that is the microscopic details of the interaction between system and gravitational bath are unspecified; in this sense, the adopted dissipative time-evolution for the **B** mesons is completely general. We have explicitly evaluated the expressions of the decay probabilities of neutral mesons for some selected decay channels and, by means of them, some asymmetries, that is quantities usually employed to evaluate the violation of the discrete symmetries (parity, time reversal and charge conjugation). These asymmetries can be directly used to fit experimental data. In this respect, entangled **B** mesons produced in the decay of spin-1 resonances are particularly useful: in such a case, the dissipative effects

induced by the non-standard dynamics produce characteristic signatures, that, at least in principle, can be experimentally probed. Indeed, the experiments involving entangled neutral **B** mesons at the so-called **B**-factories have reached a high level of sensitivity, very close to the one required for detecting the non-standard, dissipative phenomena. This is a further motivation for the investigations presented here. Moreover, since complete positivity is tightly related to the quantum correlations expressed by entanglement, the **B**-factories offer the possibility of performing experimental tests of this fundamental property.

The analysis of neutrino interactions and oscillations in matter described in Chapter 4 is not purely phenomenological in character, rather it is based on a microscopic model. The formalism is nevertheless very general and, provided the Markovian approximation is valid, it can be applied to study dissipative phenomena in neutrino physics induced by various matter effects, e.g. the one induced by the thermal fluctuations inside the Sun. We have computed the modified form of the transition probability between two different flavors; for thermal fluctuations in the Sun we find no meaningful changes to the standard expression, in agreement with previous results coming from different approaches.

Finally, in the last chapter, we devote our attention to neutron interferometers and discuss how these apparatus can be used to study dissipative dynamics, in a controlled environment. The basic idea is to insert the interferometer in an external stochastic magnetic field. By suitably adjusting the parameters characterizing the fluctuating external field, one can then reproduce the physical situations adopted to justify widely used dynamics (e.g. of the Redfield-Bloch type). It turns out that such time-evolutions are not completely positive and sometimes not even positive; neutron interferometry would then allow direct experimental tests of the inconsistencies that such dynamics necessarily produce.

Neutron interferometers have been proposed as one of the possible set-ups to study non-contextuality in quantum mechanics, by testing suitable Bell's inequalities. In presence of dissipative effects leading to loss of quantum coherence, one expects a weakening of the effectiveness of these Bell's inequality tests. As a byproduct of the previous investigations, we have studied in detail how the expressions of these inequalities change under the action of dissipative dynamics.

In this thesis, elementary particle phenomenology and quantum interferometry are shown to provide, under specific conditions, very good laboratories to investigate the properties of reduced dynamics of open systems, in particular the notion of complete positivity. The motivations about relevance and necessity of complete positivity result strengthened by the arguments developed in what follows.

Chapter 1

Open Quantum Systems

In this chapter we introduce the concept of quantum open systems and their reduced time-evolutions. After some basic definitions, we focus our attention on the properties of positivity and complete positivity of reduced time-evolutions, stressing their relevance and the problems that arise if they are not positive or not completely positive. Complete positivity shall be the leit-motiv of this thesis, since it is usually believed that any dynamical map describing the time-evolution of an open quantum system should enjoy this property in order to avoid any fallacy in the interpretation of the formalism. However, in the literature about open quantum systems many examples of not completely positive reduced dynamics are encountered; we will present some of them to show the problems they suffer.

In Section 1.3 we explicitly show how, by means of the projector formalism, it is possible to extract the dynamics of a subsystem of a larger system whose time-evolution is known. The reduced dynamics so obtained is in general rather complex because it contains memory effects negligible only when some particular conditions are fulfilled, typically involving the weakness of the coupling between system and environment. In view of this, in Section 1.4 we introduce the so-called Markovian dynamics, that is the class of time-evolutions in which memory terms are made to disappear; they are described by linear maps satisfying the semigroup composition law. This is the class of dynamics we shall consider in the following chapters.

A different approach to the time-evolution of an open quantum system is presented in Section 1.7, based on a Langevin-type equation.

In the final part of this chapter some considerations about the long-time behavior of states evolving under a Markovian dynamics are included.

1.1 States, observables and dynamical maps

In this section we introduce the basic elements to describe a quantum system: the states, the observables and their time-evolutions.

Given a Hilbert space $\mathfrak{H}$, we recall the following definitions:

Definition 1.1 A linear operator A acting on $\mathfrak{H}$ is said to be *bounded* iff $\|A\|_\infty < \infty$, where

$$\|A\|_\infty \equiv \sup_{\|\psi\| \leq 1} \|A\psi\|, \quad \psi \in \mathfrak{H}.$$

Definition 1.2 A linear operator A acting on $\mathfrak{H}$ is said to be *trace class* iff $A \in \mathcal{B}(\mathfrak{H})$ and $\|A\|_1 \equiv \text{Tr}(A^\dagger A)^{1/2} < +\infty$.

We introduce two complex linear spaces of operators acting on $\mathfrak{H}$ [14]:

- $\mathcal{B}(\mathfrak{H})$: the complex linear space of bounded operators, which is complete in the operator norm $\|\cdot\|_\infty$;
- $\mathcal{T}(\mathfrak{H})$: the complex linear space of trace-class operators, which is complete in the trace norm $\|\cdot\|_1$.

Remark 1.1 If $\mathfrak{H}$ is finite-dimensional, say $\dim(\mathfrak{H}) = n$, then $\mathcal{B}(\mathfrak{H}) = \mathcal{T}(\mathfrak{H}) = M_n(\mathbf{C})$, the algebra of $n \times n$ matrices. If $\dim(\mathfrak{H}) = \infty$, $\mathcal{T}(\mathfrak{H}) \subset \mathcal{B}(\mathfrak{H})$; in particular $\mathcal{B}(\mathfrak{H})$ is an algebra whereas $\mathcal{T}(\mathfrak{H})$ is not.

The physical states of quantum systems with finitely many degrees of freedom can always be represented by positive (then Hermitian) trace-class operators with trace one¹, called density matrices or statistical operators:

$$\rho \in \mathcal{T}(\mathfrak{H}), \quad \rho \geq 0, \quad (\rho^\dagger = \rho), \quad \text{Tr}(\rho) = 1. \quad (1.1)$$

They constitute a convex set in $\mathcal{T}(\mathfrak{H})$, denoted by $\mathcal{P}(\mathfrak{H})$, whose extremal points are projectors on one-dimensional subspaces; in spectral representation, they read

$$\rho = \sum_i \lambda_i |e_i\rangle \langle e_i|; \quad \langle e_i | e_j \rangle = \delta_{ij}, \quad \lambda_i \geq 0, \quad \sum_i \lambda_i = 1. \quad (1.2)$$

Definition 1.3 A state ρ is said to be *pure* if the sum in (1.2) can be reduced to only one term (after a suitable choice of the basis vectors $|e_i\rangle$), otherwise it is said to be *mixed* and it represents a statistical convex superposition of pure states. In such a case the eigenvalues are the statistical weights of the corresponding eigenprojectors in the mixture.

Remark 1.2 Density matrices are not uniquely expressible as convex linear combinations of other states. For instance any mixture $\sum_i \mu_i \sigma_i$ with generic density matrices σ_i and $\mu_i > 0$, $\sum_i \mu_i = 1$, can always be spectralized and written also in the form (1.2)

Pure states are fully characterized as follows.

¹Or, as we shall shortly see, less than one if the state represents an unstable system

Remark 1.3 A state $\rho \in \mathcal{T}(\mathfrak{H})$ is pure iff $\rho^2 = \rho$.

The system observables are represented by Hermitian operators $A = A^\dagger$ belonging to $\mathcal{B}(\mathfrak{H})$. The expectation value of physical observables A with respect to a state ρ , in this formalism, is given by the trace operation:

$$\langle A \rangle_\rho = \text{Tr}(\rho A) = \sum_i \lambda_i \langle e_i | A | e_i \rangle. \quad (1.3)$$

In order to describe the time-evolution of quantum systems, two approaches are possible. In the Schrödinger picture a state evolves as $\rho(0) \rightarrow \rho(t) = \Lambda_t[\rho(0)]$ where Λ_t is a linear operator, $\Lambda_t : \mathcal{P}(\mathfrak{H}) \rightarrow \mathcal{P}(\mathfrak{H})$ and t is time; the one-parameter family of linear maps $\{\Lambda_t, t \in \mathfrak{R}\}$ is called time-evolution of the system; $\mathfrak{R}$ is the range over which t can vary and it depends on the characteristics of the considered dynamics.

In the Heisenberg picture, we introduce the one-parameter family of linear maps $\{\Lambda_t^*, t \in \mathfrak{R}\}$, producing the time-evolution of the observables: $A(0) \rightarrow A(t) = \Lambda_t^*[A(0)]$. Λ_t^* is called the dual map to Λ_t , defined by the duality relation:

$$\text{Tr}[\rho(t)A] = \text{Tr}[\rho A(t)]. \quad (1.4)$$

We distinguish the following classes of dynamical maps.

Definition 1.4 A dynamical map is called *reversible* iff Λ_t admits an inverse Λ_t^{-1} , $\forall t \in \mathfrak{R}$. Otherwise it is called *irreversible*.

Definition 1.5 A dynamical map is called *trace-preserving* iff $\text{Tr}[\rho(t)] = \text{Tr}[\rho(0)]$, $\forall t \in \mathfrak{R}$.

For the physical consistency of the formalism, any dynamical map must be Hermiticity-preserving, that is $\Lambda_t[\rho]^\dagger = \Lambda_t[\rho]$, $\forall \rho \in \mathcal{P}(\mathcal{H})$, $\forall t \in \mathfrak{R}$. Instead, reversibility and trace preservation may be satisfied or not, depending on the physical systems we are dealing with.

In particular, trace-preserving maps describe stable systems in which the total probability is conserved; on the other hand, the time-evolution of unstable systems (that is decaying systems) is modeled by maps that do not preserve the trace of states: the total probability has indeed to decrease in time.

The standard quantum dynamics is given by the time-evolution of stable closed systems, that in the Schrödinger picture reads (putting $\hbar = 1$):

$$\rho(t) = \Lambda_t[\rho(0)] = e^{-iHt} \rho(0) e^{iHt} \quad (1.5)$$

or, in differential form,

$$\dot{\rho}(t) = -i[H, \rho(t)], \quad (1.6)$$

where $H = H^\dagger$ is the Hamiltonian describing the system. In the following we denote by $L_H[\cdot] = -i[H, \cdot]$ the Liouville operator which generates of the time-evolution maps (or dynamical maps):

$$\Lambda_t = e^{L_H t} \quad \forall t \in \mathbf{R}. \quad (1.7)$$

The same evolution in the Heisenberg picture reads

$$A(t) = \Lambda_t^*[A(0)] = e^{iHt} A(0) e^{-iHt}, \quad (1.8)$$

which is generated by

$$\dot{A}(t) = i[H, A(t)], \quad (1.9)$$

where A is any observable of the system.

In such a case, the set of linear operators $\{\Lambda_t, t \in \mathbf{R}\}$ (and the set of their dual maps $\{\Lambda_t^*, t \in \mathbf{R}\}$) forms a group of transformations. The group structure implies that for each Λ_t there is an inverse map $\Lambda_t^{-1} = \Lambda_{-t}$, that is the dynamics is reversible; moreover the composition law $\Lambda_{t_1+t_2} = \Lambda_{t_1} \circ \Lambda_{t_2}$, $\forall t_1, t_2 \in \mathbf{R}$ is satisfied, as can be seen from (1.7). The invariance of the trace under cyclic permutations produces the trace-preserving property for the dynamical maps. Moreover, it is Hermiticity-preserving and it maintains the coherence of the states over which it acts: indeed, from (1.5), $\rho^2(t) = \rho(t)$. This is another expression of the fact that Hilbert space vectors $|\psi\rangle$ are mapped into themselves: $|\psi\rangle \rightarrow |\psi_t\rangle = e^{-iHt}|\psi\rangle$.

Another example of dynamical map is the one describing a closed unstable system. The description of decaying systems is performed introducing a non-Hermitian Hamiltonian H in the Weisskopf-Wigner form [20], that is composed by a Hermitian part and an anti-Hermitian one:

$$H = M - \frac{i}{2}\Gamma; \quad M = M^\dagger, \quad \Gamma = \Gamma^\dagger \quad (1.10)$$

with positive M and Γ characterizing masses and natural widths of decaying states. In this case, in the Schrödinger picture, the states evolution reads

$$\rho(t) = \Lambda_t[\rho(0)] = e^{-iHt} \rho(0) e^{iH^\dagger t} \quad (1.11)$$

or, in differential form,

$$\dot{\rho}(t) = -iH\rho(t) + i\rho(t)H^\dagger = -i[M, \rho(t)] - \frac{1}{2}\{\Gamma, \rho(t)\} \quad (1.12)$$

leading to the property

$$\frac{d}{dt} \text{Tr}[\rho(t)] = -\text{Tr}[\rho(t)\Gamma] < 0. \quad (1.13)$$

In the Heisenberg picture the time-evolution of the observables is described by equations analogous to (1.11) and (1.12).

In this case the range $\mathfrak{R}$ over which the dynamics makes sense is not the real axis $\mathbf{R}$ but rather the positive semi-axis $\mathbf{R}^+$. Indeed for $t < 0$ equation (1.13) should imply $\text{Tr}[\rho(t)] > 1$ since $\text{Tr}[\rho(0)] = 1$, and thus the statistical interpretation of the eigenvalues of $\rho(t)$ would be compromised. Since this time-evolution is defined only for positive time, it is irreversible; moreover it is not trace-preserving, as can be seen from (1.13), and it maintains the coherence of the states, i.e. Hilbert space vectors transform into Hilbert space vectors, though not normalized, since $H \neq H^\dagger$.

In the study of open quantum systems we will meet irreversible time-evolutions either trace-preserving or not, and decohering, that is causing the transition from pure to mixed states.

We have to discuss the fundamental property of the preservation of positivity of the states subject to a dynamical map. For its relevance and its prominent interest we reserve to this issue the next section.

1.2 Positivity and Complete Positivity

Let us start with the definitions of positivity and complete positivity:

Definition 1.6 A linear transformation $\Gamma : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ is said to be *positive* (or *positivity-preserving*) iff $A \geq 0 \Rightarrow \Gamma(A) \geq 0$, $\forall A \in \mathcal{B}(\mathfrak{H})$.

Definition 1.7 A linear transformation $\Gamma : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ is said to be *n-positive* iff $\Gamma \otimes \mathbf{I}_n : \mathcal{B}(\mathfrak{H} \otimes \mathfrak{H}_n) \rightarrow \mathcal{B}(\mathfrak{H} \otimes \mathfrak{H}_n)$ is positive, with $n \in \mathbf{N}$ and $\mathbf{I}_n$ the identity on the n -dimensional Hilbert space $\mathfrak{H}_n$.

Definition 1.8 A linear transformation $\Gamma : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ is said to be *completely positive* (CP) iff it is n -positive $\forall n \in \mathbf{N}$.

Remark 1.4 Complete positivity is a stronger property than positivity which amounts to 1-positivity.

Remark 1.5 Historically, the notion of complete positivity was introduced for linear transformations acting on bounded operators, that is, in a dynamical context, not for the time-evolution of states, Λ_t , but for its dual Λ_t^* , the time-evolution of operators. In this thesis, the notion of complete positivity will be referred to dynamical maps on states in the sense that a map $\Lambda_t : \mathcal{P}(\mathfrak{H}) \rightarrow \mathcal{P}(\mathfrak{H})$ is CP if and only if its dual map $\Lambda_t^* : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ is CP.

In the context of dynamical maps describing generic time-evolutions of states, the property of positivity is required in order to preserve at any time the positivity of the

eigenvalues of the statistical operator representing the system. As previously stated, these eigenvalues represent the weights of a statistical superposition; then they must be positive in order to guarantee the correct probabilistic interpretation of the formalism. For any non-positive time-evolution Λ_t there necessarily exists a positive initial state $\rho(0)$ mapped by Λ_t in a non-positive one, $\rho(t) = \Lambda_t[\rho(0)]$, leading to physical inconsistency of the formalism.

It is less intuitive to understand the relevance and necessity of complete positivity for time-evolutions of states. Let Λ_t be a generic dynamical map describing the time-evolution of a state ρ_S ; following the definition of CP, we have to consider the map $\Lambda_t \otimes \mathbf{I}_n$, where $\mathbf{I}_n$ is the identity in a n -dimensional Hilbert space and n is any natural number. This map represents the factorized time-evolution of a composite system $T = S + S_n$, where S is the system described by ρ_S and evolving under Λ_t while S_n is a generic n -level system characterized by a trivial Hamiltonian $H_n = 0$. Since the dynamics of the composite system has the factorized form $\Lambda_t \otimes \mathbf{I}_n$ there is no dynamical coupling between S and S_n . Now, the positivity of any state ρ_T , describing the composite system T , must be preserved under the action of $\Lambda_t \otimes \mathbf{I}_n$, but this is not guaranteed by the positivity of Λ_t : a completely positive Λ_t is needed.

To clarify this point, we define a fundamental property of the state describing the composite system T , tightly connected to the notion of complete positivity.

Definition 1.9 Given a state ρ_T describing a composite system $T = S_1 + S_2$, it is called *separable* if it can be written as $\rho_T = \sum_{i,j} c_{ij} \rho_{S_1}^i \otimes \rho_{S_2}^j$ with $c_{ij} > 0, \forall i, j$; otherwise it is called *entangled*.

It turns out that the request of complete positivity is necessary when the evolution of entangled states ρ_T is taken into account. Indeed, any factorized statistical operator ρ_T of the bipartite system $T = S + S_n$, evolving under the time-evolution $\Lambda_t \otimes \mathbf{I}_n$, where Λ_t is positive-definite, is mapped into a positive operator. In fact,

$$\rho_T(0) = \sum_{i,j} c_{ij} \rho_S^i(0) \otimes \rho_{S_n}^j(0) \quad (1.14)$$

with $c_{ij} > 0$, so that, since $\Lambda_t \geq 0$,

$$\rho_T(t) = (\Lambda_t \otimes \mathbf{I}_n)[\rho_T(0)] = \sum_{i,j} c_{ij} \Lambda_t[\rho_S^i(0)] \otimes \rho_{S_n}^j(0) \geq 0. \quad (1.15)$$

Then it is the presence of quantum entanglement that asks for complete positivity, that is only entangled states of $S + S_n$ can lose their positivity under the dynamical map $\Lambda_t \otimes \mathbf{I}_n$ when Λ_t is not completely positive. We shall shortly come back to this point after discussing the structural characterization of completely positive maps.

The following theorem is due to Kraus [16] who used a result by Stinespring [17], obtained in the general framework of completely positive maps on C^* -algebras.

Theorem 1.1 (Kraus, Stinespring) A linear transformation $\Gamma : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ is CP iff $\Gamma(B) = \sum_{k \in K} A_k^\dagger B A_k$, $\forall B \in \mathcal{B}(\mathfrak{H})$, where K is a finite or countably infinite set; the operators $A_k \in \mathcal{B}(\mathfrak{H})$ satisfy $\sum_{k \in K} A_k^\dagger A_k \leq \mathbf{I}$. The set $\{A_k, k \in K\}$ is not uniquely determined by Γ .

Corollary 1.1 A dynamical map $\Lambda_t : \mathcal{P}(\mathfrak{H}) \rightarrow \mathcal{P}(\mathfrak{H})$ is CP iff $\Lambda_t[\rho] = \sum_{k \in K} A_k \rho A_k^\dagger$, $\forall \rho \in \mathcal{P}(\mathfrak{H})$, with $\sum_{k \in K} A_k^\dagger A_k \leq \mathbf{I}$.

Corollary 1.2 For a trace-preserving completely positive map the operators A_k in Theorem 1.1 must satisfy $\sum_{k \in K} A_k^\dagger A_k = \mathbf{I}$.

When $\mathfrak{H}_S$ is finite-dimensional, say $\dim(\mathfrak{H}_S) = d$, one doesn't need to check all the maps $\Lambda_t \otimes \mathbf{I}_n$, but can use the following result.

Theorem 1.2 (Choi [18,19]) Given a finite d -dimensional Hilbert space $\mathfrak{H}_d$, a linear transformation $\Gamma : \mathcal{B}(\mathfrak{H}_d) \rightarrow \mathcal{B}(\mathfrak{H}_d)$ is CP iff it is d -positive.

Remark 1.6 Note that the standard evolutions for systems, either stable or unstable (equations (1.5) and (1.11)), are in the form of Corollary 1.1, thus they are completely positive.

Remark 1.7 A simple example of a positive but not completely positive map is given by the transposition in a two-dimensional Hilbert space, $\tau : M_2(\mathbf{C}) \rightarrow M_2(\mathbf{C})$. Positivity holds for $\tau[\rho] \geq 0$, $\forall \rho \in \mathcal{P}(\mathcal{H}_2)$; to see that complete positivity does not hold, consider the singlet state $\rho = |\psi\rangle\langle\psi|$ where $|\psi\rangle = (|e_1\rangle \otimes |e_2\rangle - |e_2\rangle \otimes |e_1\rangle)/\sqrt{2}$. ρ is represented by the 4×4 positive matrix:

$$\rho = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (1.16)$$

After explicit computation, we find that $\rho_\tau \equiv \tau \otimes \mathbf{I}_2$ is given by

$$\rho_\tau = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \quad (1.17)$$

that is no longer positive, since one of its eigenvalues is $-1/2$. Consequently, τ is not completely positive.

Using Theorem 1.2, we stress the tight connection between complete positivity and quantum entanglement.

Proposition 1.1 If Λ_t is not completely positive, there exists an entangled state ρ_T in $S + S_d$ whose positivity is not preserved by $\Lambda_t \otimes \mathbf{I}_d$.

Proof: If Λ_t is not completely positive, the same holds for its dual Λ_t^* . Then, because of Theorem 1.2, there must be a positive bounded operator $A \in M_d(\mathbf{C}) \otimes M_d(\mathbf{C})$ such that $(\Lambda_t^* \otimes \mathbf{I}_d)[A]$ has a negative eigenvalue. Let ρ_T be the corresponding eigenprojection, then

$$\text{Tr} [\rho_T (\Lambda_t^* \otimes \mathbf{I}_d)[A]] = \text{Tr} [(\Lambda_t \otimes \mathbf{I}_d)[\rho_T] A] \leq 0. \quad (1.18)$$

Thus, $(\Lambda_t \otimes \mathbf{I}_d)[\rho_T]$ must have negative eigenvalues since $A \geq 0$.

To summarize, positivity of a dynamical map Λ_t guarantees that the eigenvalues of any state ρ_S evolving under Λ_t remain positive at any time. Complete positivity of the dynamical map Λ_t guarantees that the eigenvalues of any entangled state ρ_T of $S + S_n$ remain positive at any time, where S_n is an arbitrary n -level system, possibly very far from S , and the coupling between S and S_n is statistical, not dynamical.

Because of the abstractness of the definition of CP, the necessity of positivity of maps of the form $\Lambda_t \otimes \mathbf{I}_n$, and thus of complete positivity of Λ_t , is often refused as a technical artifact rather than a physical constraint.

In fact, the motivations for considering the system of interest S jointly with an arbitrary n -level system S_n , not evolving and without interaction with S , are not obvious. All the more so, since this coupling should motivate the special form that, according to Corollary 1.1, the time-evolution would inherit.

In Section 1.6 we will show that complete positivity of Λ_t is a necessary and sufficient condition to preserve the positivity of factorized maps in the form $\Lambda_t \otimes \Lambda_t$. This gives the opportunity to study the notion of CP in the treatment of the dissipative evolution of entangled pairs of neutral mesons, given in Chapter 2. More in general, time-evolutions of the form $\Lambda_t \otimes \Lambda_t$ are definitely physically plausible and less abstract than $\Lambda_t \otimes \mathbf{I}_n$.

Nevertheless, in Chapter 5 we shall deal with a physical setup reproducing a factorized dynamics in the form $\Lambda_t \otimes \mathbf{I}_n$ and involving entanglement between the degrees of freedom of two subsystems: then, it will be possible to show explicitly, even with $\Lambda_t \otimes \mathbf{I}_n$, that lack of complete positivity of Λ_t produces necessarily physically concrete inconsistencies.

1.3 Reduced dynamics: the Generalized Master Equation

In the previous sections we have described the general features characterizing the time-evolutions of quantum systems $\rho \in \mathcal{P}(\mathcal{H})$. Now, we proceed to describe the time-evolution equations for open systems.

We denote the system of interest by S , the environment by B and assume that the Hamiltonian H_T of the compound system $T = S + B$ is given. Such a Hamiltonian consists of contributions H_S and H_B from system and environment alone, plus an interaction term H_I :

$$H_T = H_S + H_B + H_I. \quad (1.19)$$

H_I is usually assumed to be of the linear form

$$H_I = \sum_j V_S^j \otimes V_B^j, \quad (1.20)$$

with V_S^j, V_B^j Hermitian operators of S , respectively B .

We want to describe states and observables belonging to S only (because of lack of control over all the degrees of freedom of B). The formalism we follow in order to obtain the dynamics of S when the dynamics of T is known, consists in the projector technique developed independently by Nakajima [21] and Zwanzig [22] (see [23] for a review). There are also different approaches [24, 25, 26, 27, 28, 29, 30] to describe the time-evolution of open systems; in Section 1.7 we shall present a stochastic approach [125].

The Hilbert spaces characterizing S , B and T are such that $\mathfrak{H}_T = \mathfrak{H}_S \otimes \mathfrak{H}_B$. We are interested in observables $A \in \mathcal{B}(\mathfrak{H}_S)$, whose expectation values are:

$$\langle A \rangle_S = \text{Tr}(A \rho_T) = \text{Tr}_S \text{Tr}_B[(A \otimes \mathbf{I}_B) \rho_T] = \text{Tr}_S(A \rho_S), \quad (1.21)$$

where the statistical operator of S is obtained from the one of T by the partial trace over the degrees of freedom of B :

$$\rho_S = \text{Tr}_B(\rho_T). \quad (1.22)$$

We are interested in reduced dynamics of the system S alone, that is in linear maps Λ_t describing the time-evolution $\rho_S(0) \rightarrow \Lambda_t[\rho_S(0)]$. In full generality, the state of S at time t , denoted by $\rho_S(t)$, is obtained from the unitary evolution of $T = S + B$, $U_t = e^{-iH_T t}$, as follows:

$$\rho_S(t) = \text{Tr}_B[U_t \rho_T(0) U_t^\dagger]. \quad (1.23)$$

However, $\rho_S(t)$ is not always expressible as $\Lambda_t[\rho_S(0)]$, because there is no general prescription for extracting $\rho_S(0)$ from the initial global state $\rho_T(0)$. Indeed, there can be two different initial global states $\rho_T^a(0) \neq \rho_T^b(0)$ that lead to the same reduced state of S , $\rho_S^a(0) = \rho_S^b(0)$, and nevertheless at time t $\rho_S^a(t) \neq \rho_S^b(t)$. In this case, the dynamics Λ_t is not properly defined because a state of S at time $t = 0$ would evolve in two distinct states at time t . It is thus necessary to define an assignment map $\Phi : \mathcal{P}(\mathfrak{H}_S) \rightarrow \mathcal{P}(\mathfrak{H}_T)$ giving, for each initial state $\rho_S(0)$, a state of the total system T such that

$$\rho_S(t) = \Lambda_t[\rho_S(0)] = \text{Tr}_B(U_t \Phi[\rho_S(0)] U_t^\dagger), \quad (1.24)$$

and the diagram in Figure 1.3 commute. Following this procedure, it is possible to define the reduced dynamics Λ_t . The conditions usually imposed on the map Φ are: 1. Φ preserves mixtures: $\Phi[\sum_i \lambda_i \rho_S^i] = \sum_i \lambda_i \Phi[\rho_S^i]$; 2. Φ is consistent: $\text{Tr}_B(\Phi[\rho_S]) = \rho_S$; 3. $\Phi[\rho_S]$ is positive for any positive ρ_S . It has been shown that, for a two-dimensional Hilbert space $\mathfrak{H}_S$, the only assignment maps satisfying these three conditions for all initial density

$$\begin{array}{ccc}
\rho_S(0) \in \mathcal{P}(\mathcal{H}_S) & \xrightarrow{\Lambda_t} & \rho_S(t) \in \mathcal{P}(\mathcal{H}_S) \\
\downarrow \Phi & & \uparrow \text{Tr}_B \\
\Phi[\rho_S(0)] \in \mathcal{P}(\mathcal{H}_T) & \longrightarrow & U_t \Phi[\rho_S(0)] U_t^\dagger \in \mathcal{P}(\mathcal{H}_T)
\end{array}$$

Figure 1.1: The procedure required in order to define consistently the reduced dynamics Λ_t acting on the states of the system S using the projector technique.

matrices are the product maps $\Phi[\rho_S] = \rho_S \otimes \rho_B$, where ρ_B is a fixed reference state in the bath [24]. This factorized state is usually justified in the context of a weak coupling between S and B ; however there are physical situations in which such a factorized initial state cannot be used [25].

Remark 1.8 Since Λ_t in (1.24) is obtained by the composition of three maps, its properties depends on those of the trace over the environment degrees of freedom, of the standard evolution and of the assignment map Φ . If the choice $\Phi[\rho_S] = \rho_S \otimes \rho_B$ is assumed, the reduced dynamics Λ_t is completely positive since it is the composition of maps that are CP. However, in some cases the initial state has not the form of a product state, namely, when there exist initial correlations between S and B [24, 25]. In [24] an argument against the complete positivity of a reduced dynamics has been discussed. It has been argued that, with correlated initial conditions, dynamical maps Λ_t can be properly defined only on a subset of initial system states and the linear extension of these maps on all possible initial states may not be physically realizable and may not even be positive. In this case the condition 3 is not fulfilled; an example of this procedure is given by the assignment map proposed in [26]. In literature many examples of not completely positive dynamical maps are encountered [27, 28, 29, 30, 31]; however, in [32] it has been shown that it is possible in any case to preserve the complete positivity of Λ_t , giving up either the condition 1 or 2; it has also been stressed that, beyond the weak coupling regime, there exists no unique definition of the reduced dynamics Λ_t for the system S .

Let us now present the Nakajima-Zwanzig procedure. We assume the dynamics of the total system T to be given; furthermore, to be as general as possible, we do not consider the latter to be necessarily closed. Namely, we consider, for the states ρ_T of T , time-evolution equations of the form

$$\dot{\rho}_T(t) = L_T[\rho_T(t)], \quad (1.25)$$

where $L_T = L_S + L_B + L_I$ is the generator of the time-evolution of ρ_T ; only when T is a closed system, L_T is proportional to the commutator with the total Hamiltonian $H_T = H_S + H_B + H_I$, see equation (1.6).

We define the two projectors $P, Q : \mathcal{P}(\mathfrak{H}_T) \rightarrow \mathcal{P}(\mathfrak{H}_T)$ that map states of T into themselves as follows:

$$P[\rho_T] \equiv \text{Tr}_B(\rho_T) \otimes \rho_B = \rho_S \otimes \rho_B; \quad Q = \mathbf{I} - P. \quad (1.26)$$

They depend on a normalized reference state ρ_B of the bath, $\text{Tr}_B(\rho_B) = 1$, that one usually choose to be invariant under the dynamics of B : $L_B[\rho_B] = 0$. The following manipulations are independent from the choice of this state. Inserting in equation (1.25) $\rho_T(t) = P[\rho_T(t)] + Q[\rho_T(t)]$, we obtain

$$\begin{cases} P[\dot{\rho}_T(t)] = PL_TP[\rho_T(t)] + PL_TQ[\rho_T(t)], \\ Q[\dot{\rho}_T(t)] = QL_TP[\rho_T(t)] + QL_TQ[\rho_T(t)]. \end{cases} \quad (1.27)$$

Formally, solving the second of these equations, one gets

$$Q[\rho_T(t)] = e^{QL_Tt}Q[\rho_T(0)] + \int_0^t dt' e^{QL_Tt'} QL_TP[\rho_T(t-t')]. \quad (1.28)$$

Then upon inserting (1.28) in the first equation in (1.27), we find a closed equation for the relevant part $P[\rho_T(t)]$; after a partial trace over B we are left with the equation of motion of S alone, a so-called generalized master equation (GME):

$$\dot{\rho}_S(t) = L_S^{eff}[\rho_S(t)] + \int_0^t dt' \mathcal{K}(t')[\rho_S(t-t')] + \mathcal{I}(t) \quad (1.29)$$

with

$$\begin{aligned} L_S^{eff} &\equiv L_S + \text{Tr}_B(L_I[\rho_B]); \\ \mathcal{K}(t)[\rho_S] &\equiv \text{Tr}_B\left(L_I e^{QL_Tt} QL_T[\rho_S \otimes \rho_B]\right); \\ \mathcal{I}(t) &\equiv \text{Tr}_B\left(L_I e^{QL_Tt} Q[\rho_T(0)]\right). \end{aligned} \quad (1.30)$$

These expressions have been obtained using that $PL_{S,B}Q = 0$.

The GME is the equation we were looking for; it describes in general the behavior of an open system S because no assumptions have been made about the nature of the system and of the bath, and the interaction is completely arbitrary; we have only assumed the existence of a reference bath-equilibrium state ρ_B in $\mathcal{P}(\mathfrak{H}_B)$. Notice that in absence of interaction between S and B (i.e. if $L_I = 0$) this equation reduces to the time-evolution for S isolated from B , that is $\dot{\rho}_s(t) = L_S[\rho_S(t)]$; in fact $L_S^{eff} = L_S$, $\mathcal{K}(t) = 0$ and $\mathcal{I}(t) = 0$.

Let us now describe the various terms appearing in the GME. L_S^{eff} generates the effective time-evolution in ρ_S , redefined by $\text{Tr}_B(L_I[\rho_B])$ which represents a mean field term. $\mathcal{K}(t)$ is the kernel of the convolution term, that is the memory term depending on the whole history of ρ_S . The inhomogeneity $\mathcal{I}(t)$ is related to initial conditions: if they

are factorized, $\rho_T(0) = \rho_S(0) \otimes \rho_B$, then $Q\rho_T(0) = 0$ and consequently $\mathcal{I}(t) = 0$. In the following we will assume this condition to be fulfilled.

The reduced dynamics Λ_t is well defined and it is completely positive.

If T is a closed system it is possible to simplify (1.29) and (1.30). In such a case the operators L are proportional to the commutators with the Hamiltonians and the following relations hold: $[P, L_S] = 0$; $PL_B = L_BP = 0$. Then the GME becomes:

$$\begin{aligned}\dot{\rho}_S(t) &= L_S^{eff}[\rho_S(t)] + \int_0^t dt' \mathcal{K}(t')[\rho_S(t-t')]; \\ L_S^{eff}[\rho_S] &= -i[H_S + \text{Tr}_B(H_I \rho_B), \rho_S], \\ \mathcal{K}(t)[\rho_S] &= \text{Tr}_B \left(L_I e^{Q L_T Q t} Q L_I [\rho_S \otimes \rho_B] \right).\end{aligned}\tag{1.31}$$

Let us focus our attention on cases in which the interaction between S and B is weak. We introduce a small dimensionless parameter ε in the interaction Hamiltonian; consequently $H_T = H_S + H_B + \varepsilon H_I$. In this way we can write a perturbative expansion in ε for the kernel $\mathcal{K}(t)$ in equation (1.31) [40]:

$$\mathcal{K}(t) = e^{L_S t} \left[\mathcal{K}_0(t) + \sum_{n=1}^{\infty} \varepsilon^n \int dt_1 \dots dt_n \mathcal{K}_n(t|t_1 \dots t_n) \right],\tag{1.32}$$

with $0 \leq t_n \leq \dots \leq t_1 \leq t$, and

$$\begin{aligned}\mathcal{K}_0(t)[\rho_S] &= \text{Tr}_B(L_I^i(t) Q L_I^i[\rho_S \otimes \rho_B]), \\ \mathcal{K}_n(t|t_1 \dots t_n)[\rho_S] &= \text{Tr}_B(L_I^i(t) Q L_I^i(t_1) Q \dots Q L_I^i(t_n) Q L_I^i[\rho_S \otimes \rho_B])\end{aligned}\tag{1.33}$$

where

$$L_I^i(t_j)[\cdot] = -i[H_I^i(t_j), \cdot] = -i[e^{i(H_S+H_B)t_j} H_I e^{-i(H_S+H_B)t_j}, \cdot]\tag{1.34}$$

is the Liouville generator in the interaction picture. $\mathcal{K}_n(t|t_1 \dots t_n)$ contains the multi-time correlation function of the bath

$$h_{j,j_1 \dots j_{n+1}}(t, t_1, \dots t_n) = \text{Tr}_B(e^{iH_B t} V_B^j e^{-iH_B t} e^{iH_B t_1} V_B^{j_1} e^{-iH_B t_1} \dots V_B^{j_{n+1}} \rho_B),\tag{1.35}$$

where, as already mentioned in (1.20), we have assumed the interaction Hamiltonian to be $H_I = \sum_j V_S^j \otimes V_B^j$.

Usually, one proceeds with a Born approximation of the GME, namely with neglecting in the kernel expansion (1.32) all the terms except $\mathcal{K}_0(t)$; in this case only two-time correlation functions appear.

1.4 The Markovian approximation. The Weak Coupling Limit

As we have seen, the GME obtained in the previous section, eq. (1.31) (or, more in general, eq. (1.29)) contains a memory term that depends on the kernel $\mathcal{K}(t)$. If $\mathcal{K}(t)$ decays rather fastly, it is possible to extract the density matrix from the convolution integral, then obtaining an equation without memory term [1, 33, 34]; in such a case it is possible to introduce a time-independent generator of the reduced dynamics.

Definition 1.10 The dynamical map $\{\Lambda_t, t \in \mathfrak{R}\}$ is said to be *Markovian* iff there exists a linear bounded operator L on the states space $\mathcal{P}(\mathcal{H})$ which is time-independent, such that $\Lambda_t = e^{Lt} \ \forall t \in \mathfrak{R}$; L is the generator of the dynamical map.

We want to extract a Markovian time-evolution from the GME presented in the previous section: in this way we obtain a more manageable time-evolution for the system considered.

Remark 1.9 A Markovian time-evolution $\{\Lambda_t, t \in \mathfrak{R}\}$ satisfies the semigroup composition law $\Lambda_{t_1+t_2} = \Lambda_{t_1} \circ \Lambda_{t_2}$, $\forall t_1, t_2 \in \mathfrak{R}$, as can be seen from Definition 1.10.

As already stated, in order to perform a Markovian approximation of the dynamics (1.29) or (1.31) we need the decay time of the kernel $\mathcal{K}(t)$ to be much shorter than the typical evolution time of the system S . We find this situation when the interaction Hamiltonian can be parametrized by a suitable coefficient ε which is then let to vanish: $H_I \rightarrow \varepsilon H_I$. The total Hamiltonian is thus scaled as $H_T \rightarrow H_T^\varepsilon$:

$$H_T^\varepsilon = H_S + H_B + \varepsilon H_I, \quad (1.36)$$

and the corresponding dynamical map is denoted by Λ_t^ε :

$$\Lambda_t^\varepsilon[\rho_S(0)] = \text{Tr}_B(U_t^\varepsilon \rho_S(0) \otimes \rho_B U_t^{\varepsilon\dagger}), \quad (1.37)$$

where $U_t^\varepsilon = e^{-iH_T^\varepsilon t}$.

As we shall shortly see, for small ε the reduced dynamics Λ_t^ε becomes Markovian [35] over times scaling as ε^{-2} : this procedure is called Weak Coupling Limit.

Since the memory term is of order ε^2 (being quadratic in the interaction Hamiltonian), the evolution has to be studied over a time-scale of order ε^{-2} . It is then appropriate to rescale the time as $t^\varepsilon = \varepsilon^{-2}t$; in other words, the evolution is observed over a time long enough so that the memory effects can be neglected. On this time scale the convolution term in the GME heuristically behaves as:

$$\int_0^{t^\varepsilon} \mathcal{K}(t')[\rho_S(t^\varepsilon - t')]dt' \approx \left(\int_0^{+\infty} \mathcal{K}(t')dt' \right) [\rho_S(t^\varepsilon)] \equiv K[\rho_S(t^\varepsilon)] \quad (1.38)$$

and thus the GME becomes

$$\dot{\rho}_S(t) = L_S^{eff}[\rho_S(t)] + \varepsilon^2 \int_0^t \mathcal{K}(t')[\rho_S(t-t')]dt' \approx (L_S^{eff} + \varepsilon^2 K)[\rho_S(t)], \quad (1.39)$$

whose solution is

$$\rho_S(t) = \Lambda_t^\varepsilon[\rho_S(0)] = e^{(L_S^{eff} + \varepsilon^2 K)t}[\rho_S(0)]. \quad (1.40)$$

Thus $L = L_S^{eff} + \varepsilon^2 K$ is the time-independent generator, and the time-evolution Λ_t^ε is Markovian. This is the basic idea underlying any Markovian approximation of the generalized master equation; there are however many ways to perform the Markov limit, leading to Markovian reduced dynamics with different properties [35].

Remark 1.10 The prescription (1.38) generally breaks the complete positivity and even positivity of the reduced dynamics [35]

The convergence condition of the weak coupling limit depends on the time decay of the correlations of the bath. If we suppose $H_I = \sum_j V_S^j \otimes V_B^j$ this condition reads

$$\int_0^\infty dt |h_{jk}(t)|(1+t)^\delta < \infty, \quad \forall \delta > 0, \quad (1.41)$$

where

$$h_{jk}(t) = \text{Tr}_B \left(e^{iH_B t} V_B^j e^{-iH_B t} V_B^k \rho_B \right). \quad (1.42)$$

The condition (1.41) can be fulfilled only if the reservoir B is infinite [33, 34].

The Markovian approximations of the GME usually does not preserve the positivity of the states over which they act [35]. The following theorem defines Markovian approximations that do not suffer from lack of positivity. Noticeably, they turn out to be not only positive but also completely positive.

Theorem 1.3 (Davies [36]) Suppose the condition (1.41) is valid and the spectrum of H_S is discrete, then

$$\lim_{\varepsilon \rightarrow 0} \sup_{0 \leq \varepsilon^2 t \leq \tau} \|\Lambda_t^\varepsilon[\rho_S(0)] - e^{(L_S^{eff} + \varepsilon^2 K^\natural)t}[\rho_S(0)]\|_1 = 0, \quad (1.43)$$

where $\|\cdot\|_1$ is the norm introduced in Definition 1.2 and

$$K^\natural[\rho_S] \equiv \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T dt e^{-L_S t} K e^{L_S t}[\rho_S] \quad (1.44)$$

with

$$K[\rho_S] \equiv \int_0^\infty dt' e^{-L_S t'} \text{Tr}_B(L_I e^{(L_S + L_B)t'} L_I[\rho_B])[\rho_S]. \quad (1.45)$$

The Markovian evolution defined by (1.43) is completely positive.

In (1.43) we have to consider the averaged evolver $K^\sharp$ rather than K since the free motion and the dissipation have different time scales.

Another kind of Markovian approximation is the so called Singular Coupling Limit (or Singular Reservoir Limit), introduced in [37]; in this case we do not rescale the Hamiltonian H_T but the two-time correlation functions of the environment: $h_{jk}(t) \rightarrow h_{jk}^\varepsilon(t)$, with $h_{jk}^\varepsilon \rightarrow c_{jk}\delta(t)$ as $\varepsilon \rightarrow 0$. It has been shown in [38] that such a case is equivalent to a weak coupling limit with a rescaled Hamiltonian in the form

$$H_T^\varepsilon = \varepsilon^2 H_S + H_B + \varepsilon H_I, \quad (1.46)$$

where the free Hamiltonian H_S is scaled by a factor ε^2 and then the uncoupled and dissipative evolution are of the same order.

Remark 1.11 The singular coupling limit is particularly relevant in the study of Markovian quantum open systems. Indeed, it is possible to show that any family of completely positive maps satisfying the semigroup composition law, describing the dynamics of a n -level system, can be derived by a singular coupling [39].

Remark 1.12 Another relevant property of the singular coupling is that, in this limit, the correlation functions of order higher than 2 vanish, that is the generalized master equation reduces to its Born approximation, and it is Markovian [40].

Historically the Markovian approximations have been studied considering two typical times, τ_S and τ_B , characterizing the system and the bath respectively; the density matrix ρ_S can be extracted from the integral kernel $\mathcal{K}(t)$ under the hypothesis $\tau_S/\tau_B \rightarrow \infty$. There are two possibilities to realize this condition: either $\tau_S \rightarrow \infty$ and τ_B remains constant, or $\tau_B \rightarrow 0$ while τ_S remains constant [34]. The two alternatives correspond to weak coupling and singular coupling respectively.

1.5 Quantum Dynamical Semigroups

In the previous section we have introduced the set of dynamical maps we are interested in, namely, the set of Markovian time-evolutions describing the irreversible evolution of an open system under the influence of an external environment. It turns out that the structure of the generators of these dynamical maps is fully characterized by two theorems that we shall present after introducing the notion of quantum dynamical semigroups.

Definition 1.11 A set of time-evolutions $\{\Lambda_t, t \geq 0\}$ forms a *quantum dynamical semigroup* [1, 33] iff the following properties are fulfilled:

- 1.i) Λ_t is completely positive $\forall t \geq 0$;
- 2.i) Λ_t is trace-preserving: $\text{Tr}(\Lambda_t[\rho]) = \text{Tr}(\rho) \quad \forall t \geq 0, \forall \rho \in \mathcal{P}(\mathfrak{H})$;
- 3.i) Λ_t satisfies the forward in time composition law: $\Lambda_{t_1+t_2}[\rho] = \Lambda_{t_1} \circ \Lambda_{t_2}[\rho] \quad \forall t_1, t_2 \geq 0, \forall \rho \in \mathcal{P}(\mathfrak{H})$;
- 4.i) the Λ_t are strongly continuous: $\lim_{t \rightarrow 0} \|\Lambda_t[\rho] - \rho\|_1 = 0 \quad \forall \rho \in \mathcal{P}(\mathfrak{H})$.

In the Heisenberg picture, defined by (1.4), the corresponding properties of the dual map Λ_t^* are:

- 1.ii) Λ_t^* is completely positive $\forall t \geq 0$;
- 2.ii) $\Lambda_t^* \mathbf{I} = \mathbf{I}$, where $\mathbf{I}$ is the identity on $\mathcal{B}(\mathfrak{H})$;
- 3.ii) Λ_t^* satisfies the forward in time composition law: $\Lambda_{t_1+t_2}^*[A] = \Lambda_{t_1}^* \circ \Lambda_{t_2}^*[A] \quad \forall t_1, t_2 \geq 0, \forall A \in \mathcal{B}(\mathfrak{H})$;
- 4.ii) Λ_t^* ultraweakly continuous: $\lim_{t \rightarrow 0} \text{Tr}[\rho(\Lambda_t^*[A] - A)] = 0 \quad \forall \rho \in \mathcal{P}(\mathfrak{H}), \forall A \in \mathcal{B}(\mathfrak{H})$;
- 5.ii) Λ_t^* is normal.

Remark 1.13 The properties 3.i) and 4.i) imply that $\Lambda_t = e^{Lt}$, where L is the generator of the semigroup (and $\dot{\rho}(t) = L[\rho(t)]$); then the maps $\{\Lambda_t, t \geq 0\}$ are Markovian. Analogously, $\Lambda_t^* = e^{L^*t}$ (i.e. $\dot{A}(t) = L^*[A(t)]$) and L^* is the generator of the Markovian semigroup $\{\Lambda_t^*, t \geq 0\}$.

Remark 1.14 The property 5.ii) guarantees that Λ_t^* is the dual map of Λ_t [33].

The form of the generator of a quantum dynamical semigroup, if it is bounded, as a linear operator on the space of states $\mathcal{P}(\mathcal{H})$, is completely determined as follows.

Theorem 1.4 (Lindblad [41]) Given a generic separable Hilbert space $\mathfrak{H}$, a bounded linear transformation $L : \mathcal{P}(\mathfrak{H}) \rightarrow \mathcal{P}(\mathfrak{H})$ is the generator of a quantum dynamical semigroup on $\mathcal{P}(\mathfrak{H})$ iff

$$L[\rho] = -i[H, \rho] + \frac{1}{2} \sum_j \left([V_j \rho, V_j^\dagger] + [V_j, \rho V_j^\dagger] \right), \quad (1.47)$$

where $H = H^\dagger$ and the $V_j \in \mathcal{B}(\mathcal{H})$ satisfy $\sum_j V_j^\dagger V_j \in \mathcal{B}(\mathfrak{H})$. The dual generator L^* , in the Heisenberg picture, then reads

$$L^*[A] = i[H, A] + \frac{1}{2} \sum_j \left(V_j^\dagger [A, V_j] + [V_j^\dagger, A] V_j \right). \quad (1.48)$$

Independently, Gorini *et al.* arrived at (1.47) in the case of finite dimensional system S .

Theorem 1.5 (Gorini, Kossakowski, Sudarshan [42]) Given a finite n -dimensional Hilbert space, $\mathfrak{H}_n$, a bounded operator $L : \mathcal{P}(\mathfrak{H}_n) \rightarrow \mathcal{P}(\mathfrak{H}_n)$ is the generator of a quantum dynamical semigroup on $\mathcal{P}(\mathfrak{H}_n)$ iff

$$L[\rho] = -i[H, \rho] + \frac{1}{2} \sum_{k,l=1}^{n^2-1} c_{kl} \left([F_k \rho, F_l^\dagger] + [F_k, \rho F_l^\dagger] \right), \quad (1.49)$$

with $H = H^\dagger$, $\{F_k, k = 0, \dots, n^2 - 1\}$ a basis in $M_n(\mathbf{C})$ (the space of $n \times n$ matrices with complex entries), $F_0 = \mathbf{I}_n / \sqrt{n}$, $\text{Tr}(F_k) = 0 \ \forall k \neq 0$, $\text{Tr}(F_i^\dagger F_j) = \delta_{ij}$ and $C = [c_{kl}]$ a positive matrix (whose entries depend on the choice of the basis $\{F_k\}$). The Heisenberg generator $L^* : \mathcal{B}(\mathfrak{H}_n) \rightarrow \mathcal{B}(\mathfrak{H}_n)$ is written as

$$L^*[A] = i[H, A] + \frac{1}{2} \sum_{k,l=1}^{n^2-1} c_{kl} \left(F_l^\dagger [A, F_k] + [F_l^\dagger, A] F_k \right), \quad [c_{kl}] \geq 0. \quad (1.50)$$

Corollary 1.3 Given a n -dimensional Hilbert space, $\mathfrak{H}_n$, with n finite, a bounded operator $L : \mathcal{P}(\mathfrak{H}_n) \rightarrow \mathcal{P}(\mathfrak{H}_n)$ is the generator of a trace and Hermiticity-preserving semigroup of maps iff it has the form in (1.49), with $H = H^\dagger$ the effective Hamiltonian and $C = [c_{kl}]$ a Hermitian matrix.

Remark 1.15 The form of the generator of a generic semigroup does not change if we relax the request of complete positivity, but the positivity of C is lost.

The entries of the matrix C are not all independent because of the Hermiticity (or positivity) condition. A relation between these entries we shall use in the following is expressed by Lemma 1.1.

Lemma 1.1 The entries of the matrix C in the generator L of a trace and Hermiticity-preserving semigroup of maps acting on $\mathcal{P}(\mathfrak{H})$, eq. (1.49), satisfy the constraints

$$|c_{kl}| \leq \frac{1}{2}(c_{kk} + c_{ll}), \quad k, l = 1, \dots, n^2 - 1 \quad (1.51)$$

Proof: Expressing the operators in (1.47) as $V_j = \sum_{k=0}^{n^2-1} v_k^j F_k$ we have $c_{kl} = \sum_j v_k^j v_l^{j*}$. Then, applying Schwartz inequality,

$$|c_{kl}| \leq \sqrt{c_{kk} c_{ll}} \leq \frac{1}{2}(c_{kk} + c_{ll}).$$

Remark 1.16 By rearranging equations (1.47) and (1.49), the generator L can also be written as

$$L[\rho] = -i[H, \rho] + \sum_j \left(V_j \rho V_j^\dagger - \frac{1}{2} \{ \rho, V_j^\dagger V_j \} \right) \quad (1.52a)$$

$$L[\rho] = -i[H, \rho] + \sum_{k,l=1}^{n^2-1} c_{kl} \left(F_k \rho F_l^\dagger - \frac{1}{2} \{ \rho, F_l^\dagger F_k \} \right); \quad (1.52b)$$

all these expressions for L are generically called Lindblad forms.

Remark 1.17 The Hamiltonian H is not the free Hamiltonian H_S of the system of interest; indeed, in general it contains dissipative terms produced by the interaction with the environment: contributions of first order in this interaction, the mean field term $\text{Tr}_B(L_I[\rho_B])$ in (1.30), and of second order, as we shall see explicitly in Chapter 4.

Let us now analyze separately the contributions that appear in (1.52a).

The commutator with the Hamiltonian H generates a time-evolution trace and coherence preserving. The anticommutator between ρ and the positive operator $R \equiv \sum_j V_j^\dagger V_j$ may be interpreted as a friction term; indeed, let us consider the time-evolution produced only by the anticommutator and the commutator:

$$\begin{aligned} \dot{\rho}(t) &= -i[H, \rho] - \frac{1}{2} \{ R, \rho \} = -iK\rho(t) + i\rho(t)K^\dagger, \\ \rho(t) &= e^{-iKt} \rho(0) e^{iK^\dagger t}, \end{aligned} \quad (1.53)$$

where $K = H - \frac{i}{2}R$. This evolution has the Weisskopf-Wigner form and it is typical of an unstable system, as previously remarked; in fact it causes a loss of probability, as we have seen in Section 1.1:

$$\frac{d}{dt} \text{Tr}[\rho(t)] = \text{Tr}[\dot{\rho}(t)] = -\text{Tr}[\rho(t)R] \leq 0 \quad \forall t \geq 0.$$

Notice that, despite loss of probability, the time-evolution (1.53) preserves the coherence of states. In fact, in order to get decoherence, the term $\sum_j V_j \rho V_j^\dagger$ in (1.52a), called noise, is needed. Starting with a pure state, $\rho = |\psi\rangle\langle\psi|$,

$$\begin{aligned} \frac{d}{dt} \text{Tr}[\rho(t)^2] |_{t=0} &= 2 \text{Tr}[\dot{\rho}(0)\rho(0)] = 2 \sum_j \text{Tr}[V_j \rho(0) V_j^\dagger - V_j^\dagger V_j \rho(0)] \\ &= -2 \sum_j (\langle \psi | V_j^\dagger V_j | \psi \rangle - \langle \psi | V_j | \psi \rangle \langle \psi | V_j^\dagger | \psi \rangle) \leq 0 \end{aligned} \quad (1.54)$$

Then, in general $\rho(t)^2 \neq \rho(t)$, so that pure states become mixtures. Moreover, the noise term balances the probability loss due to pure friction and the overall evolution is probability preserving:

$$\frac{d}{dt} \text{Tr}[\rho(t)] = \sum_j \text{Tr} \left[V_j \rho(t) V_j^\dagger - \frac{1}{2} \{V_j^\dagger V_j, \rho(t)\} \right] = 0, \quad \forall t \geq 0. \quad (1.55)$$

1.6 Factorized dynamics of quantum dynamical semigroups

In this section we consider two systems of the same kind, S_1 and S_2 , in weak interaction with a same environment. We further suppose that the composite system $S_1 + S_2$ evolves in time according to a reduced dynamics of factorized form $\Lambda_t \otimes \Lambda_t$, where Λ_t is the dynamics of either systems S_1 or S_2 and $\{\Lambda_t, t \geq 0\}$ is assumed to form a Markovian semigroup so that the same is true for the set of maps $\{\Lambda_t \otimes \Lambda_t, t \geq 0\}$. We focus our attention on the positivity properties of these maps; in particular, in Theorem 1.6 we shall prove that, in the case of S_1 and S_2 n -dimensional systems, the positivity of $\Lambda_t \otimes \Lambda_t$ necessarily requires the complete positivity of Λ_t [43].

Given a semigroup $\{\Lambda_t, t \geq 0\}$ generated by (1.49), the justification why Λ_t should be completely positive and thus the matrix $C = [c_{kl}]$ positive, is based on the fact that, otherwise, $\Lambda_t \otimes \mathbf{I}_n$ would fail to preserve the positivity of entangled states in $\mathcal{P}(\mathfrak{H}_n \otimes \mathfrak{H}_n)$ (see Proposition 1.1).

However, while the first factor in $\mathfrak{H}_n \otimes \mathfrak{H}_n$ refers to a concrete open quantum system evolving in time according to Λ_t , because of the possible interaction with a certain environment, the second factor represents a mere possibility of entanglement with anything described by a n -dimensional system and generically out of physical control.

Instead, we argue that complete positivity is necessary to avoid physical inconsistencies in compound systems consisting of two n -dimensional systems that interact with a same environment, but not among themselves, neither directly, nor indirectly, that is through the environment itself. In such a case, the two systems are expected to evolve according to semigroups of linear maps $\{\Lambda_t \otimes \Lambda_t, t \geq 0\}$, where Λ_t is the single open system dynamics obtained when only one of them is present in the environment.

A necessary request for the physical consistency of such dynamics is that the $\Lambda_t \otimes \Lambda_t$'s preserve the positivity of all separable and entangled states of the compound system, which now describe physically concrete and controllable settings.

Then for Markovian dynamics of open quantum systems the argument in favour of the necessity of complete positivity results strengthened with respect to the argument based on $\Lambda_t \otimes \mathbf{I}_n$ (see the definition of complete positivity in Section 1.2).

Theorem 1.6 If $\{\Lambda_t, t \geq 0\}$ is a semigroup of linear maps on $\mathcal{P}(\mathfrak{H}_n)$ generated as in (1.49), then the semigroup $\{\Lambda_t \otimes \Lambda_t, t \geq 0\}$ of linear maps over $\mathcal{P}(\mathfrak{H}_n \otimes \mathfrak{H}_n)$ is positivity-preserving if and only if $\{\Lambda_t, t \geq 0\}$ consists of completely positive maps [43].

The proof of Theorem 1.6 is divided into several steps. We need just show the only if part; indeed, if Λ_t is completely positive, $\Lambda_t \otimes \mathbf{I}_n$ and $\mathbf{I}_n \otimes \Lambda_t$ are both positive and such is the composite map $\Lambda_t \otimes \Lambda_t = (\Lambda_t \otimes \mathbf{I}_n) \circ (\mathbf{I}_n \otimes \Lambda_t)$. Actually, it is completely positive (see e.g. [14]).

Before starting with the proof of this theorem, some remarks are needed.

Remark 1.18 If the Λ_t 's preserve the positivity of states of $\mathfrak{H}_n$, $\Lambda_t \otimes \Lambda_t$ preserves the positivity of separable states of $\mathfrak{H}_n \otimes \mathfrak{H}_n$: this follows by the definition of separability (Def. 1.9).

Remark 1.19 For generic positive linear maps Λ_t on the states of $\mathfrak{H}_n$, it does not follow that, if $\Lambda_t \otimes \Lambda_t$ is positivity-preserving, then Λ_t is completely positive. A counter example is the transposition τ over $M_2(\mathbf{C})$, presented in Remark 1.7: $\tau \otimes \tau$ is positivity-preserving, but τ is not completely positive. We notice, however, that τ cannot be among the Λ_t of a continuous semigroup over the states of $M_2(\mathbf{C})$ since it is not connected to the identity map.

Remark 1.20 Actually, there are experimental situations that are describable by semigroups $\{\Lambda_t \otimes \Lambda_t, t \geq 0\}$. For instance, as we shall see in Chapter 3, neutral mesons may be imagined to suffer from dissipative effects due to a noisy background determined by Planck's scale physics. As decay products of spin 1 resonances, these mesons are produced in maximally entangled states and, while independently flying apart back to back, they arguably evolve according to semigroups $\{\Lambda_t \otimes \Lambda_t, t \geq 0\}$. In such a context, whether $\Lambda_t \otimes \Lambda_t$ is positivity-preserving is crucial for concrete physical consistency [44, 45, 46, 47, 48, 49].

The proof of Theorem 1.6 proceeds as follows.

Lemma 1.2 If $\{\Lambda_t, t \geq 0\}$ is a semigroup generated by (1.49), the set $\{\Lambda_t \otimes \Lambda_t, t \geq 0\}$ consists of positivity-preserving maps only if

$$\mathcal{L}_{\phi, \psi} \equiv \langle \phi | (L \otimes \mathbf{I}_n + \mathbf{I}_n \otimes L) [|\psi\rangle\langle\psi|] | \phi \rangle \geq 0 \quad (1.56)$$

for all orthogonal vector states $|\phi\rangle, |\psi\rangle$ in $\mathfrak{H}_n$, where L is the generator on the right hand side of (1.49) and $\mathbf{I}_n$ is the identity operation on $M_n(\mathbf{C})$.

Proof: The request of positivity preservation demands that

$$\mathcal{G}_{\phi, \psi}(t) \equiv \langle \phi | (\Lambda_t \otimes \Lambda_t) [|\psi\rangle\langle\psi|] | \phi \rangle \geq 0, \quad (1.57)$$

for all $|\phi\rangle$ and $|\psi\rangle \in \mathfrak{H}_n \otimes \mathfrak{H}_n$. Choosing $\langle \phi | \psi \rangle = 0$, if $d\mathcal{G}_{\phi, \psi}(t)/dt|_{t=0} < 0$, then $\mathcal{G}_{\phi, \psi}(t) \geq 0$ is violated in a neighborhood of $t = 0$. Thus (1.56) follows.

Lemma 1.3 In the hypothesis of Lemma 1.2, let $\{|j\rangle, j = 1, n\}$ be an orthonormal basis of $\mathfrak{H}_n$, and Φ, Ψ the $n \times n$ matrices $\Phi = [\varphi_{ij}]$, $\Psi = [\psi_{ij}]$ consisting of the coefficients of

the expansion of $|\phi\rangle$ and $|\psi\rangle$ with respect to the basis $\{|j\rangle \otimes |k\rangle; j, k = 1, n\}$ of $\mathfrak{H}_n \otimes \mathfrak{H}_n$. Then

$$\mathcal{L}_{\phi, \psi} = \sum_{a, b=1}^{n^2-1} c_{ab} \left[\text{Tr}(\Psi \Phi^\dagger F_a) \text{Tr}(\Phi \Psi^\dagger F_b^\dagger) + \text{Tr}((\Phi^\dagger \Psi)^T F_a) \text{Tr}((\Psi^\dagger \Phi)^T F_b^\dagger) \right], \quad (1.58)$$

where $C = [c_{ab}]$ is the matrix of coefficients and F_a, F_b the traceless matrices appearing in (1.49), while X^T denotes transposition of X with respect to the chosen basis.

Proof: Let $|\phi\rangle = \sum_{j,k=1}^n \varphi_{jk} |j\rangle \otimes |k\rangle$, $|\psi\rangle = \sum_{j,k=1}^n \psi_{jk} |j\rangle \otimes |k\rangle$; then, one calculates

$$\begin{aligned} \mathcal{L}_{\phi, \psi} &= \sum_{ij} \sum_{kl} \sum_{pr} (\varphi_{ij}^* \varphi_{kl} \psi_{pj} \psi_{rl}^* + \varphi_{ji}^* \varphi_{lk} \psi_{jp} \psi_{lr}^*) \langle i | L[|p\rangle \langle r|] | k \rangle \\ &= \sum_{ik} \sum_{pr} [(\Psi \Phi^\dagger)_{pi} (\Phi \Psi^\dagger)_{kr} + (\Phi^\dagger \Psi)_{ip} (\Psi^\dagger \Phi)_{rk}] \langle i | L[|p\rangle \langle r|] | k \rangle. \end{aligned} \quad (1.59)$$

The commutator and the anticommutator in the generator L drop from this equation since, given any $K \in M_n(\mathbf{C})$, $\langle i | (K |p\rangle \langle r|) | k \rangle = K_{ip} \delta_{rk}$. In (1.59), we can further sum over either $r = k$ or $i = p$; in either cases, as $\langle \phi | \psi \rangle = 0$, we find $\text{Tr} \Psi \Phi^\dagger = (\text{Tr} \Phi \Psi^\dagger)^* = 0$ and the result follows.

Lemma 1.4 In the hypothesis of Lemma 1.2, the matrix $C = [c_{ab}]$ in (1.58) must be positive definite.

Proof: With any $\vec{w} = \{w_a, a = 1, n^2 - 1\} \in \mathbf{C}^{n^2-1}$, we consider $W = \frac{1}{2} \sum_{a=1}^{n^2-1} w_a^* F_a$, which is a traceless $n \times n$ matrix. If matrices Ψ and Φ exist such that $\Phi \Psi^\dagger = W$ and $\Psi^\dagger \Phi = W^T$, then, from Lemmas 1.2 and 1.3 and from the orthogonality of the matrices F_a it follows

$$\mathcal{L}_{\phi, \psi} = \sum_{a, b=1}^{n^2-1} c_{ab} w_a^* w_b \geq 0, \quad (1.60)$$

whence the positivity of $C = [c_{ab}]$ and the proof of Theorem 1.6. Any matrix W and its transposed with respect to the given basis, W^T , have the same elementary divisors; therefore, they are similar to the same canonical Jordan form and thus similar to each other [50]. Let Φ such that $\Phi^{-1} W \Phi = W^T$, that is we take as vector $|\phi\rangle \in \mathfrak{H}_n \otimes \mathfrak{H}_n$ the one whose components ϕ_{ij} are the elements of the similarity matrix transforming the given W into its transposed W^T . It then follows that $\Psi^\dagger = \Phi^{-1} W$ and moreover $\Psi^\dagger \Phi = \Phi^{-1} W \Phi = W^T$, which is what we need.

1.7 The stochastic approach

The description of an open quantum system can be performed employing an approach different from the projector technique presented in Section (1.3), that is using a stochastic

equation [125]. The point of view is similar to that developed by Langevin to describe the Brownian motion: we introduce a time dependent term in the system Hamiltonian: $H_S(t) = H_0 + H_I(t)$, where $H_I(t)$ characterizes the interaction between S and reservoir. The statistical operator that describes S is obtained by an averaging procedure over the noise: $\rho_S(t) = \langle \Sigma_S(t) \rangle$, and $\Sigma_S(t)$ is the stochastic density matrix of S :

$$\dot{\Sigma}_S(t) = L_S(t)[\Sigma_S(t)] = (L_0 + L_I(t))[\Sigma_S(t)] \quad (1.61)$$

with Liouville operators $L_0[\Sigma_S(t)] = -i[H_0, \Sigma_S(t)]$ and $L_I(t)[\Sigma_S(t)] = -i[H_I(t), \Sigma_S(t)]$. In the interaction picture (1.61) reads

$$\dot{\Sigma}_S^i(t) = L_I^i(t)[\Sigma_S^i(t)], \quad (1.62)$$

where

$$\Sigma_S^i(t) = e^{itH_0} \Sigma_S(t) e^{-itH_0}, \quad L_I^i(t) = e^{-L_0 t} L_I(t) e^{L_0 t}. \quad (1.63)$$

The solution of (1.62) is

$$\Sigma_S^i(t) = \mathbf{T} e^{\int_0^t L_I^i(t') dt'} \Sigma_S^i(0), \quad (1.64)$$

where $\mathbf{T}$ means time-ordering. Since $\rho_S^i(t) = \langle \Sigma_S^i(t) \rangle$, it follows that

$$\rho_S^i(t) = \sum_{k=0}^{\infty} M_k(t) [\rho_S^i(0)] \quad (1.65)$$

with

$$M_k(t) \equiv \int_0^t ds_1 \int_0^{s_1} ds_2 \dots \int_0^{s_{k-1}} ds_k \langle L_I^i(s_1) L_I^i(s_2) \dots L_I^i(s_k) \rangle. \quad (1.66)$$

We assume that the initial state $\Sigma_S(0)$ is decoupled from the noisy environment so that $\rho_S(0) = \langle \Sigma_S(0) \rangle = \Sigma_S(0)$ (and then $\rho_S^i(0) = \Sigma_S^i(0)$). Eq. (1.65) is a series expansion in the interaction, containing the correlation functions of the fields defining $H_I(t)$. If we suppose these stochastic fields to be Gaussian, only the even terms in (1.65) survive.

Setting $M(t) \equiv \sum_k M_k(t) = \sum_k M_{2k}(t)$, we rewrite the differential form of equation (1.65) as follows:

$$\dot{\rho}_S^i(t) = \dot{M}(t) M^{-1}(t) [\rho_S^i(t)] = \left(\dot{M}_2(t) + \dot{M}_4(t) - \dot{M}_2(t) M_2(t) + \dots \right) [\rho_S^i(t)] \quad (1.67)$$

using the formal expressions

$$M(t) = \mathbf{I} + M_2(t) + M_4(t) + \dots, \quad M^{-1}(t) = \mathbf{I} - M_2(t) + [M_2(t)]^2 - M_4(t) + \dots \quad (1.68)$$

Assuming the system S to be a n -level system, it is possible to express the interaction Hamiltonian as $H_I(t) = \sum_{k=0}^{n^2-1} V_k(t) F_k$, where F_k is the basis of $M_n(\mathbf{C})$ introduced in (1.49) and $V_k(t)$ are Gaussian stochastic fields with correlation functions $h_{kl}(t-s) = \langle V_k(t) V_l(s) \rangle$. Then, up to the second order in the interaction, eq. (1.67) assumes the form

$$\dot{\rho}_S^i(t) = - \sum_{k,l=1}^{n^2-1} \int_0^t ds h_{kl}(s) [F_k(t), [F_l(t-s), \rho_S^i(t)]], \quad (1.69)$$

with $F_k(t) \equiv e^{L_0 t} [F_k] = e^{itH_0} F_k e^{-itH_0}$.

Turning back to the Schrödinger picture, $\rho_S(t) = e^{-itH_0} \rho_S^i(t) e^{itH_0}$ and we find:

$$\dot{\rho}_S(t) = -i[H_0, \rho_S(t)] + \sum_{k,j=1}^{n^2-1} d_{kj}(t) [F_k, [F_j, \rho_S(t)]], \quad (1.70)$$

with

$$d_{kj}(t) = - \sum_{l=1}^{n^2-1} \int_0^t h_{kl}(s) u_{lj}(-s) ds \quad (1.71)$$

and $u_{lj}(-s)$ defined by $F_l(-s) = \sum_{j=0}^{n^2-1} u_{lj}(-s) F_j$.

The Markovian approximation of the reduced dynamics (1.70) is performed with the hypothesis that the correlation functions $h_{kl}(t)$ decay so fast that we can substitute $+\infty$ for the parameter t in (1.71); in this way the coefficients $d_{kj}(t)$ become time independent. This assumption, as we have already seen, is the same as considering the system evolution for times long enough, so that the transient terms (the memory contributions, non Markovian) disappear.

The reduced dynamics so obtained, although Markovian, in general is not completely positive nor positive (in fact the generator of the dissipative part in general has a different form than that expressed in Theorems 1.4 and 1.5): the stochastic approach suffers from this failure. Equations of motion in the form (1.70), in the two-dimensional case, after the Markovian limit, are called Redfield-Bloch equations [27, 125, 126]; in Chapter 5 we shall explicitly show instances where lack of positivity occurs.

It has been argued that it is possible to consistently use these time-evolutions, even if they are not positive, provided they act only on a suitable subset of $\mathcal{P}(\mathfrak{H}_n)$, constituted by statistical operators whose positivity is preserved. Such a restriction is usually explained in terms of a slippage of initial conditions [30, 31]: the initial transient, neglected in the Markov limit, is supposed to select from $\mathcal{P}(\mathfrak{H}_n)$ exactly the set of states which remain positive under the dynamical map considered. In such a perspective, the long-time Markov approximation of the dynamics, possibly not positive, should be coupled to a redefinition of initial conditions, generated by the action, on $\mathcal{P}(\mathfrak{H}_n)$, of the short-time non-Markovian dynamics. In such a way the positivity of $\rho_S(t)$ should be saved. However, it can be shown

that the prescription needed to guarantee the consistency of the action of a Markovian dynamical map Λ_t in general does not hold when two systems are immersed in the same environment and evolve with the factorized dynamics $\Lambda_t \otimes \Lambda_t$, starting from an entangled state: a new prescription has to be enforced [51]. This argument can be repeated when three or more systems, living in the same environment, are taken into account: there does not exist a general prescription, valid for any number of systems interacting with the same environment. In this sense, complete positivity appears at least as a cheaper tool to avoid physical inconsistencies.

1.8 Entropy and long time behavior in open quantum systems

Typically, an open system exchanges energy and entropy (also matter in general) with its environment. Because of the vastly different number of their degrees of freedom, one expects the noise coming from the environment to always increase the randomness of the system and thus its entropy.

For later use, in this section we shortly review some results about the long time behavior of a state evolving according to quantum dynamical semigroups, and of its entropy.

Definition 1.12 Given a statistical operator $\rho \in \mathcal{P}(\mathfrak{H})$, its von Neumann entropy $S(\rho)$ is defined by $S(\rho) \equiv -\text{Tr}(\rho \log \rho)$.

For a review about entropy and its properties, see [56]; here it is sufficient to remark that the von Neumann entropy is always non-negative, $S(\rho) \geq 0$, $\forall \rho \in \mathcal{P}(\mathfrak{H})$, and $S(\rho) = 0$ if and only ρ is a pure state. Furthermore, if $\mathfrak{H}$ is finite dimensional, say $\dim(\mathfrak{H}) = n$, the entropy is greatest for the maximally mixed state $\rho_n \equiv \mathbf{I}_n/n$: $S(\rho_n) = \log n$.

The time-evolution of entropy is related to the long time behavior of states evolving under quantum dynamical semigroups, to the existence of invariant states, and to the approach to equilibrium of the considered open systems.

Definition 1.13 A state $\rho_0 \in \mathcal{P}(\mathcal{H})$ is said *invariant* (or *stationary*) under the action of a quantum dynamical semigroup $\{\Lambda_t, t \geq 0\}$ iff $\Lambda_t \rho_0 = \rho_0$, $\forall t \geq 0$ or, equivalently, $L[\rho_0] = 0$, where L is the generator of the semigroup, expressed in (1.47).

Definition 1.14 The dynamical semigroup $\{\Lambda_t, t \geq 0\}$ is called *relaxing* iff there exists an invariant state $\rho_0 \in \mathcal{P}(\mathcal{H})$ such that $\forall \rho \in \mathcal{P}(\mathcal{H})$, $\lim_{t \rightarrow \infty} \Lambda_t \rho = \rho_0$.

A complete classification of all types of stationary states associated to relaxing semigroups is given in [52]; see also [53, 54]. For later reference we are only interested in the following considerations.

As we have seen in Section 1.5, the dual time-evolution Λ_t^* always preserves the identity $\mathbf{I}$, and then its generator, denoted by L^* , maps $\mathbf{I}$ into the null operator: $L^*[\mathbf{I}] = 0$. In

general, the same does not hold for the evolution map Λ_t and $L[\mathbf{I}] \neq 0$. However, as we shall shortly see, there is a tight connection between the entropy time-evolution of the system and the invariance of $\mathbf{I}$, technically called unitality.

Definition 1.15 A quantum dynamical semigroup $\{\Lambda_t, t \geq 0\}$ is called *unital* iff the identity operator $\mathbf{I} \in \mathcal{P}(\mathcal{H})$, is invariant under its action.

Remark 1.21 If Λ_t is unital, the maximally mixed state ρ_n is invariant.

Remark 1.22 If Λ_t is unital, the operators A_k defined in Corollary 1.1 must satisfy $\sum_k A_k A_k^\dagger = \mathbf{I}$.

Remark 1.23 If Λ_t is unital, the operators V_k defined in Theorem 1.4 must satisfy $\sum_k [V_k, V_k^\dagger] = 0$.

Now, it turns out that, for an n -level system, the unitality of the dynamics is a necessary condition for the non-decreasing behavior of its entropy. Indeed, if unitality is not fulfilled, the maximally mixed state ρ_n is not invariant under the reduced dynamics considered: $\dot{\rho}_n = L[\rho_n] \neq 0$; therefore $\rho_n(t) = \Lambda_t[\rho_n] \neq \rho_n$ and then, necessarily, the entropy decreases: $S[\rho_n(t)] < S(\rho_n)$.

In the next chapter we shall prove that, for $n = 2$, the unitality of quantum dynamical semigroups is also a sufficient condition in order to have a non-decreasing entropy.

Chapter 2

Bidimensional Systems

In view of later applications, in this chapter we restrict our attention to quantum open systems described by two-dimensional Hilbert spaces. The parameters describing general Markovian dynamics are introduced; positivity and complete positivity of the time-evolution are expressed by sets of inequalities among them.

Although the equations of motion can be integrated exactly, in many cases it is preferable to express their solutions as perturbative expansions in the dissipative parameters, provided they are small with respect to the parameters describing the dynamics of the system alone, in absence of the environment. This computation, to leading order in the dissipative parameters, is performed in Section 2.4.

2.1 Evolution of states

The physical systems we shall consider in the next chapters will be described by means of two-dimensional Hilbert spaces. Therefore, their states will be represented by density matrices corresponding to positive elements of $M_2(\mathbf{C})$ with unit trace. We will use two different basis for the algebra $M_2(\mathbf{C})$, namely:

- the set $\{P_i, i = 1, \dots, 4\}$, where

$$P_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad P_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad P_3 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad P_4 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad (2.1)$$

with $P_{1,2}^\dagger = P_{1,2}$ and $P_{3,4}^\dagger = P_{4,3}$;

- the set $\{\sigma_i, i = 0, \dots, 3\}$, that is the set of Pauli matrices plus the identity:

$$\sigma_0 = \mathbf{I}_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (2.2)$$

satisfying $\sigma_i^\dagger = \sigma_i$, $i = 0, \dots, 3$; $\text{Tr}(\sigma_i) = 0$, $i = 1, \dots, 3$; $\text{Tr}(\sigma_i^\dagger \sigma_j) = 2\delta_{ij}$, $i, j = 0, \dots, 3$.

Accordingly, states ρ will be written, with respect to these two basis, as

$$\begin{aligned} \rho &= \sum_{i=1}^4 \rho_i P_i = \begin{pmatrix} \rho_1 & \rho_3 \\ \rho_4 & \rho_2 \end{pmatrix} \\ &= \sum_{i=0}^3 \rho^i \sigma_i = \begin{pmatrix} \rho^0 + \rho^3 & \rho^1 - i\rho^2 \\ \rho^1 + i\rho^2 & \rho^0 - \rho^3 \end{pmatrix}. \end{aligned} \quad (2.3)$$

We use upper and lower indices to distinguish the coefficients in the two different expansions. We have $\rho_i \in \mathbf{C}$ and $\rho_4 = \rho_3^\dagger$, while $\rho^i \in \mathbf{R}$.

We are interested in Markovian dynamics for the states ρ ; they are fully characterized by Theorem 1.5 and Corollary 1.3. The basis $\{F_i \equiv \sigma_i/\sqrt{2}, i = 0, \dots, 3\}$ fulfills the properties required in Theorem 1.5: $F_0 = \mathbf{I}_2/\sqrt{2}$, $\text{Tr}(F_i) = 0 \ \forall i \neq 0$, $\text{Tr}(F_i^\dagger F_j) = \delta_{ij}$; then, the more general form for the generator of a Markovian dynamics is

$$L[\rho] = (L_H + L_D)[\rho], \quad (2.4)$$

where $L_H[\cdot]$ and $L_D[\cdot]$ are the Hamiltonian, respectively dissipative part, of the generator. L_H , for a stable system, is proportional to the commutator with a Hamiltonian $H = H^\dagger$ that can be chosen traceless without loss of generality, then $H = \sum_{i=1}^3 h_i \sigma_i$, $h_i \in \mathbf{R}$; L_D can be written in the Kossakowski form:

$$L_H[\rho] \equiv -i[H, \rho], \quad L_D[\rho] \equiv \frac{1}{2} \sum_{k,l=1}^3 c_{kl} \left(\sigma_k \rho \sigma_l - \frac{1}{2} \{\rho, \sigma_l \sigma_k\} \right) \quad (2.5)$$

with $C = [c_{kl}]$ a Hermitian matrix.

For unstable systems H is non-Hermitian (i.e. $h_i \in \mathbf{C}$) of Weisskopf-Wigner type (see the discussion in Section 1.1) and the time-evolution does not preserve the trace of the states ρ ; the generator has thus the form (2.4) with a modified Hamiltonian part:

$$L_H[\rho] \equiv -iH\rho + i\rho H^\dagger \quad (2.6)$$

and will be used when dealing with neutral mesons systems; at present we restrict our attention to trace preserving dynamics with L_H in form (2.5).

It is convenient to introduce a vector notation and write the density matrix ρ as a 4-component vector $|\rho\rangle \equiv (\rho^0, \rho^1, \rho^2, \rho^3)$ whose entries are the coefficients of the expansion of ρ in the basis (2.2); in fact, since equation (2.4) is linear in ρ , it can be rewritten in vectorial form as a kind of Schrödinger-like equation:

$$\frac{d}{dt} |\rho(t)\rangle = \mathcal{L} |\rho(t)\rangle = (\mathcal{H} + \mathcal{D}) |\rho(t)\rangle, \quad (2.7)$$

where $\mathcal{H}$ and $\mathcal{D}$ are 4×4 real matrices, representing the Hamiltonian, respectively the dissipative part of (2.4). These matrices have the form

$$\mathcal{H} = -2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & h_3 & -h_2 \\ 0 & -h_3 & 0 & h_1 \\ 0 & h_2 & -h_1 & 0 \end{pmatrix}, \quad \mathcal{D} = -2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ u & a & b & c \\ v & b & \alpha & \beta \\ w & c & \beta & \gamma \end{pmatrix}. \quad (2.8)$$

In (2.8) we have introduced nine real coefficients to express the dissipative part $\mathcal{D}$. They depend on the entries of the matrix C in (2.5); explicitly [1, 55]:

$$\mathcal{D} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 2 \operatorname{Im} c_{32} & -c_{22} - c_{33} & \operatorname{Re} c_{12} & \operatorname{Re} c_{13} \\ 2 \operatorname{Im} c_{13} & \operatorname{Re} c_{12} & -c_{11} - c_{33} & \operatorname{Re} c_{23} \\ 2 \operatorname{Im} c_{21} & \operatorname{Re} c_{13} & \operatorname{Re} c_{23} & -c_{11} - c_{22} \end{pmatrix}; \quad (2.9)$$

conversely, the matrix C can be written in terms of these dissipative parameters:

$$C = \begin{pmatrix} \alpha + \gamma - a & -2b + iw & -2c - iv \\ -2b - iw & a + \gamma - \alpha & -2\beta + iu \\ -2c + iv & -2\beta - iu & a + \alpha - \gamma \end{pmatrix}. \quad (2.10)$$

Remark 2.1 The Hamiltonian contribution $\mathcal{H}$ is skew-symmetric since it represent a commutator.

Remark 2.2 The entries of the first row of both $\mathcal{H}$ and $\mathcal{D}$ vanish since we have required the dynamics to be trace preserving. Indeed, from (2.8) it follows that $\dot{\rho}^0(t) = 0$, as it should be since $\operatorname{Tr}[\rho(t)] = 2\rho^0(t)$ and then $d\operatorname{Tr}[\rho(t)]/dt = 0$.

Using the basis (2.1) ρ gets represented by the 4-vector $|\hat{\rho}\rangle \equiv (\rho_1, \rho_2, \rho_3, \rho_4)$, which evolves in time according to

$$\frac{d}{dt}|\hat{\rho}(t)\rangle = \hat{\mathcal{L}}|\rho(t)\rangle = (\hat{\mathcal{H}} + \hat{\mathcal{D}})|\hat{\rho}(t)\rangle, \quad (2.11)$$

where $\hat{\mathcal{H}}$ and $\hat{\mathcal{D}}$ are respectively the Hamiltonian and the dissipative contribution in this representation¹. From the relations in (2.3) it follows that the two vectors representing the state ρ are related by

$$|\hat{\rho}(t)\rangle = \mathcal{V}|\rho(t)\rangle, \quad \mathcal{V} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & -1 \\ 0 & 1 & -i & 0 \\ 0 & 1 & i & 0 \end{pmatrix}. \quad (2.12)$$

¹We use hats to characterize states as well as operators in this representation

Thus,

$$\hat{\mathcal{H}} = \mathcal{V}\mathcal{H}\mathcal{V}^{-1} = \begin{pmatrix} 0 & 0 & -h_2 + ih_1 & -h_2 - ih_1 \\ 0 & 0 & h_2 - ih_1 & h_2 + ih_1 \\ h_2 + ih_1 & -h_2 - ih_1 & -2ih_3 & 0 \\ h_2 - ih_1 & -h_2 + ih_1 & 0 & 2ih_3 \end{pmatrix} \quad (2.13)$$

and

$$\hat{\mathcal{D}} = \mathcal{V}\mathcal{D}\mathcal{V}^{-1} = \begin{pmatrix} -\gamma - w & \gamma - w & -C & -C^* \\ \gamma + w & w - \gamma & C & C^* \\ -C^* - U^* & C^* - U^* & -A & B \\ -C - U & C - U & B^* & -A \end{pmatrix} \quad (2.14)$$

with

$$A \equiv a + \alpha, \quad B \equiv \alpha - a + 2ib, \quad C \equiv c + i\beta, \quad U \equiv u + iv. \quad (2.15)$$

Before discussing solutions of the equations of motion we discuss the positivity of the dynamics.

2.2 Positivity and Complete Positivity

The requests of positivity or complete positivity for the time-evolutions described in the previous section are expressed by different constraints on the dissipative parameters appearing in $\mathcal{D}$.

Remark 2.3 The positivity of $A \in M_2(\mathbf{C})$ is equivalent to $\text{Tr}(A) \geq 0$ and $\text{Det}(A) \geq 0$, since we are in two dimensions. Since states ρ have $\text{Tr}(\rho) = 1$, so that $\rho^0 = 1/2$, they are positive if and only if $\text{Det}(\rho) = 1/4 - \sum_{j=1}^3 (\rho^j)^2 \geq 0$. A state ρ is pure (i.e. a one-dimensional projection) if and only if $\text{Det}(\rho) = 0$. A mixed state has $\sum_{j=1}^3 (\rho^j)^2 < 1/4$.

Theorem 2.1 If the trace and Hermiticity preserving time-evolution (2.7) with $\mathcal{H}$ and $\mathcal{D}$ in (2.8) is positive, then the real entries of $\mathcal{D}$ satisfy the inequalities:

$$\begin{cases} a \geq 0 \\ \alpha \geq 0 \\ \gamma \geq 0 \end{cases} ; \quad \begin{cases} a\alpha \geq b^2 \\ a\gamma \geq c^2 \\ \alpha\gamma \geq \beta^2 \end{cases} ; \quad a\alpha\gamma + 2bc\beta - a\beta^2 - \alpha c^2 - \gamma b^2 \geq 0. \quad (2.16)$$

Proof: $\rho(t) \geq 0$ if and only if $\text{Det}[\rho(t)] \geq 0$ (Remark 2.3). Choosing the initial state $|\rho_a(0)\rangle = (1/2, \rho^1, \rho^2, \rho^3)$ to be pure, that is $\sum_{j=1}^3 (\rho^j)^2 = 1/4$, then $d\text{Det}[\rho_a(t)]/dt|_{t=0} \geq 0$

must hold if $\rho_a(t)$ has to remain a state. Explicitly,

$$\begin{aligned} \frac{d}{dt} \text{Det}[\rho_a(t)]|_{t=0} &= -2 \sum_{j=1}^3 \rho^j(t) \dot{\rho}^j(t)|_{t=0} \\ &= -2 \sum_{j,k=1}^3 (\mathcal{H} + \mathcal{D})_{jk} \rho^j(0) \rho^k(0) - \sum_{j=1}^3 \mathcal{D}_{j0} \rho^j(0) \geq 0. \end{aligned} \quad (2.17)$$

Analogously, starting with the pure state $|\rho_b(0)\rangle = (1/2, -\rho^1, -\rho^2, -\rho^3)$ we find

$$\frac{d}{dt} \text{Det}[\rho_b(t)]|_{t=0} = -2 \sum_{j,k=1}^3 (\mathcal{H} + \mathcal{D})_{jk} \rho^j(0) \rho^k(0) + \sum_{j=1}^3 \mathcal{D}_{j0} \rho^j(0) \geq 0; \quad (2.18)$$

adding equations (2.17) and (2.18) and considering that $\mathcal{H}$ is a skew symmetric matrix, we are left with

$$\sum_{j,k=1}^3 \mathcal{D}_{jk} \rho^k(0) \rho^j(0) \leq 0. \quad (2.19)$$

Introducing the sub-matrix $\tilde{\mathcal{D}}$ as

$$\mathcal{D} = -2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ u & & & \\ v & \tilde{\mathcal{D}} & & \\ w & & & \end{pmatrix}, \quad \tilde{\mathcal{D}} \equiv \begin{pmatrix} a & b & c \\ b & \alpha & \beta \\ c & \beta & \gamma \end{pmatrix}, \quad (2.20)$$

the condition (2.19) becomes $\tilde{\mathcal{D}} \geq 0$; which is equivalent to (2.16) by considering the minors of $\tilde{\mathcal{D}}$.

Theorem 2.2 [1] The trace and Hermiticity preserving time-evolution (2.7) is completely positive iff the entries of $\mathcal{D}$ satisfy the inequalities:

$$\begin{cases} 2R \equiv \alpha + \gamma - a \geq 0 \\ 2S \equiv a + \gamma - \alpha \geq 0 \\ 2T \equiv a + \alpha - \gamma \geq 0 \end{cases} ; \quad \begin{cases} RS \geq b^2 + w^2/4 \\ RT \geq c^2 + v^2/4 \\ ST \geq \beta^2 + u^2/4 \end{cases} ;$$

$$RST - 2bc\beta + \frac{1}{2}buw + \frac{1}{2}\beta vw + \frac{1}{2}cuw - S(c^2 + \frac{1}{4}v^2) - R(\beta^2 + \frac{1}{4}u^2) - T(b^2 + \frac{1}{4}w^2) \geq 0. \quad (2.21)$$

Proof: The thesis follows from the fact that the time-evolution is completely positive if and only if the matrix C in (2.10) is positive (Theorem 1.5).

Remark 2.4 Note that the constraints expressed in these two theorems are not all independent. Also, the constraints (2.21) imply those in (2.16). Indeed, CP is stronger than positivity.

2.3 Positivity and CP for entropy increasing evolutions

There are physical situations in which the reduced time-evolution of states does necessarily increase the von Neumann entropy (see Definition 1.12) and states becomes less ordered as the time passes. For instance, in Chapters 3 and 5 we shall describe irreversible time-evolutions that are entropy increasing.

As anticipated in Section 1.8, any entropy non decreasing time-evolution can be fully characterized as follows.

Theorem 2.3 The trace and Hermiticity preserving time-evolution (2.7) with $\mathcal{H}$ and $\mathcal{D}$ in (2.8) has a non decreasing von Neumann entropy iff $u = v = w = 0$ and $\tilde{\mathcal{D}} \geq 0$ (i.e. the time evolution (2.7) is unital), where u , v and w are the parameters defined in equations (2.8) and (2.9) and $\tilde{\mathcal{D}}$ is the matrix in eq. (2.20). Moreover, the matrix C in (2.10) is symmetric [39, 55].

Proof: The eigenvalues, respectively eigenvectors, of a generic initial state represented by the vector $|\rho(0)\rangle = (1/2, \rho^1, \rho^2, \rho^3)$ with $|\rho|^2 \equiv \sum_{j=1}^3 (\rho^j)^2 \leq 1/4$, are

$$\rho_{\pm} = \frac{1}{2}(1 \pm 2|\rho|); \quad |\rho_{\pm}\rangle = \frac{1}{2|\rho|}(|\rho|, \pm\rho^1, \pm\rho^2, \pm\rho^3). \quad (2.22)$$

Consider now

$$\frac{d}{dt}S[\rho(t)]|_{t=0} = -\frac{d}{dt}\text{Tr}[\rho(t)\log\rho(t)]|_{t=0} = -\text{Tr}[\dot{\rho}(t)|_{t=0}\log\rho(0)] \quad (2.23)$$

where we have made use of the trace preserving property of the time-evolution. Evaluating the trace over the states $|\rho_+\rangle$ and $|\rho_-\rangle$ we find

$$\begin{aligned} \frac{d}{dt}S[\rho(t)]|_{t=0} &= -\log\rho_+ \text{Tr}[\dot{\rho}(t)|_{t=0}|\rho_+\rangle\langle\rho_+|] - \log\rho_- \text{Tr}[\dot{\rho}(t)|_{t=0}|\rho_-\rangle\langle\rho_-|] = \\ &= -\frac{1}{|\rho|} \left[\frac{1}{2} \sum_{i=1}^3 \mathcal{D}_{i0} \rho^i(0) + \sum_{i,j=1}^3 \mathcal{D}_{ij} \rho^i(0) \rho^j(0) \right] \log \frac{\rho_+}{\rho_-}, \end{aligned} \quad (2.24)$$

having taken into account the skew-symmetry of $\mathcal{H}$.

Suppose $u = v = w = 0$ (that is $\mathcal{D}_{i0} = 0 \forall i$) and $\tilde{\mathcal{D}} \geq 0$. Then

$$\frac{d}{dt}S[\rho(t)]|_{t=0} = \frac{2}{|\rho|} \sum_{i,j=1}^3 \tilde{\mathcal{D}}_{ij} \rho^i(0) \rho^j(0) \log \frac{\rho_+}{\rho_-} \quad (2.25)$$

and, as $\rho_+ \geq \rho_-$, the von Neumann entropy is not decreasing.

Vice versa, suppose $\frac{d}{dt}S[\rho(t)]|_{t=0} \geq 0$ and choose $\rho^1 = \varepsilon$, $\rho^2 = \rho^3 = 0$, with $|\varepsilon| \leq 1/2$. Then

$$\frac{d}{dt}S[\rho(t)]|_{t=0} = \frac{2}{|\varepsilon|} \left(\frac{1}{2} u \varepsilon + a \varepsilon^2 \right) \log \frac{\rho_+}{\rho_-}. \quad (2.26)$$

Since $|u| \leq a$ (see Lemma 1.1), the right hand side of (2.26) could be made negative by choosing $\text{sign}(\varepsilon) = -\text{sign}(u)$ and $|\varepsilon| \leq |u|/2a$, unless $u = 0$. The same argument implies $v = 0$ and $w = 0$ choosing $\rho^2 = \varepsilon$, $\rho^1 = \rho^3 = 0$ and respectively $\rho^3 = \varepsilon$, $\rho^1 = \rho^2 = 0$. Then eq. (2.25) holds, whose positivity imply $\tilde{\mathcal{D}}_{ij} \geq 0$.

Theorem 2.4 The trace and Hermiticity preserving, entropy non decreasing, time evolution (2.7) is positive iff the real parameters of $\mathcal{D}$ satisfy the inequalities (2.16).

Proof: The if part is a direct consequence of Theorem 2.1. Vice versa, considering a generic initial state $|\rho(0)\rangle = (1/2, \rho^1, \rho^2, \rho^3)$ with $\text{Det}[\rho(0)] \geq 0$, we have

$$\frac{d}{dt} \text{Det}[\rho(t)]|_{t=0} = \sum_{i,j=1}^3 \tilde{\mathcal{D}}_{ij} \rho^i(0) \rho^j(0) \geq 0 \quad (2.27)$$

since $\tilde{\mathcal{D}} \geq 0$. Consequently $\text{Det}[\rho(t)] \geq 0$, $\forall t$; thus the evolution is positive (Remark 2.3). Indeed, suppose there is a time $\bar{t}$ for which $\text{Det}[\rho(\bar{t})] < 0$. By continuity, there must be a time t' , with $0 < t' < \bar{t}$ such that $\text{Det}[\rho(t')] = 0$ and $d\text{Det}[\rho(t')]/dt < 0$. But this can never happen, if $\tilde{\mathcal{D}} \geq 0$.

Theorem 2.5 [1, 44, 45, 46, 47, 48, 49, 55] The trace and Hermiticity preserving, entropy non decreasing, time-evolution (2.7) is completely positive iff the real parameters of $\mathcal{D}$ satisfy the inequalities

$$\begin{cases} 2R \equiv \alpha + \gamma - a \geq 0 \\ 2S \equiv a + \gamma - \alpha \geq 0 \\ 2T \equiv a + \alpha - \gamma \geq 0 \end{cases} ; \begin{cases} RS \geq b^2 \\ RT \geq c^2 \\ ST \geq \beta^2 \end{cases} ; RST - 2bc\beta - Sc^2 - R\beta^2 - Tb^2 \geq 0. \quad (2.28)$$

Proof: The thesis follows from Theorem 2.2 putting $u = v = w = 0$.

2.4 Perturbative expansion

We want to integrate the equation of motion for ρ , i.e. we want to solve either (2.7) or (2.11). In vectorial notation the solutions will be denoted by

$$|\rho(t)\rangle = \mathcal{G}_t |\rho(0)\rangle, \quad \mathcal{G}_t \equiv e^{\mathcal{L}t} = e^{(\mathcal{H}+\mathcal{D})t} \quad (2.29)$$

and

$$|\hat{\rho}(t)\rangle = \hat{\mathcal{G}}_t |\hat{\rho}(0)\rangle, \quad \hat{\mathcal{G}}_t \equiv e^{\hat{\mathcal{L}}t} = e^{(\hat{\mathcal{H}}+\hat{\mathcal{D}})t} \quad (2.30)$$

respectively. Both $\mathcal{G}_t$ and $\hat{\mathcal{G}}_t$ are 4×4 matrices and $\hat{\mathcal{G}}_t = \mathcal{V} \mathcal{G}_t \mathcal{V}^{-1}$, with $\mathcal{V}$ expressed in (2.12).

Although it is always possible, at least in principle, to compute the form of $\mathcal{G}_t$ and $\hat{\mathcal{G}}_t$ without approximations, often these explicit expressions are rather involved and not particularly useful in order to capture the essence of the dissipative processes.

Usually, however, the parameters appearing in the dissipative part of either (2.7) or (2.11) can be considered small with respect to those characterizing the Hamiltonian contribution. It is thus possible to express both $\mathcal{G}_t$ and $\hat{\mathcal{G}}_t$ as series expansions in these parameters [44, 45, 46, 47, 48, 49],

$$\mathcal{G}_t = \mathcal{G}_t^0 + \mathcal{G}_t^1 + \mathcal{G}_t^2 + \dots \quad (2.31)$$

with

$$\begin{aligned} \mathcal{G}_t^0 &\equiv e^{\mathcal{H}t}, \quad \mathcal{G}_t^1 \equiv \int_0^t ds_1 e^{\mathcal{H}(t-s_1)} \mathcal{D} e^{\mathcal{H}s_1}, \\ \mathcal{G}_t^2 &\equiv \int_0^t ds_1 \int_0^{s_1} ds_2 e^{\mathcal{H}(t-s_1)} \mathcal{D} e^{\mathcal{H}(s_1-s_2)} \mathcal{D} e^{\mathcal{H}s_2}, \quad \dots \end{aligned} \quad (2.32)$$

Equivalent expressions hold for $\hat{\mathcal{G}}_t$.

Remark 2.5 Note that $\hat{\mathcal{G}}_t^j = \mathcal{V} \mathcal{G}_t^j \mathcal{V}^{-1}$ for any j , that is the similarity relation holds at any order in the perturbative expansion.

In this section we write the time-evolutor to leading order in the dissipative coefficients. Indeed, in the coming chapters, we will encounter physical situations in which the dissipative parameters are small with respect to the parameters describing the free time-evolution (i.e. without environment) and the formalism here developed will be applied. We perform the computation in the representation (2.11).

We suppose H as general as possible, so that we allow for a non Hermitian part. Hence its eigenvalues ϵ_1 and ϵ_2 of H can be complex and its eigenstates non orthogonal:

$$\begin{cases} H|\epsilon_1\rangle = \epsilon_1|\epsilon_1\rangle \\ H|\epsilon_2\rangle = \epsilon_2|\epsilon_2\rangle \end{cases} \quad \epsilon_i = m_i - \frac{i}{2}\gamma_i \quad m_i, \gamma_i \in \mathbf{R}, \quad \langle \epsilon_1 | \epsilon_2 \rangle \neq 0; \quad (2.33)$$

m_i and γ_i are the energies, respectively the widths, of the eigenstates.

Since the underlying Hilbert space is two-dimensional, in any given basis the two eigenvectors can be written as $|\epsilon_1\rangle = (p_1, q_1)$ and $|\epsilon_2\rangle = (p_2, -q_2)$, with the normalization conditions

$$|p_1|^2 + |q_1|^2 = 1, \quad |p_2|^2 + |q_2|^2 = 1. \quad (2.34)$$

Remark 2.6 If H is Hermitian then $\langle \epsilon_1 | \epsilon_2 \rangle = 0$ and $\gamma_i = 0$, that is $\epsilon_i \in \mathbf{R}$. The condition of orthogonality of the two eigenstates reads

$$p_1 p_2^* - q_1 q_2^* = 0. \quad (2.35)$$

It is convenient to work in the basis that diagonalizes the matrix $\hat{\mathcal{H}}$, hence $\hat{\mathcal{D}}$ will be transformed in a more complex matrix²; the whole computation will result simpler [49]. Accordingly we choose as basis for our bidimensional Hilbert space $\mathfrak{H}_2$ the pair of eigenstates of the Hamiltonian, given in (2.33). H is diagonalized by means of an invertible transformation U (unitary if H is Hermitian); its diagonal form will be denoted by an upper index d , as any operator represented in the basis (2.33):

$$H^d = U^{-1} H U; \quad H^d = \begin{pmatrix} \epsilon_1 & 0 \\ 0 & \epsilon_2 \end{pmatrix}, \quad U = \begin{pmatrix} p_1 & p_2 \\ q_1 & -q_2 \end{pmatrix}. \quad (2.36)$$

It is useful to introduce the two ratios

$$r_1 \equiv \frac{p_1}{q_1}, \quad r_2 \equiv \frac{p_2}{q_2}; \quad (2.37)$$

this allows to write $U = RQ$, with

$$R = \begin{pmatrix} r_1 & r_2 \\ 1 & -1 \end{pmatrix}, \quad Q = \begin{pmatrix} q_1 & 0 \\ 0 & q_2 \end{pmatrix}, \quad (2.38)$$

so that the similarity transformation connecting H and H^d depends solely on the parameters (2.37):

$$H^d = R^{-1} H R. \quad (2.39)$$

Remark 2.7 Note that R is not unitary even if H is Hermitian. In this case U is unitary and then $r_1 r_2^* = 1$.

The evolution of a state ρ , expressed in (2.4) with Hamiltonian and dissipative contributions as in (2.5) or (2.6), in the new basis takes the form

$$\dot{\rho}^d(t) = (L_H^d + L_D^d)[\rho^d(t)], \quad (2.40)$$

where

$$\rho^d(t) = R^{-1} \rho(t) R^{-1\dagger}, \quad L_{H,D}^d[\cdot] = R^{-1} L_{H,D}[R \cdot R^\dagger] R^{-1\dagger}. \quad (2.41)$$

Using the vector representation of ρ^d we have

$$\frac{d}{dt} |\hat{\rho}^d(t)\rangle = \hat{\mathcal{L}}^d |\rho(t)\rangle = (\hat{\mathcal{H}}^d + \hat{\mathcal{D}}^d) |\hat{\rho}^d(t)\rangle, \quad (2.42)$$

where $|\hat{\rho}^d(t)\rangle$ is the vector whose components are the coefficients of the expansion of $\rho^d(t)$ in the basis (2.1). It is obtained from $|\hat{\rho}(t)\rangle$ by the linear transformation

$$|\hat{\rho}^d(t)\rangle = \mathcal{R} |\hat{\rho}(t)\rangle, \quad (2.43)$$

²Notice that the choice of the representation of the state and the choice of the basis of the Hilbert space are completely independent

whose explicit expressions can be obtained from (2.41):

$$\mathcal{R} = \frac{1}{|r_1 + r_2|^2} \begin{pmatrix} 1 & |r_2|^2 & r_2^* & r_2 \\ 1 & |r_1|^2 & -r_1^* & -r_1 \\ 1 & -r_2 r_1^* & -r_1^* & r_2 \\ 1 & -r_1 r_2^* & r_2^* & -r_1 \end{pmatrix}. \quad (2.44)$$

The 4×4 matrices in (2.42) can now be explicitly obtained:

$$\hat{\mathcal{H}}^d = \mathcal{R} \hat{\mathcal{H}} \mathcal{R}^{-1} = \begin{pmatrix} -\gamma_1 & 0 & 0 & 0 \\ 0 & -\gamma_2 & 0 & 0 \\ 0 & 0 & -\Gamma_- & 0 \\ 0 & 0 & 0 & -\Gamma_+ \end{pmatrix}, \quad (2.45)$$

with $\Gamma_{\pm} \equiv \Gamma \pm i\Delta m$, $\Gamma \equiv (\gamma_1 + \gamma_2)/2$ and $\Delta m \equiv m_2 - m_1$; and

$$\hat{\mathcal{D}}^d = \mathcal{R} \hat{\mathcal{D}} \mathcal{R}^{-1} = \begin{pmatrix} \Lambda + \lambda_1 & \Sigma + \sigma & \Delta + \delta_1 & \Delta^* + \delta_1^* \\ \Xi + \xi & \Lambda + \lambda_2 & \Phi + \phi_1 & \Phi^* + \phi_1^* \\ \phi_2 - \Phi^* & \delta_2 - \Delta^* & \Omega + \omega & \Theta + \theta \\ \phi_2^* - \Phi & \delta_2^* - \Delta & \Theta^* + \theta^* & \Omega + \omega^* \end{pmatrix}, \quad (2.46)$$

where small and capital Greek letters refer respectively to contributions related to the entropy behavior (i.e. functions of u , v and w in (2.8)) and remaining parameters (i.e. those that are functions of a , b , c , α , β and γ in (2.8)). Their expressions are listed in Appendix A, both for stable and unstable systems.

Remark 2.8 For later convenience we observe that $\lambda_1, \lambda_2, \sigma, \xi \in \mathbf{R}$ and $\delta_1, \delta_2, \phi_1, \phi_2, \omega, \theta \in \mathbf{C}$; on the other hand $\Lambda, \Xi, \Sigma, \Omega \in \mathbf{R}$ and $\Delta, \Phi, \Theta \in \mathbf{C}$.

In the basis of the eigenvectors of H and in the chosen vectorial representation, the evolution operator is denoted by $\hat{\mathcal{G}}_t^d$. Up to first order contributions in the dissipative parameters, $\hat{\mathcal{G}}_t^d = \hat{\mathcal{G}}_t^{d0} + \hat{\mathcal{G}}_t^{d1}$; from (2.45) we find

$$\hat{\mathcal{G}}_t^{d0} = e^{\hat{\mathcal{H}}^d t} = \begin{pmatrix} e^{-\gamma_1 t} & 0 & 0 & 0 \\ 0 & e^{-\gamma_2 t} & 0 & 0 \\ 0 & 0 & e^{-\Gamma_- t} & 0 \\ 0 & 0 & 0 & e^{-\Gamma_+ t} \end{pmatrix}. \quad (2.47)$$

The next order contribution is obtained, accordingly with (2.32), as

$$\hat{\mathcal{G}}_t^{d1} = \int_0^t e^{\hat{\mathcal{H}}^d(t-s)} \hat{\mathcal{D}}^d e^{\hat{\mathcal{H}}^d s} ds.$$

The operator $\hat{\mathcal{G}}_t^d$ so obtained contains terms which are linear in t ; they disappear if we reconstruct the exponential dependences out of them. For example, summing the contributions from $\hat{\mathcal{G}}_t^{d0}$ and $\hat{\mathcal{G}}_t^{d1}$, the entry in first column and row of $\hat{\mathcal{G}}_t^d$ for an unstable system reads

$$(\hat{\mathcal{G}}_t^d)_{11} = [1 + (\Lambda + \lambda_1)t]e^{-\gamma_1 t} \approx e^{(\Lambda + \lambda_1)t}e^{-\gamma_1 t};$$

by looking at this expression we observe that, at the first order in the dissipative contributions, what we physically mean by width of the state $|\epsilon_1\rangle$ is not the γ_1 appearing in the Hamiltonian H , but rather $\gamma_1 - \Lambda - \lambda_1$. Accordingly, we redefine γ_1 so that

$$(\hat{\mathcal{G}}_t^d)_{11} = e^{-(\gamma_1 - \Lambda - \lambda_1)t} \rightarrow e^{-\gamma_1 t}.$$

In the same way also γ_2 and $\Delta\Gamma_{\pm} \equiv \Delta\Gamma \pm 2i\Delta m$ get redefined. The final, explicit expressions of the components of $\hat{\mathcal{G}}_t^d$, both for stable or unstable systems, are collected in Appendix B.

Remark 2.9 In general, the n -th term in the expansion (2.31) contains polynomials in t of order n ; these terms represent truncated exponential behaviors and it is possible to consistently re-absorb them at any order in the perturbative expansion by reconstructing the exponentials they represent. A truncated perturbative expansions does not hold for a too long time since the neglected terms become more and more relevant as the time passes, then the reabsorption of the contributions polynomial in t could seem useless rather than physically justified. However, this procedure allows to redefine the parameters m_i and γ_i by dissipative terms; in such a way they coincide with physical quantities (see Chapter 3 for further details).

Remark 2.10 The time-evolutors in the physical basis are obtained as $\hat{\mathcal{G}}_t = \mathcal{R}^{-1}\hat{\mathcal{G}}_t^d\mathcal{R}$ and $\mathcal{G}_t = \mathcal{V}^{-1}\hat{\mathcal{G}}_t^d\mathcal{V}$ in the two different representations adopted. However their explicit expressions are unnecessary for the later developments; indeed in the formalism of density matrices the mean value of any observable, represented by a Hermitian operator A , is obtained by a trace operation over the evolving state $\rho(t)$; this operation is invariant under any change of basis in the underlying Hilbert space:

$$\langle A \rangle_{\rho} \equiv \text{Tr}[A\rho(t)] = \text{Tr}[A^d\rho^d(t)]. \quad (2.48)$$

It turns out to be more convenient to represent the operator A in the basis of the eigenstates of H and to compute $\langle A \rangle_{\rho}$ via (2.48), rather than re-expressing $\rho^d(t)$ in the physical basis.

Remark 2.11 In the case of a stable system, the reabsorption of the terms linear in time is more difficult because it involves also the off diagonal entries of $\hat{\mathcal{G}}_t^d$. The final expressions are reported in Appendix B.

These results will be used in the coming chapters for the description of the phenomenology of stable and unstable particles immersed in various specific environments.

Chapter 3

Neutral B Mesons

In this chapter we concentrate on the study of the neutral **B** mesons system treated as an open system.

This system is presently under both theoretical and experimental investigation since it enables the study of some specific properties of the Standard Model, e.g. the topic of the violation of the CP symmetry [57, 58] (charge conjugation plus space reversal), but also of the CPT symmetry (CP plus time reversal) and the flavors oscillations. In particular single **B** mesons or entangled pairs of **B** mesons physics are intensively studied at the colliders or at the so-called **B** mesons factories [87].

Usually, the neutral **B** mesons system is considered as a closed system. Instead, in the following, we assume this system to be immersed in a thermal bath of gravitational origin. We shall adopt a phenomenological approach, that does not make any hypothesis about the microscopic nature of the interaction between **B** mesons and environment; we suppose solely the coupling to be weak and the entropy non-decreasing. It turns out that the resulting irreversible dynamics for the **B** mesons can be parameterized by six phenomenological coefficients that affect in a very specific way the relevant physical quantities (decays rates and asymmetries) [49]. In principle, an experimental study of these quantities enable us to evaluate the magnitude of these dissipative parameters.

As usual, we shall describe our system by a bidimensional Hilbert space, using the formalism developed in Chapter 2 and in particular the time evolution operator calculated in Section 2.4.

This dissipative approach has been also applied to describe the neutral **K** mesons system [66, 67, 68, 69, 44, 45, 46, 47, 48]. For another approach to decoherence in the neutral **B** mesons system, characterized by only one dissipative parameter, see [70].

3.1 The physics of B mesons

In the standard phenomenological approach the neutral **B** meson system can be described by means of a state or, more in general, a statistical operator, in a two-dimensional Hilbert

space $\mathfrak{H}_2$. The effective Hamiltonian H of this system is a non Hermitian operator of the Wigner-Weisskopf type [57, 58],

$$H = M - \frac{i}{2}\Gamma; \quad H = \begin{pmatrix} H_1 & H_3 \\ H_4 & H_2 \end{pmatrix} \quad (3.1)$$

Several basis in $\mathfrak{H}_2$ are of interest for the discussion that follows:

- The set $\{|\mathbf{B}^0\rangle, |\overline{\mathbf{B}}^0\rangle\}$ of the physical states (called also flavor states);
- The set $\{|\mathbf{B}_L\rangle, |\mathbf{B}_H\rangle\}$ of the eigenstates of the Hamiltonian of the system:

$$\begin{cases} H|\mathbf{B}_L\rangle = \epsilon_L|\mathbf{B}_L\rangle, & \epsilon_L = m_L - \frac{i}{2}\gamma_L \\ H|\mathbf{B}_H\rangle = \epsilon_H|\mathbf{B}_H\rangle, & \epsilon_H = m_H - \frac{i}{2}\gamma_H \end{cases} \quad (3.2)$$

where m and γ denote respectively the masses and the widths of the two eigenstates and the subscripts L, H means “light”, “heavy” respectively, since it turns out that $m_L \ll m_H$ [57, 58]. These states are not ortogonal: $\langle \mathbf{B}_L | \mathbf{B}_H \rangle \neq 0$;

- The set $\{|\mathbf{B}_1\rangle, |\mathbf{B}_2\rangle\}$ of the eigenstates of CP , the operator performing the CP transformation (charge conjugation plus parity):

$$\begin{aligned} CP|\mathbf{B}_1\rangle &= |\mathbf{B}_1\rangle \\ CP|\mathbf{B}_2\rangle &= -|\mathbf{B}_2\rangle \end{aligned} \quad (3.3)$$

The eigenvalues of CP are 1 and -1 since $(CP)^2 = \sigma_0$. The CP operator exchange the two physical states:

$$\begin{aligned} CP|\mathbf{B}^0\rangle &= e^{i\varphi}|\overline{\mathbf{B}}^0\rangle \\ CP|\overline{\mathbf{B}}^0\rangle &= e^{-i\varphi}|\mathbf{B}^0\rangle \end{aligned} \quad (3.4)$$

and φ is a phase depending on the convention adopted in defining the physical states.

Remark 3.1 In the treatement of kaon phenomenology the distinction between the two eigenstates of H is based on the large difference between the two widths and the terms “short” and “long” are used.

The relations between the basis (3.2) and (3.3) are:

$$\begin{aligned} |\mathbf{B}_L\rangle &= p_L|\mathbf{B}^0\rangle + q_L|\overline{\mathbf{B}}^0\rangle, & |p_L|^2 + |q_L|^2 &= 1 \\ |\mathbf{B}_H\rangle &= p_H|\mathbf{B}^0\rangle - q_H|\overline{\mathbf{B}}^0\rangle, & |p_H|^2 + |q_H|^2 &= 1 \end{aligned} \quad (3.5)$$

and

$$\begin{aligned} |\mathbf{B}_1\rangle &= \frac{1}{\sqrt{2}}(|\mathbf{B}^0\rangle + |\overline{\mathbf{B}}^0\rangle) \\ |\mathbf{B}_2\rangle &= \frac{1}{\sqrt{2}}(|\mathbf{B}^0\rangle - |\overline{\mathbf{B}}^0\rangle) \end{aligned} \quad (3.6)$$

The distinction between these three basis is one of the reasons of the physical relevance of the **B** mesons system.

Since the physical states, produced in the interactions, are linear superpositions of the eigenstates of H , an initially pure state, e.g. $|\mathbf{B}^0\rangle$, would develop a non-vanishing component on the other state, e.g. $|\overline{\mathbf{B}}^0\rangle$, as time passes. We have then transitions between the states $|\mathbf{B}^0\rangle$ and $|\overline{\mathbf{B}}^0\rangle$, called flavor oscillations, as the probability of transition between the two flavors behaves like a periodic function.

On the other side the distinction between eigenstates of CP and H is a consequence of CP violation: it is not possible to diagonalise at the same time both these operators. In the standard Model, CP violation is due to the presence, in the Cabibbo Kobayashi Maskawa matrix¹, of a complex phase that cannot be reabsorbed by a redefinition of the fields describing the quarks [57, 58]; this violation is expected to be greater in the **B** mesons sector with respect to the **K** mesons case. CP invariance is equivalent to $|p_H| = |q_H| = |p_L| = |q_L|$.

The adopted phenomenological approach can further account for violations of other symmetries that are preserved by the Standard Model, like CPT invariance and the phenomenological rule $\Delta B = \Delta Q$. In this relation, B is the beauty quantum number and Q the charge of the decaying quarks.

Further, it is possible to describe time-evolutions different from the standard ones, taking into account a possible interaction with an external environment. As discussed below, the modifications of the relevant quantities produced by the irreversible dynamics turn out to have a characteristic signature: the open system dynamics cannot be confused with the others possible symmetries violations. To be as general as possible in our discussion, we shall consider all of these different possibilities together.

After the definition of the two ratios

$$r_L = \frac{p_L}{q_L}, \quad r_H = \frac{p_H}{q_H}; \quad (3.7)$$

the Hamiltonian (3.1) can be rewritten as

$$H = \frac{1}{r_L + r_H} \begin{pmatrix} r_L \epsilon_L + r_H \epsilon_H & r_L r_H (\epsilon_L - \epsilon_H) \\ \epsilon_L - \epsilon_H & r_L \epsilon_H + r_H \epsilon_L \end{pmatrix}. \quad (3.8)$$

¹This is the matrix giving the mixing in the hadronic sector of the Standard Model

For future reference we find useful to define three complex parameters, σ , ξ and θ [57, 58]:

$$\sigma = r_H r_L = \frac{H_3}{H_4}, \quad \xi = \frac{|\sigma| - 1}{|\sigma| + 1}, \quad \theta = \frac{r_L - r_H}{r_L + r_H} = \frac{H_1 - H_2}{\epsilon_L - \epsilon_H}; \quad (3.9)$$

as we shall see in the coming sections, by means of them it is possible to parametrize in a convenient way violations of the discrete symmetries.

3.2 The gravitational bath

The physical motivation of an open quantum system treatment of the neutral **B** mesons system (or a generic neutral mesons system) comes from quantum gravity. A satisfactory and complete theory of quantum gravity has not been found yet. Nevertheless, general physical arguments involving quantum fluctuations of the gravitational field lead to the conclusion that space-time could lose its continuum character at the Planck's scale. Several models have been proposed to account for this striking conclusion, for example the existence, at the Planck's scale, of virtual black holes [59, 60, 61, 62]; also the dynamics of string and branes could cause the appearance of a foamy nature of the space-time at microscopic scales [63, 64, 65]. In all these cases the space-time could act as an effective environment and all the physical systems would necessarily interact with it. Since the degrees of freedom characterizing such an environment are not accessible, any physical system should exhibit an open system dynamics, leading to loss of quantum coherence (decoherence) [66, 67, 68, 59, 69, 71, 72, 73, 74, 75, 76].

The microscopic details of the interaction between physical systems and gravitational bath are unnecessary to parametrize this dissipative evolution: we adopt a phenomenological approach, that is independent on the underlying fundamental dynamics. Indeed, the form of the reduced dynamics can be uniquely fixed by the requirements of being a completely positive, Markovian, entropy increasing evolution.

The Markovian hypothesis is justified by the fact that the action of the gravitational thermal bath on the neutral mesons system is very weak. The magnitude of the dissipative parameters may be estimated to be of order E_S^2/E_B , where E_S is the characteristic energy of the system and E_B is the fundamental scale for the bath. In our case $E_S = m_{\mathbf{B}} \approx 5$ GeV, the neutral **B** mesons mass, and $E_B = M_P \approx 10^{19}$ GeV, the Planck mass (the scale at which the space-time is expected to exhibit a kind of granularity). The magnitude of the dissipative parameters has thus an upper bound of about 10^{-18} GeV. The Markovian dynamics is justified, since this scale is very small respect to the scale characterizing the usual Hamiltonian dynamics of the **B** mesons system.

Remark 3.2 The gravitational thermal bath should affect any physical system. However, its effects on the majority of these systems would be completely negligible, so the systems characterized by energies in the sub-nuclear range or higher are the only that could feel

it. For example, for atomic phenomena, characterized by energies of about 100 KeV, the dissipative parameters would be as small as 10^{-27} GeV.

The hypothesis of entropy non decreasing evolution is based on the choice of time-evolutions leading to states less-ordered in time. The correlations contained in the coherence of states are lost as time passes.

The accuracy of present experiments on neutral **B** mesons is very high, so that one expects the dissipative effects induced by gravitational phenomena to be in the reach of present or planned experiments. This is certainly one of the motivations that justify the coming considerations.

3.3 The dissipative evolution

The equation of motion for the state ρ representing the **B** mesons is assumed to have the form of a quantum dynamical semigroup, with a non standard contribution L_D in the generator, due to the gravitational bath:

$$\dot{\rho}(t) = -iH\rho(t) + i\rho(t)H^\dagger + L_D[\rho(t)]. \quad (3.10)$$

with H as in (3.1) and L_D expressed in (2.6).

Working in the basis of physical states $\{|\mathbf{B}^0\rangle, |\bar{\mathbf{B}}^0\rangle\}$ and using the vector notation introduced in the previous chapter, we have

$$\frac{d}{dt}|\hat{\rho}(t)\rangle = (\hat{\mathcal{H}} + \hat{\mathcal{D}}')|\hat{\rho}(t)\rangle \quad (3.11)$$

with

$$\hat{\mathcal{H}} = \begin{pmatrix} 2\text{Im } H_1 & 0 & iH_3^* & -iH_3 \\ 0 & 2\text{Im } H_2 & -iH_4 & iH_4^* \\ iH_4^* & -iH_3 & i(H_2^* - H_1) & 0 \\ -iH_4 & iH_3^* & 0 & i(H_1^* - H_2) \end{pmatrix} \quad (3.12)$$

the matrix containing the standard Hamiltonian terms, the masses being redefined to take into account possible Hamiltonian dissipative contributions, while the matrix containing the non Hamiltonian contribution, causing decoherence of states and entropy increase, is

$$\hat{\mathcal{D}}' = \begin{pmatrix} -a & a & -C' & -C'^* \\ a & -a & C' & C'^* \\ -C'^* & C'^* & -A' & B' \\ -C' & C' & B'^* & -A' \end{pmatrix} \quad (3.13)$$

where

$$A' \equiv \gamma + \alpha, \quad B' \equiv \alpha - \gamma - 2i\beta, \quad C' \equiv c - ib. \quad (3.14)$$

The matrix in (3.13) differs from that appearing in (2.14) because here we assume the dissipative contribution to have the form $\mathcal{D}$ in (2.8), with $u = v = w = 0$; $\hat{\mathcal{D}}'$ is obtained from $\hat{\mathcal{D}}$ in (2.14) by a change of basis, from the eigenstates of CP to the basis of physical states.

The dissipative parameters a, b, c, α, β and γ are small with respect to the parameters characterizing the Hamiltonian contribution; then it is possible to apply the formalism developed in Section 2.4. The only difference is the use of $\hat{\mathcal{D}}'$ instead of $\hat{\mathcal{D}}$. The parameters a, A', B' and C' take the place of γ, A, B and C respectively.

In the basis $\{|\mathbf{B}_L\rangle, |\mathbf{B}_H\rangle\}$, that diagonalizes H , the statistical operator becomes

$$\rho^d = R^{-1} \rho R^{-1\dagger}, \quad R = \begin{pmatrix} r_L & r_H \\ 1 & -1 \end{pmatrix} \quad (3.15)$$

while (3.11) takes the form

$$\frac{d}{dt} |\hat{\rho}^d(t)\rangle = (\hat{\mathcal{H}}^d + \hat{\mathcal{D}}'^d) |\hat{\rho}^d(t)\rangle \quad (3.16)$$

where $\hat{\mathcal{H}}^d$ is a diagonal Hamiltonian part, see (2.45), and $\hat{\mathcal{D}}'^d$ is a more involved dissipative part:

$$\hat{\mathcal{D}}'^d = \begin{pmatrix} \Lambda' & \Sigma' & \Delta' & \Delta'^* \\ \Xi' & \Lambda' & \Phi' & \Phi'^* \\ -\Phi'^* & -\Delta'^* & \Omega' & \Theta' \\ -\Phi' & -\Delta' & \Theta'^* & \Omega' \end{pmatrix}; \quad (3.17)$$

the coefficients $\Lambda', \Sigma', \Xi', \Delta', \Phi', \Theta'$ and Ω' are expressed as the corresponding parameters in Appendix A with the substitutions $\gamma \rightarrow a, A \rightarrow A', B \rightarrow B'$ and $C \rightarrow C'$. They are linear superpositions of a, b, c, α, β and γ with coefficients that depend on r_L and r_H , or, equivalently, on θ and σ . In the next section we will show that θ and σ are related to violations of the discrete symmetries, that are expected to be small; it is then justified to neglect the terms where these coefficients multiply the dissipative parameters. We finally find

$$\begin{aligned} \Lambda' &= \frac{1}{2} \left\{ \text{Re}[(\alpha - \gamma + 2i\beta)\sigma] - (\alpha + \gamma) \right\}, \\ \Sigma' &= -\Lambda', \\ \Delta' &= -\frac{1}{2} \left\{ 2 \text{Re}[(c - ib)\sqrt{\sigma}] + i \text{Im}[(\alpha - \gamma + 2i\beta)\sigma] \right\}, \\ \Xi' &= -\Lambda', \\ \Phi' &= -\Delta', \\ \Omega' &= -\frac{1}{2} \left\{ (\alpha + \gamma) + 2a + \text{Re}[(\alpha - \gamma + 2i\beta)\sigma] \right\}, \\ \Theta' &= -\frac{1}{2} \left\{ 2a - (\alpha + \gamma) - \text{Re}[(\alpha - \gamma + 2i\beta)\sigma] + 4i \text{Im}[(c - ib)\sqrt{\sigma}] \right\}. \end{aligned} \quad (3.18)$$

Since we have request that entropy be non decreasing, the remaining coefficients in Appendix A vanish. (see Section 2.3)

By means of the coefficients (3.18) we can compute the time evoluter $\hat{\mathcal{G}}_t^d$ acting on our states; see Appendix B:

$$|\hat{\rho}_{\mathbf{B}^0}^d(t)\rangle = \hat{\mathcal{G}}_t^d |\hat{\rho}_{\mathbf{B}^0}^d(0)\rangle, \quad |\hat{\rho}_{\bar{\mathbf{B}}^0}^d(t)\rangle = \hat{\mathcal{G}}_t^d |\hat{\rho}_{\bar{\mathbf{B}}^0}^d(0)\rangle \quad (3.19)$$

with the initial conditions

$$\rho_{\mathbf{B}^0}(0) = |\mathbf{B}^0\rangle\langle\mathbf{B}^0|, \quad \rho_{\bar{\mathbf{B}}^0}(0) = |\bar{\mathbf{B}}^0\rangle\langle\bar{\mathbf{B}}^0|. \quad (3.20)$$

The operators $\rho_{\mathbf{B}^0}^d(t)$ and $\rho_{\bar{\mathbf{B}}^0}^d(t)$ so obtained will be used to compute the decay rates of the neutral $\mathbf{B}$ mesons.

As already observed in Remark 2.10, it is convenient to perform all the computations in the basis of the eigenstates of H .

3.4 Invariance properties

Any symmetry transformation on our system can be modeled by a unitary or antiunitary transformation acting on the basis vectors $|\mathbf{B}^0\rangle$ and $|\bar{\mathbf{B}}^0\rangle$. In our formalism, we have to consider the corresponding transformation on the vector $|\hat{\rho}\rangle$, realized by a 4×4 matrix $\hat{\mathcal{U}}$:

$$|\hat{\rho}\rangle \rightarrow \hat{\mathcal{U}}|\hat{\rho}\rangle. \quad (3.21)$$

The request of invariance of the formalism under the transformation expressed by $\hat{\mathcal{U}}$ has the form

$$\hat{\mathcal{U}}(\hat{\mathcal{H}} + \hat{\mathcal{D}}')\hat{\mathcal{U}}^{-1} = (\hat{\mathcal{H}} + \hat{\mathcal{D}}'), \quad \hat{\mathcal{U}}(\hat{\mathcal{H}} + \hat{\mathcal{D}}')\hat{\mathcal{U}}^{-1} = (\hat{\mathcal{H}} + \hat{\mathcal{D}}')^\dagger \quad (3.22)$$

for unitary, respectively antiunitary, transformations.

For the discrete CPT, T and CP transformations the form of the matrix $\hat{\mathcal{U}}$ is

$$\hat{\mathcal{U}}_{CPT} = \begin{pmatrix} \sigma_1 & 0 \\ 0 & \sigma_0 \end{pmatrix}, \quad \hat{\mathcal{U}}_T = \begin{pmatrix} \sigma_0 & 0 \\ 0 & \sigma_\varphi \end{pmatrix}, \quad \hat{\mathcal{U}}_{CP} = \begin{pmatrix} \sigma_1 & 0 \\ 0 & \sigma_\varphi \end{pmatrix}, \quad (3.23)$$

where the σ_i are the Pauli matrices and the 2×2 identity and

$$\sigma_\varphi = \begin{pmatrix} 0 & e^{-2i\varphi} \\ e^{2i\varphi} & 0 \end{pmatrix} \quad (3.24)$$

contains the phase appearing in the CP transformation of the physical states, eq. (3.4). In the standard case $\hat{\mathcal{D}}' = 0$; the conditions of invariance under these discrete transformations reads

$$r_L = r_H \Rightarrow \theta = 0 \quad (3.25)$$

for the CPT symmetry,

$$|r_L r_H| = 1 \Rightarrow |\sigma| = 1, \xi = 0 \quad (3.26)$$

for the T symmetry; the CP invariance is obtained imposing both CPT and T invariance,

$$r_L = r_H = \sqrt{\sigma}, \sigma = e^{-2i\varphi}. \quad (3.27)$$

Remark 3.3 The parameters σ , ξ and θ characterize possible violations of the discrete symmetries during the time evolution of our system; they are called violations in mixing to distinguish them from violations of the same symmetries that can occur during the decay of the **B** mesons and which are due to the decay mechanism (see Section 3.5).

Also the dissipative part $\hat{\mathcal{D}}'$ can be responsible for violations of the discrete symmetries; choosing $\hat{\mathcal{H}} = 0$ the conditions of invariance are

$$c = b = 0 \quad (3.28a)$$

$$C' = e^{2i\varphi} C'^*, \quad B'^* = e^{4i\varphi} B' \quad (3.28b)$$

$$C' = -e^{2i\varphi} C'^*, \quad B'^* = e^{4i\varphi} B' \quad (3.28c)$$

holding for CPT, T and CP transformations respectively.

We consider now the invariance properties of our formalism under a redefinition of the phases of the two physical states. If the states $|\mathbf{B}^0\rangle$ and $|\bar{\mathbf{B}}^0\rangle$ undergo an independent change of phase,

$$|\mathbf{B}^0\rangle \rightarrow e^{i\phi} |\mathbf{B}^0\rangle, \quad |\bar{\mathbf{B}}^0\rangle \rightarrow e^{i\bar{\phi}} |\bar{\mathbf{B}}^0\rangle, \quad (3.29)$$

correspondingly the 4-vector $|\hat{\rho}\rangle$ undergoes the transformation described by

$$\mathcal{U}_\phi = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & e^{i(\phi-\bar{\phi})} & 0 \\ 0 & 0 & 0 & e^{-i(\phi-\bar{\phi})} \end{pmatrix}, \quad (3.30)$$

that causes a phase shift of the off-diagonal entries of the density matrix ρ :

$$\rho_3 \rightarrow e^{i(\phi-\bar{\phi})} \rho_3, \quad \rho_4 \rightarrow e^{-i(\phi-\bar{\phi})} \rho_4. \quad (3.31)$$

This transformation is not an invariance for $\hat{\mathcal{H}}$; indeed, the off-diagonal elements H_3 and H_4 of the effective Hamiltonian H do change. In particular r_H , r_L are found to transform as

$$r_L \rightarrow e^{i(\phi-\bar{\phi})} r_L, \quad r_H \rightarrow e^{i(\phi-\bar{\phi})} r_H; \quad (3.32)$$

consequently,

$$\sigma \rightarrow e^{2i(\phi-\bar{\phi})}\sigma \quad (3.33)$$

whereas ξ and θ are invariant. The phase redefinition is not an invariance also for $\hat{\mathcal{D}}'$: the dissipative coefficients are found to transform as

$$C' \rightarrow e^{-i(\phi-\bar{\phi})}C', \quad B' \rightarrow e^{2i(\phi-\bar{\phi})}B' \quad (3.34)$$

whereas a and A' are invariant. Then equation (3.11) is not invariant under $\mathcal{U}_\phi$, although the physically meaningful quantities (i.e. mean values of observables) are. Some problems can arise when this equation is solved by perturbative methods: the physically meaningful quantities could exhibit an explicit dependence on the phases ϕ and $\bar{\phi}$.

However, combining the results (3.32) and (3.34) it is possible to show that the entries of the matrix $\hat{\mathcal{D}}'^d$ are rephasing invariant. The same is true for the matrix $\hat{\mathcal{H}}^d$ since it is diagonal. These results ensure the invariance of our formalism under any phase redefinition, also for a truncated perturbative expansion of the time evolver. This is another advantage in the use of the eigenstates of the effective Hamiltonian as basis of the two-dimensional Hilbert space.

3.5 Observables

In this section we want to introduce the observables usually used to describe the physics of the neutral $\mathbf{B}$ mesons system. In the formalism of density matrices, these are represented by Hermitian operators O_f related to the decay of the $\mathbf{B}$ mesons into a generic final state f . In the $|\mathbf{B}^0\rangle, |\bar{\mathbf{B}}^0\rangle$ basis, these operators are represented by 2×2 matrices:

$$O_f = \begin{pmatrix} O_1 & O_3 \\ O_4 & O_2 \end{pmatrix} \quad (3.35)$$

whose entries, by construction, are related to the two independent decay amplitudes $\mathcal{A}(\mathbf{B}^0 \rightarrow f)$ and $\mathcal{A}(\bar{\mathbf{B}}^0 \rightarrow f)$:

$$\begin{aligned} O_1 &= |\mathcal{A}(\mathbf{B}^0 \rightarrow f)|^2, & O_3 &= [\mathcal{A}(\mathbf{B}^0 \rightarrow f)]^* \mathcal{A}(\bar{\mathbf{B}}^0 \rightarrow f), \\ O_2 &= |\mathcal{A}(\bar{\mathbf{B}}^0 \rightarrow f)|^2, & O_4 &= [\mathcal{A}(\bar{\mathbf{B}}^0 \rightarrow f)]^* \mathcal{A}(\mathbf{B}^0 \rightarrow f). \end{aligned} \quad (3.36)$$

with

$$\mathcal{A}(\mathbf{B}^0 \rightarrow f) \equiv \langle f|T|\mathbf{B}^0\rangle, \quad \mathcal{A}(\bar{\mathbf{B}}^0 \rightarrow f) \equiv \langle f|T|\bar{\mathbf{B}}^0\rangle, \quad (3.37)$$

T being the scattering operator.

Accordingly to (1.3), the mean value of O_f is obtained by a trace operation over the state describing the system at time t , regardless of the basis of the Hilbert space considered:

$$\langle O_f(t) \rangle = \text{Tr}[O_f \rho(t)] = \text{Tr}[O_f^d \rho^d(t)], \quad (3.38)$$

where O_f^d is the matrix representation of O_f in the $|\mathbf{B}_L\rangle, |\mathbf{B}_H\rangle$ basis: $O_f^d = R^\dagger O_f R$ and R as in (3.15). This mean value gives the probability rate for the decay of the **B** meson into the final state f .

In the analysis of the decays of neutral **B** mesons, it is customary to introduce the following rephasing invariant phenomenological parameters [57, 58]:

$$\begin{aligned} \lambda_L^f &\equiv \frac{q_L}{p_L} \frac{\mathcal{A}(\overline{\mathbf{B}}^0 \rightarrow f)}{\mathcal{A}(\mathbf{B}^0 \rightarrow f)}, & \mu_L^f &\equiv \frac{1}{\lambda_L^f}; \\ \lambda_H^f &\equiv \frac{q_H}{p_H} \frac{\mathcal{A}(\overline{\mathbf{B}}^0 \rightarrow f)}{\mathcal{A}(\mathbf{B}^0 \rightarrow f)}, & \mu_H^f &\equiv \frac{1}{\lambda_H^f}. \end{aligned} \quad (3.39)$$

By means of them the matrix O_f^d can be written as

$$\begin{aligned} O_f^d &= |\mathcal{A}(\mathbf{B}^0 \rightarrow f)|^2 \begin{pmatrix} |r_L|^2 |1 + \lambda_L^f|^2 & r_H r_L^* (1 - \lambda_H^f)(1 + \lambda_L^f)^* \\ r_L r_H^* (1 + \lambda_L^f)(1 - \lambda_H^f)^* & |r_H|^2 |1 - \lambda_H^f|^2 \end{pmatrix} \\ &= |\mathcal{A}(\overline{\mathbf{B}}^0 \rightarrow f)|^2 \begin{pmatrix} |1 + \mu_L^f|^2 & -(1 - \mu_H^f)(1 + \mu_L^f)^* \\ -(1 + \mu_L^f)(1 - \mu_H^f)^* & |1 - \mu_H^f|^2 \end{pmatrix}. \end{aligned} \quad (3.40)$$

A class of decays widely studied is the set of the so-called semileptonic decays, namely the decays where the final state f is constituted by a charged hadron h , a charged lepton l and its associated neutrino ν : $\mathbf{B}^0, \overline{\mathbf{B}}^0 \rightarrow h^- l^+ \nu, h^+ l^- \bar{\nu}$. In this case, it is convenient to parametrize the decay amplitudes as follows:

$$\mathcal{A}(\mathbf{B}^0 \rightarrow h^- l^+ \nu) = \mathcal{M}_h (1 - y_h), \quad (3.41a)$$

$$\mathcal{A}(\overline{\mathbf{B}}^0 \rightarrow h^+ l^- \bar{\nu}) = \mathcal{M}_h^* (1 + y_h^*), \quad (3.41b)$$

$$\mathcal{A}(\mathbf{B}^0 \rightarrow h^+ l^- \bar{\nu}) = z_h \mathcal{A}(\overline{\mathbf{B}}^0 \rightarrow h^+ l^- \bar{\nu}), \quad (3.41c)$$

$$\mathcal{A}(\overline{\mathbf{B}}^0 \rightarrow h^- l^+ \nu) = x_h \mathcal{A}(\mathbf{B}^0 \rightarrow h^- l^+ \nu), \quad (3.41d)$$

where $\mathcal{M}_h$ is a common factor. Possible violations of CPT invariance and of the $\Delta B = \Delta Q$ rule are parametrized by the coefficients x_h, y_h and z_h . In particular, non vanishing x_h and

z_h indicate violations of $\Delta B = \Delta Q$, because this rule would forbid the decays $\mathbf{B}^0 \rightarrow h^+ l^- \bar{\nu}$ and $\bar{\mathbf{B}}^0 \rightarrow h^- l^+ \nu$, while a non-vanishing y_h signals violation of CPT invariance, since this symmetry implies equal probability for the processes $\mathbf{B}^0 \rightarrow h^- l^+ \nu$ and $\bar{\mathbf{B}}^0 \rightarrow h^+ l^- \bar{\nu}$.

We assume the violations of the discrete symmetries (in mixing or in decay) and of the $\Delta B = \Delta Q$ rule to be small. In this way we can neglect all terms in which the coefficients $\theta, \xi, x_h, y_h, z_h$ are multiplied with the dissipative parameters a, b, c, α, β and γ . With this assumption, for the semileptonic decays, the parameters introduced in (3.39) assume a simplified form:

$$\lambda_L^{h^-} = \lambda_H^{h^-} = \sqrt{\sigma^*} x_h \equiv \lambda_h, \quad \mu_L^{h^+} = \mu_H^{h^+} = \sqrt{\sigma} z_h \equiv \mu_h. \quad (3.42)$$

Another class of decays we shall examine are characterized by a final state f that is an eigenstate of CP , with a definite CP parity ζ_f :

$$CP|f\rangle = \zeta_f|f\rangle, \quad \zeta_f = \pm 1. \quad (3.43)$$

At lowest order one finds $\lambda_{H,L}^f = \zeta_f$ [57].

3.6 Decay rates for a single **B** meson

In our formalism, the probability rate for the decay of an initial $\mathbf{B}^0$ or $\bar{\mathbf{B}}^0$ meson in a final state f at time t is given by

$$\mathcal{P}_f(\mathbf{B}^0; t) = \text{Tr}[O_f^d \rho_{\mathbf{B}^0}^d(t)], \quad \mathcal{P}_f(\bar{\mathbf{B}}^0; t) = \text{Tr}[O_f^d \rho_{\bar{\mathbf{B}}^0}^d(t)]. \quad (3.44)$$

Together with these probability rates, it is useful to define integrated rates as well:

$$\mathcal{P}_f(\mathbf{B}^0) = \frac{1}{\Gamma} \int_0^\infty \mathcal{P}_f(\mathbf{B}^0; \tau) d\tau, \quad \mathcal{P}_f(\bar{\mathbf{B}}^0) = \frac{1}{\Gamma} \int_0^\infty \mathcal{P}_f(\bar{\mathbf{B}}^0; \tau) d\tau. \quad (3.45)$$

As an example, we will explicitly list the probability rates for the semileptonic decays. It is convenient to define the parameters

$$\omega \equiv \frac{\Delta m}{\Gamma}, \quad \delta \equiv \frac{\Delta \Gamma}{2\Gamma} \quad (3.46)$$

where $\Delta m = m_H - m_L$, $\Gamma = (\gamma_H + \gamma_L)/2$ and $\Delta \Gamma = \gamma_L - \gamma_H$; it is also convenient to rescale the time as $\tau = t\Gamma$ (that is, we measure the time in units of **B** lifetime). Note that in the case of $\mathbf{B}_d$ mesons δ is expected to be small, of the order of 10^{-2} ; this allows to simplify some expressions in the coming sections.

Explicitely, one finds for the four relevant semileptonic decays $\mathbf{B}^0 \rightarrow h^- l^+ \nu$, $h^+ l^- \bar{\nu}$ and $\bar{\mathbf{B}}^0 \rightarrow h^- l^+ \nu$, $h^+ l^- \bar{\nu}$; for example,

$$\begin{aligned} \mathcal{P}_{h^-}(\mathbf{B}^0; \tau) = |\mathcal{M}_h|^2 e^{-\tau} \Big\{ & \sinh \delta \tau \left[\frac{|r_H|^2 - |r_L|^2}{|r_H + r_L|^2} - \text{Re } \lambda_h - \frac{\Lambda'}{\Delta \Gamma} \right] + \\ & + \cosh \delta \tau \left[\frac{|r_H|^2 + |r_L|^2}{|r_H + r_L|^2} - \text{Re } y_h + \frac{2}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Re } \Delta' \right] + \\ & + \sin \omega \tau \left[2 \frac{\text{Im}(r_H r_L^*)}{|r_H + r_L|^2} + \text{Im } \lambda_h - \frac{1}{2\Delta m} \text{Re } \Theta' \right] + \\ & + \cos \omega \tau \left[2 \frac{\text{Re}(r_H r_L^*)}{|r_H + r_L|^2} e^{-(\Omega' + \Lambda') \frac{\tau}{\Gamma}} - \text{Re } y_h - \frac{2}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Re } \Delta' \right] \Big\}; \end{aligned} \quad (3.47)$$

and the others are reported in Appendix C.

In these expressions we can identify the contributions coming from the standard evolution, from the hypothetical violations of CPT and $\Delta B = \Delta Q$ and from the dissipation. The probabilities $\mathcal{P}_h$ are directly accessible and can be studied in experiments performed at colliders: a fit of the time dependence of these quantities should permit, in line of principle, the evaluation of dissipative parameters Λ' , Δ' , Ω' and $\text{Re } \Theta'$. In the next section we shall define some quantities, called asymmetries, more suitable to extract informations about the dissipative parameters.

Analogously we can compute the probability rates for the decay of $\mathbf{B}^0$ and $\bar{\mathbf{B}}^0$ into final states with a definite CP parity, for example $\mathbf{D}^+ \mathbf{D}^-$, $\pi^+ \pi^-$, $\mathbf{J}/\Psi \mathbf{K}$. Their expressions are more involved and can be found in Appendix C.

3.7 Asymmetries for a single **B** meson

In order to reduce systematic errors, two experimental studies of the neutral **B** mesons system are usually performed using suitable combinations of the decay probabilities introduced in the previous section. They are called asymmetries and give a direct information about interesting physical quantities.

Given the process $\mathbf{B}^0 \rightarrow \mathbf{B}^0$, its CPT conjugate is $\bar{\mathbf{B}}^0 \rightarrow \bar{\mathbf{B}}^0$; given instead the process $\mathbf{B}^0 \rightarrow \bar{\mathbf{B}}^0$ its T conjugate is $\bar{\mathbf{B}}^0 \rightarrow \mathbf{B}^0$. Since, in absence of violation of the $\Delta B = \Delta Q$ rule, h^+ and h^- signal the decay of $\bar{\mathbf{B}}^0$ and $\mathbf{B}^0$ respectively, CPT and T violations are studied by the asymmetries

$$A_{CPT}(\tau) \equiv \frac{\mathcal{P}_{h^+}(\bar{\mathbf{B}}^0; \tau) - \mathcal{P}_{h^-}(\mathbf{B}^0; \tau)}{\mathcal{P}_{h^+}(\bar{\mathbf{B}}^0; \tau) + \mathcal{P}_{h^-}(\mathbf{B}^0; \tau)}, \quad (3.48)$$

respectively

$$A_T(\tau) \equiv \frac{\mathcal{P}_{h-}(\bar{\mathbf{B}}^0; \tau) - \mathcal{P}_{h+}(\mathbf{B}^0; \tau)}{\mathcal{P}_{h-}(\bar{\mathbf{B}}^0; \tau) + \mathcal{P}_{h+}(\mathbf{B}^0; \tau)}. \quad (3.49)$$

In the standard case (i.e. without dissipative evolution) they vanish if CPT and T are preserved. In our case, using the expressions in Appendix C, we find

$$\begin{aligned} A_{CPT}(\tau) = & \frac{(|r_L|^2 - |r_H|^2) \sinh \delta\tau + 2 \operatorname{Im}(r_L r_H^*) \sin \omega\tau}{(|r_L|^2 + |r_H|^2) \cosh \delta\tau + 2 \operatorname{Re}(r_L r_H^*) \cos \omega\tau} + 2 \operatorname{Re} y_h + \\ & + \frac{1}{\cosh \delta\tau + \cos \omega\tau} [\operatorname{Im}(\lambda_h - \mu_h) \sin \omega\tau + \operatorname{Re}(\lambda_h - \mu_h) \sinh \delta\tau] + \\ & - \frac{\cosh \delta\tau - \cos \omega\tau}{\cosh \delta\tau + \cos \omega\tau} \frac{4}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \operatorname{Re} \Delta' \end{aligned} \quad (3.50)$$

and

$$\begin{aligned} A_T(\tau) = & \frac{|r_L r_H|^2 - 1}{|r_L r_H|^2 + 1} - 2 \operatorname{Re} y_h + \\ & + \frac{1}{\cosh \delta\tau - \cos \omega\tau} \left\{ \sinh \delta\tau \left[\operatorname{Re}(\mu_h - \lambda_h) + \frac{4}{\Gamma} \frac{\omega}{\delta^2 + \omega^2} \operatorname{Im} \Delta' \right] + \right. \\ & \left. + \sin \omega\tau \left[\operatorname{Im}(\lambda_h - \mu_h) - \frac{4}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \operatorname{Im} \Delta' \right] \right\} \end{aligned} \quad (3.51)$$

We observe that these quantities are affected by the presence of dissipation, since they contain both the real and imaginary part of Δ' . However, in their expressions the dissipative parameters appear multiplied by the coefficient δ , that, as already noticed, is expected to be small; the dissipative effects are suppressed with respect to the standard contributions. As a consequence, these asymmetries are not in general suitable for probing the presence of dissipative effects in the neutral **B** mesons system.

To isolate the dissipative contribution, more complicated combinations of the probability rates are needed. For example, the asymmetry

$$A_{\Delta m}(\tau) \equiv \frac{[\mathcal{P}_{h-}(\mathbf{B}^0; \tau) - \mathcal{P}_{h-}(\bar{\mathbf{B}}^0; \tau)] - [\mathcal{P}_{h+}(\mathbf{B}^0; \tau) - \mathcal{P}_{h+}(\bar{\mathbf{B}}^0; \tau)]}{\mathcal{P}_{h-}(\mathbf{B}^0; \tau) + \mathcal{P}_{h-}(\bar{\mathbf{B}}^0; \tau) + \mathcal{P}_{h+}(\mathbf{B}^0; \tau) + \mathcal{P}_{h+}(\bar{\mathbf{B}}^0; \tau)} \quad (3.52)$$

used in the standard case to fit the mass parameter Δm , in our case presents a characteristic dependence on the dissipative terms; in the limit of a vanishing δ , it takes a very simple form:

$$A_{\Delta m}(\tau) = (\operatorname{Re} \theta)^2 (1 - \cos \omega\tau) + e^{-\Omega' \frac{\tau}{\Gamma}} \cos \omega\tau - \sin \omega\tau \left[\frac{1}{\Delta m} \operatorname{Re} \Theta' + \operatorname{Im}(\lambda_h + \mu_h) \right] \quad (3.53)$$

Assuming the validity of the $\Delta B = \Delta Q$ rule (that forbids the decays $\mathbf{B}^0 \rightarrow h^+ l^- \bar{\nu}$ and $\bar{\mathbf{B}}^0 \rightarrow h^- l^+ \nu$), and neglecting the quadratic dependence on the CPT-violating parameter θ , the different time behaviors in $A_{\Delta m}(\tau)$ should allow a determination of the non-standard parameters Ω' and the real part of Θ' . The actual accuracy of such a determination highly depends on the sensitivity of the measure of $A_{\Delta m}(\tau)$. From the results of available simulations one can expect an accuracy in the determination of the various terms in $A_{\Delta m}(\tau)$ of about a few percent [79,80]. This sensitivity is already enough to give interesting bounds on Ω' and $\text{Re } \Theta'$.

The assumption $\delta = 0$ has some important consequences. Indeed, in the limit $\Delta\Gamma = 0$ the non-Hermitian part Γ of the Weisskopf-Wigner Hamiltonian H becomes proportional to the identity. In this case the eigenstates of H , $|\mathbf{B}_H\rangle$ and $|\mathbf{B}_L\rangle$ becomes orthogonal, so $r_L r_H^* = 1$, producing $|\sigma| = 1$ (or $\xi = 0$) and, accordingly to (3.9), $\text{Im } \theta = 0$. Then T violation in mixing is forbidden and CPT violation in mixing is parametrized by $\text{Re } \theta$. This fact has important consequences for experimental tests of CPT and T invariance in mixing [81, 82, 83, 84, 85, 86].

Another interesting asymmetry is defined using the decay of both $\mathbf{B}^0$ and $\bar{\mathbf{B}}^0$ in an eigenstate f of CP :

$$A_f(\tau) \equiv \frac{\mathcal{P}_f(\mathbf{B}^0; \tau) - \mathcal{P}_f(\bar{\mathbf{B}}^0; \tau)}{\mathcal{P}_f(\mathbf{B}^0; \tau) + \mathcal{P}_f(\bar{\mathbf{B}}^0; \tau)}; \quad (3.54)$$

in the standard case, this observable is particularly suitable for studying violations of CP invariance. In presence of dissipation, we get

$$\begin{aligned} A_f(\tau) = & \frac{2\text{Re } \lambda_f}{1 + |\lambda_f|^2} \text{Re } \theta + \frac{2\zeta_f}{\omega\Gamma} \text{Im } \Delta' + \\ & - \sin \omega\tau \left[\frac{2\text{Im } \lambda_f}{1 + |\lambda_f|^2} + \frac{2\zeta_f}{\omega\Gamma} \text{Re } \Delta' \right] + \\ & + \cos \omega\tau \left[\frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2} - \frac{2\text{Re } \lambda_f}{1 + |\lambda_f|^2} \text{Re } \theta - \frac{2\zeta_f}{\omega\Gamma} \text{Im } \Delta' \right], \end{aligned} \quad (3.55)$$

where the condition $\delta \approx 0$ has been used; this enables us to write λ_H^f and λ_L^f in terms of a unique amplitude ratio λ_f :

$$\lambda_f \equiv \sqrt{\sigma^*} \frac{\mathcal{A}(\bar{\mathbf{B}}^0 \rightarrow f)}{\mathcal{A}(\mathbf{B}^0 \rightarrow f)} \quad (3.56)$$

and

$$\lambda_L^f = \lambda_f(1 - \text{Re } \theta), \quad \lambda_H^f = \lambda_f(1 + \text{Re } \theta). \quad (3.57)$$

The presence of dissipation in (3.55) manifests itself in the modification of the coefficients of the oscillating terms and in the presence of a τ -independent piece. Dedicated **B**

experiments at colliders should be able to identify the various τ -dependence and therefore provide stringent bounds on the parameter Δ' .

Asymmetries analogous to the ones just discussed can be constructed using integrated rates, see eq. (3.46), rather than the probability rates. The quantities we obtain in this way are time independent and are denoted with a prime index to distinguish them from the previously defined ones. Two examples are

$$A'_{\Delta m} = \frac{1}{1 + \omega^2} \left\{ 1 + \omega^2 (\text{Re } \theta)^2 - \omega \left[\frac{1}{\Delta m} \text{Re } \Theta' + \text{Im}(\lambda_h + \mu_h) \right] + \frac{1}{\Gamma(1 + \omega^2)} [(\omega^2 - 1)\Omega' + 2\omega^2\Lambda'] \right\} \quad (3.58)$$

and

$$A'_f = \frac{2 \text{Re } \lambda_f}{1 + |\lambda_f|^2} \text{Re } \theta + \frac{2\zeta_f}{\omega\Gamma} \text{Im } \Delta' + \frac{1}{1 + \omega^2} \left\{ \omega \left[\frac{2 \text{Im } \lambda_f}{1 + |\lambda_f|^2} + \frac{2\zeta_f}{\omega\Gamma} \text{Re } \Delta' \right] + \frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2} + \frac{2 \text{Re } \lambda_f}{1 + |\lambda_f|^2} \text{Re } \theta + \frac{2\zeta_f}{\omega\Gamma} \text{Im } \Delta' \right\}. \quad (3.59)$$

The result in equation (3.58) is particularly interesting: assuming the validity of the $\Delta B = \Delta Q$ rule and CPT-invariance, a measure of $A'_{\Delta m}(1 + \omega^2)$ not compatible with 1 would signal the presence of non standard effects. Such a test can be performed at colliders with high accuracy, in particular using the next generation dedicated **B** mesons experiments.

The integrated asymmetry (3.59) is not very useful in probing the presence of dissipative effects in neutral **B** mesons decays: since there is no time-dependence, the measure of this asymmetry would give little information on the magnitude of Δ' .

3.8 Decay rates for entangled **B** meson

In the analysis of neutral **B** mesons physics, the study of pairs of mesons produced at the so-called **B**-factories, that is e^+e^- colliders tuned at the energy of the $\Upsilon(4S)$ resonance, is of particular relevance. The pairs of **B** mesons are produced by the decay of this resonance,

$$\Upsilon(4S) \rightarrow \mathbf{B}^0 \bar{\mathbf{B}}^0, \quad (3.60)$$

in an entangled state. Since $\Upsilon(4S)$ has spin 1 whereas the two **B** mesons are spinless, they are produced, in the $\Upsilon(4S)$ rest frame, in an antisymmetric spatial state. In this

reference frame $\mathbf{B}^0$ and $\bar{\mathbf{B}}^0$ are flying apart with opposite momenta $\vec{p}$ and $-\vec{p}$; the state encoding this configuration is the singlet state:

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|\mathbf{B}^0, -\vec{p}\rangle \otimes |\bar{\mathbf{B}}^0, \vec{p}\rangle - |\bar{\mathbf{B}}^0, -\vec{p}\rangle \otimes |\mathbf{B}^0, \vec{p}\rangle), \quad (3.61)$$

or, in the density matrix formalism, the operator $\rho = |\psi\rangle\langle\psi|$, in this case a 4×4 matrix. Its irreversible time evolution is assumed to have a factorized form:

$$\rho(t_1, t_2) = (\Lambda_{t_1} \otimes \Lambda_{t_2})[\rho(0)], \quad (3.62)$$

where we consider two possibly different proper times t_1 and t_2 for the two mesons since in general they won't decay at the same time.

Remark 3.4 In the standard treatment of the neutral **B** mesons system the time-evolution is assumed to have the factorized form (3.62) since there is not interaction between the two particles in the pair; in the open system approach an interaction can be induced by the common thermal bath, but it disappears at the lowest order in the dissipative effects; whence the hypothesis of factorization.

As in the previous sections, we prefer to express the statistical operator in the basis diagonalizing the Hamiltonian contribution, then

$$\rho^d(t_1, t_2) = (R^{-1} \otimes R^{-1})\rho(t_1, t_2)(R^{-1\dagger} \otimes R^{-1\dagger}), \quad (3.63)$$

which in terms of the basis $\{P_i, i = 1, \dots, 4\}$ introduced in Section 2.1, reads

$$\rho^d(t_1, t_2) = \frac{1}{2|r_L + r_H|^2}(P_1(t_1) \otimes P_2(t_2) + P_2(t_1) \otimes P_1(t_2) - P_3(t_1) \otimes P_4(t_2) - P_4(t_1) \otimes P_3(t_2)), \quad (3.64)$$

where $P_i(t) = \Lambda_t[P_i]$.

The probability that a meson decays into a final state f_1 at proper time t_1 while the other meson decays into the final state f_2 at proper time t_2 is given by

$$\mathcal{P}(f_1, t_1; f_2, t_2) = \text{Tr}[(O_{f_1}^d \otimes O_{f_2}^d)\rho^d(t_1, t_2)] \quad (3.65)$$

where $O_{f_1}^d$ and $O_{f_2}^d$ are the 2×2 matrix representations, in the basis of eigenstates of H , of the Hermitian operators O_{f_1} and O_{f_2} described in Section 3.5.

Since the neutral **B** lifetime is very short, it is difficult to identify the two proper times t_1 and t_2 : the probability (3.65) is experimentally unattractive. It is thus customary to define integrated rates, as in the case of the decays of a single meson. In particular it is possible to consider a fixed time delay $t = t_1 - t_2$ between the two decays, obtaining a one-time probability rate:

$$\mathcal{P}(f_1, f_2; t) \equiv \int_0^\infty \mathcal{P}(f_1, t' + t; f_2, t') dt' \quad (3.66)$$

when $t \geq 0$, that is when the first meson decays after the second, while, for $t < 0$, one defines

$$\mathcal{P}(f_1, f_2; t) \equiv \int_0^\infty \mathcal{P}(f_1, t' + t; f_2, t') \theta(t' + t) dt' \quad (3.67)$$

where the θ of Heavyside is required since the evolution is of semigroup type, with forward in time propagation. This definition implies

$$\mathcal{P}(f_1, f_2; -t) = \mathcal{P}(f_2, f_1; t). \quad (3.68)$$

In the following we shall always assume $t \geq 0$. It is also possible to define a time independent integrated rate as

$$\mathcal{P}(f_1, f_2) = \int_0^\infty \mathcal{P}(f_1, f_2; t) dt \quad (3.69)$$

in which we sum over all possible delays between the two decays. This quantity is relevant in the experimental analysis at the so-called symmetric **B** factories where the $\Upsilon(4S)$ decays practically at rest in the laboratory frame, and then the time information is hardly attainable. On the other side, at the so-called asymmetric **B** factories the decay of the $\Upsilon(4S)$ resonance takes place in a boosted frame respect to the laboratory; in this case the temporal information can be extracted and the one-time probability (3.66) is suitable to the data analysis.

The explicit expressions for these joint probabilities are rather involved; some particularly interesting examples are collected in Appendix C.

One can nevertheless make a general remark. Consider $\mathcal{P}(f, f; 0)$, i.e. the probability rate for the simultaneous decay of the two **B** mesons in the same final state f . Since the initial state is antisymmetric, the usual dynamics produces $\mathcal{P}(f, f; 0) = 0$ whereas the modified dynamics in general gives $\mathcal{P}(f, f; 0) \neq 0$. This quantity could then unambiguously show the presence of non standard effects.

3.9 Asymmetries for entangled **B** meson

As in the study of the decays of single **B** mesons, the experimental analysis of our system is not usually performed directly with the probability rates introduced before; rather, some suitable asymmetries are constructed. We use the assumption $\delta = 0$ to simplify our expressions and, as before, we refer to a rescaled time scale $\tau = t\Gamma$.

An interesting quantity involving semileptonic decays in both mesons is

$$\mathcal{R}(\tau) \equiv \frac{\mathcal{P}(h^+, h^+; \tau) + \mathcal{P}(h^-, h^-; \tau)}{\mathcal{P}(h^+, h^-; \tau) + \mathcal{P}(h^-, h^+; \tau)}. \quad (3.70)$$

In the given approximation, one finds

$$\begin{aligned} \mathcal{R}(\tau) = & \frac{(1 + |r_H r_L|^2)(\cosh \delta\tau)}{(|r_L|^2 + |r_H|^2) \cosh \delta\tau + 2 \operatorname{Re}(r_L r_H^*) \cos \omega\tau} + \\ & + \frac{2}{(1 + \cos \omega\tau)^2} \left[(1 + \tau) \frac{\Omega'}{\Gamma} \cos \omega\tau + \frac{1}{\Delta m} (\sin \omega\tau + \omega \cos \omega\tau) \operatorname{Re} \left(\frac{\Theta'}{1 - i\omega} \right) \right], \end{aligned} \quad (3.71)$$

and its evaluation for simultaneous decays is particularly meaningful:

$$\mathcal{R}(0) = \frac{1}{2} \left[\frac{\Omega'}{\Gamma} - \frac{1}{\Delta m} \frac{\omega}{1 + \omega^2} (\omega \operatorname{Im} \Theta' - \operatorname{Re} \Theta') \right], \quad (3.72)$$

since in the standard case $\mathcal{R}(0) = 0$. Therefore a fit of $\mathcal{R}(\tau)$ on actual experimental data that produces a non-zero intercept would clearly signal the presence of non-standard effects.

Another quantity built with the double semileptonic decays and relevant in the study of dissipative effects is the asymmetry $\mathcal{A}_{\Delta m}(\tau)$, analogous to the one defined in (3.52):

$$\mathcal{A}_{\Delta m}(\tau) \equiv \frac{[\mathcal{P}(h^+, h^-; \tau) + \mathcal{P}(h^-, h^+; \tau)] - [\mathcal{P}(h^+, h^+; \tau) + \mathcal{P}(h^-, h^-; \tau)]}{\mathcal{P}(h^+, h^+; \tau) + \mathcal{P}(h^-, h^-; \tau) + \mathcal{P}(h^+, h^-; \tau) + \mathcal{P}(h^-, h^+; \tau)}. \quad (3.73)$$

Its explicit expression is particularly simple and meaningful, even in the case of vanishing δ :

$$\begin{aligned} \mathcal{A}_{\Delta m}(\tau) = & \frac{\Gamma}{\Gamma + \Omega'} e^{-\Omega' \frac{\tau}{\Gamma}} \cos \omega\tau + (\operatorname{Re} \theta)^2 (1 - \cos \omega\tau) + \\ & - (\sin \omega\tau + \frac{1}{\Delta m} \omega \cos \omega\tau) \operatorname{Re} \left(\frac{\Theta'}{1 - i\omega} \right). \end{aligned} \quad (3.74)$$

Thanks to the different time behavior, an experimental study of this observable should allow to extract precise information on Ω' and the combination $\omega \operatorname{Im} \Theta' - \operatorname{Re} \Theta'$. The asymmetry $\mathcal{A}_{\Delta m}$ is intensively studied at **B**-factories [87, 88, 89] since it allows a precise determination of the mass difference Δm . Indeed, $\mathcal{A}_{\Delta m}(\tau) = \cos \omega\tau$ in absence of dissipation.

Another asymmetry interesting from a dissipative point of view is built with one-time integrated rates and it involves a semileptonic channel and an hadronic channel (in particular a state with definite CP -parity):

$$\mathcal{A}_f(\tau) \equiv \frac{\mathcal{P}(h^+, f; \tau) - \mathcal{P}(h^-, f; \tau)}{\mathcal{P}(h^+, f; \tau) + \mathcal{P}(h^-, f; \tau)}. \quad (3.75)$$

If we neglect possible violations of the $\Delta B = \Delta Q$ rule, the semileptonic decay identify the flavor of the decaying meson. In this case this asymmetry is equivalent to the $A_f(\tau)$ defined in (3.54). If the final state is $f = \mathbf{J}/\Psi \mathbf{K}$ these observables are directly expressed in terms of one of the angles that parametrize the CP violation in the Cabibbo Kobayashi Maskawa matrix; for this reason they are extensively studied. The expression of $\mathcal{A}_f(\tau)$ in the case of irreversible dynamics is given by:

$$\begin{aligned} \mathcal{A}_f(\tau) = & \frac{2 \operatorname{Re} \lambda_f}{1 + |\lambda_f|^2} \operatorname{Re} \theta + \zeta_f \operatorname{Re}(\lambda_h - \mu_h + 2\zeta_f y_h) - \frac{2\zeta_f}{\omega\Gamma} \operatorname{Im} \Delta' + \\ & + \cos \omega\tau \left[\frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2} - \frac{2 \operatorname{Re} \lambda_f}{1 + |\lambda_f|^2} \operatorname{Re} \theta - \frac{2\zeta_f}{\omega\Gamma} \operatorname{Re} \left(\frac{\omega + 2i}{2 + i\omega} \Delta' \right) \right] + \\ & + \sin \omega\tau \left[\frac{2 \operatorname{Im} \lambda_f}{1 + |\lambda_f|^2} - \frac{2\zeta_f}{\omega\Gamma} \operatorname{Im} \left(\frac{\omega + 2i}{2 + i\omega} \Delta' \right) \right] \end{aligned} \quad (3.76)$$

where λ_f is defined in (3.56) and ζ_f in (3.43); for simplicity in this formula only first order terms in CPT violating parameters are kept. Equation (3.76) holds for $\tau \geq 0$; for $\tau < 0$ the expression is

$$\begin{aligned} \mathcal{A}_f(\tau) = & \frac{2 \operatorname{Re} \lambda_f}{1 + |\lambda_f|^2} \operatorname{Re} \theta + \zeta_f \operatorname{Re}(\lambda_h - \mu_h + 2\zeta_f y_h) - \frac{2\zeta_f}{\omega\Gamma} \operatorname{Re} \left(\frac{\omega + 2i}{2 + i\omega} \Delta' \right) + \\ & + \cos \omega\tau \left[\frac{1 - |\lambda_f|^2}{1 + |\lambda_f|^2} - \frac{2 \operatorname{Re} \lambda_f}{1 + |\lambda_f|^2} \operatorname{Re} \theta - \frac{2\zeta_f}{\omega\Gamma} \operatorname{Im} \Delta' \right] + \\ & + \sin \omega\tau \left[\frac{2 \operatorname{Im} \lambda_f}{1 + |\lambda_f|^2} + \frac{2\zeta_f}{\omega\Gamma} \operatorname{Re} \Delta' \right]; \end{aligned} \quad (3.77)$$

the two expressions coincide for $\tau = 0$ and, in absence of dissipation, one can be obtained from the other by letting $\tau \rightarrow -\tau$. The fact that this is not longer true when $\Delta' \neq 0$ is a signal of irreversibility.

Assuming the validity of the $\Delta B = \Delta Q$ rule and neglecting CPT-violating effects in mixing ($\theta = 0$) as well as direct CP violations (i.e. $|\lambda_f| = 1$, see [57]), a fit of (3.76) with experimental data allows the determination of both $\operatorname{Re} \Delta'$ and $\operatorname{Im} \Delta'$. The study of $\mathcal{A}_f$ could result in one of the best tests on the presence of dissipative effects in **B** physics, in particular for the final state $\mathbf{J}/\Psi \mathbf{K}$ [87].

The quantities involving the fully integrated probability rates, eq. (3.69), in general are not suitable to study dissipative effects, since the dissipative parameters are multiplied by δ ; nevertheless it is possible to define two ratios unaffected by this problem:

$$\chi_B \equiv \frac{\mathcal{P}(h^+, h^+) + \mathcal{P}(h^-, h^-)}{\mathcal{P}(h^+, h^+) + \mathcal{P}(h^-, h^-) + \mathcal{P}(h^+, h^-) + \mathcal{P}(h^-, h^+)} \quad (3.78)$$

and

$$R_B \equiv \frac{\mathcal{P}(h^+, h^+) + \mathcal{P}(h^-, h^-)}{\mathcal{P}(h^+, h^-) + \mathcal{P}(h^-, h^+)}. \quad (3.79)$$

Assuming the $\Delta B = \Delta Q$ rule, χ_B represents the total $\mathbf{B}^0$ - $\overline{\mathbf{B}}^0$ mixing probability. In our framework, one explicitly finds:

$$\chi_B = \frac{\omega^2}{2(1 + \omega^2)} \left[1 - (\text{Re } \theta)^2 + \frac{2}{\omega^2(1 + \omega^2)} X \right], \quad (3.80)$$

$$R_B = \frac{\omega^2 [1 - (\text{Re } \theta)^2]}{2 + \omega^2 [1 + (\text{Re } \theta)^2]} + \frac{4}{(2 + \omega^2)^2} X; \quad (3.81)$$

the dissipative contribution is the same for the two observables and it is given by

$$X \equiv \frac{1}{\Gamma} (\Omega' - \omega^2 \Lambda') - \frac{\omega}{\Delta m} (\omega \text{Im } \Theta' - \text{Re } \Theta'). \quad (3.82)$$

Independent measures of these two quantities would provide a way to estimate both $\text{Re } \theta$ and X , and therefore give limits on both dissipative and CPT-violating effects.

3.10 Discussion and conclusions

In this chapter we have treated the neutral **B** mesons system in the framework of quantum dynamical semigroups. The formalism is very general and can be applied to describe the time evolution of this system when it is subject to a weak interaction with an external environment. As described in Section 3.2, quantum gravity could give a motivation for such an approach, since in this case the space-time itself would act as an effective environment for any physical system.

As discussed before, the open system framework allows a rough evaluation of magnitude of the dissipative parameters; they scale at most as $m_{\mathbf{B}}^2/M_F$, where $m_{\mathbf{B}}$ is the meson mass while M_F is a large fundamental mass scale. Typically, M_F coincides with the Planck scale so that $m_{\mathbf{B}}^2/M_F \sim 10^{-18}$ GeV. This value is very small; however, the sophistication of the dedicated **B**-experiments, both at colliders (CDF-II, HERA-B, BTeV, LHC-*b*) and **B**-factories (BaBar, Belle, CLEO-III), is so high that the sensitivity required to probe the presence of non-vanishing dissipative parameters can be reached in just a few years of data taking [79, 80, 87, 88, 89].

Remark 3.5 An evaluation of the dissipative parameters for a gravitational bath have been already performed in the context of the neutral **K** mesons physics [55, 90]. Upper bounds for the dissipative parameters have been obtained; these are in the range of what suggested by gravitational or stringy effects. More data are expected from ongoing experiments at the so-called Φ -factories, the analogous of **B**-factories for the **K** mesons.

To conclude, we observe that, besides the study of irreversible dynamics, these experiments, involving correlated mesons, could give the possibility to investigate the mathematical notion of complete positivity from a phenomenological point of view. Let us consider a simplified situation with a trivial Hamiltonian sector, $\mathcal{H} = 0$, and a dissipative part with vanishing a , b and c . A dissipative contribution of this kind has been used to describe non-standard effects in the $\mathbf{K}$ meson system, using a positive but not completely positive dynamics [69, 77, 78]. The eigenvalues of the evolved singlet state in (3.62), for $t_1 = t_2$, are

$$\begin{aligned}\lambda_{1,2} &= 2 \pm \{[E_+(t)]^2 + [E_-(t)]^2 + 2[F(t)]^2\}, \\ \lambda_{3,4} &= \pm[E_+(t) + E_-(t)]\{[E_+(t) - E_-(t)]^2 + 4[F(t)]^2\}^{1/2},\end{aligned}\tag{3.83}$$

where

$$\begin{aligned}E_{\pm}(t) &= \frac{1}{\nu_+ - \nu_-}[(\nu_+ + 2\alpha)e^{\nu_{\pm}t} - (\nu_- + 2\alpha)e^{\nu_{\mp}t}], \\ F(t) &= \frac{2\beta}{\nu_+ - \nu_-}[e^{\nu_-t} - e^{\nu_+t}],\end{aligned}\tag{3.84}$$

with $\nu_{\pm} = -(\alpha + \gamma) \pm \sqrt{(\alpha - \gamma)^2 + 4\beta^2}$.

The condition of positivity of this dynamics reads $\alpha\gamma \geq \beta^2$ whereas complete positivity requires $\alpha = \gamma$ and $\beta = 0$. Consequently, in the simply positivity case $\nu_{\pm}$ are both negative and thus, for $t \neq 0$ two of the eigenvalues (3.83) are always negative and the statistical interpretation of the density matrix $\rho(t, t)$ is compromised. In the completely positive case we find $\nu_+ = \nu_-$ and all the considered eigenvalues remain non negative for any time.

These results offer a further motivation for studying generalized dynamics of the form (3.10) at mesons factories. These set-ups are in fact high-performance quantum interferometers: at least in principle, they can verify from the experimental point of view the role of the condition of complete positivity in the time evolution of correlated mesons [44, 45, 46, 47, 48, 49]. As shown by the simple example above, this is not just a mere technical question: it is crucial for our physical understanding of the quantum dynamics of open systems.

Chapter 4

Neutrinos

Neutrino physics is presently a very active area of research. Recent experiments suggest neutrinos to be massive and the three flavors, electronic (ν_e), muonic (ν_μ) and tauonic (ν_τ) neutrinos, to mix among themselves. Consequently the Standard Model should be modified by the introduction of a mixing matrix in the leptonic sector (analogous to the Cabibbo Kobayashi Maskawa matrix in the hadronic sector). The first hint to the possibility of neutrino mixing came from the observation of a reduced ν_e flux from the Sun, incompatible with the usual model of solar evolution. The most favoured explanation of this phenomenon is the so-called MSW effect which is deduced from the hypothesis that electronic neutrinos interact with solar matter [91,92,93]. In this context, flying neutrinos behave as open systems in interaction with an environment (the Sun), but the MSW effect is only the Lamb-shift term (a redefinition of the neutrino effective Hamiltonian) and no dissipative effects appear.

In other approaches, dissipation has been taken into account [94,95,96,97,98,99,100,101,102,103,104,105], though paying scarce attention to internal consistency. In this chapter, following the lines introduced in Chapters 1 and 2, we will apply the rigorous formalism of open quantum systems to describe neutrino time-evolutions in matter, with particular care to the derivation of a Markovian reduced dynamics. In this way, we will get results unaffected from physical inconsistencies that otherwise can appear [99].

We shall consider an environment with constant mean density and derive the general form of the generator of the time-evolution in the Markovian approximation. Since, in the standard description of the MSW effect, it is the variation of this mean density that produces the most relevant aspects in the treatment of the flavors oscillations [92,93], we shall describe how to generalize our formalism to account for the case of environments with a varying density profile.

The transitions probabilities are explicitly written out in two different situations: a flavor-changing interaction and a flavor-preserving one (the case of solar matter).

4.1 The physics of neutrinos in vacuum

In this section we shortly outline the principal facts of neutrino physics in vacuum. For sake of simplicity, we consider the case of two neutrino's flavors, ν_e and ν_μ , and we describe the neutrino system by means of the bidimensional Hilbert space of flavor, $\mathfrak{H}_2$ [106, 107, 108]; indeed, in vacuum the neutrino momentum can be kept fixed and the related degrees of freedom does not appear in the description. This is the standard phenomenological approach to this system and the experimental data are usually analyzed in this context. This approach is justified by more rigorous treatments; see [110, 111].

As in the case of the neutral **B** mesons system we distinguish between two basis for $\mathfrak{H}_2$:

- the basis of physical states (or flavor states), $\{|\nu_e\rangle, |\nu_\mu\rangle\}$; these are the states that are actually detected in experiments;
- the basis of the eigenstates of the Hamiltonian of the system H_S , $\{|\nu_1\rangle, |\nu_2\rangle\}$, characterized by

$$\begin{cases} H_S|\nu_1\rangle = \epsilon'_1|\nu_1\rangle, \\ H_S|\nu_2\rangle = \epsilon'_2|\nu_2\rangle, \end{cases} \quad (4.1)$$

where the H_S is Hermitian since neutrinos are stable particles and $\epsilon'_{1,2}$ are real eigenvalues.

H_S generates the time-evolution of eigenstates in vacuum:

$$|\nu_i(t)\rangle = e^{-iH_S t}|\nu_i\rangle = e^{-i\epsilon'_i t}|\nu_i\rangle \quad (4.2)$$

and the eigenvalues ϵ'_i are the energies of these states, that depend on the neutrino momentum p and mass m_i . Indeed, in agreement with experimental evidence [109], the states ν_1 and ν_2 are assumed to be massive and we denote with m_1 , respectively m_2 , their masses. In the Standard Model they vanish; for later reference we assume they are not degenerate.

Since m_1 and m_2 are believed to be very small, ϵ'_i can be expanded in the ultrarelativistic limit, i.e. neglecting¹ the terms of higher order than m_1^2 and m_2^2 :

$$\epsilon'_i = \sqrt{p^2 + m_i^2} \approx p + \frac{m_i^2}{2p} \approx p + \frac{m_i^2}{2E}, \quad (4.3)$$

where $E \approx p$ is the neutrino energy.

The two basis are connected, in vacuum, by an unitary transformation U_0 (the so-called mixing matrix) that can be chosen to be orthogonal by a suitable redefinition of the states $|\nu_1\rangle$ and $|\nu_2\rangle$.

¹These terms are supposed smaller than the dissipative effects that we shall consider in the coming sections

Remark 4.1 The choice of an orthogonal U_0 matrix is a consequence of the impossibility to violate the CP symmetry in a two flavors context: in fact the CP violation is related to the existence of unavoidable complex phases in the mixing matrix and they appear necessarily in a three flavors context. However, when dealing with dissipative dynamics, in general the complex phase appearing in U_0 is relevant also in the two flavors context, because it produces physical consequences [117]. In our context, however, this phase can be consistently reabsorbed; indeed, the particular dissipative effects we shall consider in what follows does not produce any evidence of this phase. Therefore, in the following, we do not account for possible violations of CP in the leptonic sector.

We have:

$$\begin{cases} |\nu_e\rangle = U_0|\nu_1\rangle, \\ |\nu_\mu\rangle = U_0|\nu_2\rangle; \end{cases} \quad U_0 = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}; \quad (4.4)$$

where the parameter θ is the mixing angle in vacuum between the two families considered. The Hamiltonian H_S in the basis above reads:

$$H_S^d = \begin{pmatrix} \epsilon_1 & 0 \\ 0 & \epsilon_2 \end{pmatrix}, \quad H_S = \frac{\Delta m^2}{4E} \begin{pmatrix} -\cos 2\theta & \sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix} \quad (4.5)$$

where² $H_S = U_0 H_S^d U_0^{-1}$. Since we prefer to deal with a traceless matrix H_S^d , we have redefined the eigenvalues of H_S as $\epsilon_1 = -\Delta m^2/4E$ and $\epsilon_2 = \Delta m^2/4E$, where $\Delta m^2 \equiv m_2^2 - m_1^2$ and E is the neutrino energy.

The distinction between the two basis of the Hilbert space produces the phenomenon of flavor oscillations, i.e. periodical transitions between ν_e and ν_μ .

Since a physical state is given by a superposition of eigenstates of H_S , an initially pure state $|\nu_e\rangle$ will present, for $t > 0$, a non-zero component along $|\nu_\mu\rangle$. For instance

$$|\nu_e(t)\rangle = \left(e^{i\Delta m^2 t/4E} \cos^2 \theta + e^{-i\Delta m^2 t/4E} \sin^2 \theta \right) |\nu_e\rangle + \left(i \sin 2\theta \sin \frac{\Delta m^2}{4E} t \right) |\nu_\mu\rangle. \quad (4.6)$$

The probability of transition (in vacuum) is given by [106, 107, 108]:

$$P_{\nu_e \rightarrow \nu_\mu}(t) = |\langle \nu_\mu | \nu_e(t) \rangle|^2 = \sin^2 2\theta \sin^2 \frac{\Delta m^2}{4E} t \quad (4.7)$$

and the surviving probability is

$$P_{\nu_e \rightarrow \nu_e}(t) = 1 - P_{\nu_e \rightarrow \nu_\mu}(t). \quad (4.8)$$

Vacuum neutrino oscillations were the first explanation of the lack of ν_e in the solar flux.

²We use the notation introduced in Chapter 2 to characterize the operators expressed in this basis

Remark 4.2 This phenomenon should lead to lepton numbers violations, forbidden in the Standard Model.

However, the transition probabilities are modified by the interactions of neutrinos in matter [91, 130]; in a homogeneous medium a matter mixing angle and two redefined masses are needed and the probabilities of transition are formally equal to (4.7) and (4.8) but with the redefined parameters θ_m and Δm_m^2 . These transition probabilities are used to fit the parameters θ and Δm^2 to the experimental data [106, 107, 108].

In what follows, we shall describe neutrinos in flavor space by means of statistical operators since we are interested in decohering evolutions. In particular, $\rho_{\nu_e} = |\nu_e\rangle\langle\nu_e|$ and $\rho_{\nu_\mu} = |\nu_\mu\rangle\langle\nu_\mu|$, whence (4.7) reads

$$P_{\nu_e \rightarrow \nu_\mu}(t) = \text{Tr}[\rho_{\nu_e}(t)\rho_{\nu_\mu}]. \quad (4.9)$$

We shall also adopt the vector representation, developed in Chapter 2, for the density matrices.

4.2 The model of interaction

In the previous section we have described the behavior of neutrinos in vacuum. Here we consider the case of neutrinos interacting with a suitable environment, consisting of a slab of matter of volume V . The Hamiltonian describing both neutrinos and environment has the form $H_T = H_S + H_B + H_I$, where H_S is the neutrino Hamiltonian (4.5), H_B is the Hamiltonian of the environment, and H_I is the contribution containing the interaction between neutrinos and environment. In this section we shall derive its form.

Remark 4.3 In order to have a general H_I , it is not possible to consider a two-dimensional $\mathfrak{H}_S$, as in vacuum; momentum dependence needs also to be taken into account. Nevertheless, we shall see that, under certain circumstances, a two-dimensional effective description is available also in the open system context.

The interaction H_I is determined by a Hamiltonian density $\mathfrak{h}$ by

$$H_I(t) = \int_V \mathfrak{h}(x) d\vec{x}, \quad (4.10)$$

where $x \equiv (\vec{x}, t)$ (the dependence on time in H_I disappears if we suppose the layer of matter to be uniform). We assume a linear coupling of the form

$$\mathfrak{h}(x) = g \sum_{\alpha} V_S^{\alpha}(x) \otimes V_B^{\alpha}(x), \quad (4.11)$$

with

$$V_S^{\alpha}(x) \equiv V_S^{lm\mu}(x) = \overline{\nu}_l(x) \gamma_{\mu} (1 - \gamma_5) \nu_m(x), \quad (4.12)$$

where g is a suitable coupling constant (dimensionally it is an inverse energy squared), $V_{S,B}^\alpha(x)$ are the neutrino and bath currents that characterize the interactions and α is a set of flavor and Dirac indices. The γ_μ 's, $\mu = 0, \dots, 3$ are the Dirac matrices and $\gamma_5 = i\gamma_0\gamma_1\gamma_2\gamma_3$ while $\nu_l(x)$, $l = e, \mu$, are the fermionic neutrino fields. To guarantee the Hermiticity of H_I we require

$$V_B^{lm\mu\dagger}(x) = V_B^{ml\mu}(x), \quad (4.13)$$

otherwise, $V_B^\alpha(x)$ is completely arbitrary. The indices S and B denote the neutrino system and the environment respectively, according to the notation of Chapter 1.

Remark 4.4 The assumption (4.12) is quite general; for instance, it includes transitions forbidden in the Standard Model, like neutrinos flavor-changing interactions [112, 113, 114, 115]. These latter are induced by suitable environment operators $V_B^{lm\mu}$ with $l \neq m$.

Remark 4.5 To describe neutrino interactions in the Sun, a low-energy Hamiltonian density is actually introduced; it contains neutral currents and charged currents contributions in the point-interaction approximation (Fermi theory) to the Standard Model [108]:

$$\begin{aligned} \mathfrak{h}(x) = & \frac{G_F}{\sqrt{2}} \sum_l \sum_\mu \bar{\nu}_l(x) \gamma_\mu (1 - \gamma_5) \nu_l(x) \otimes \left\{ \delta_{le} \bar{e}(x) \gamma^\mu (1 - \gamma_5) e(x) + \right. \\ & \left. + \sum_f \bar{f}(x) \gamma^\mu [I_3(1 - \gamma_5) - 2Q \sin^2 \theta_W] f(x) \right\}, \end{aligned} \quad (4.14)$$

where $l = e, \mu$ identifies the neutrino fields; $f = e, p, n$ characterizes the fermionic fields (electrons, protons and neutrons respectively); I_3 and Q are the isospin and charge operators; θ_W is the weak angle and G_F the Fermi constant. In this case, only flavor-preserving transitions are allowed.

In the representation for which γ_5 is diagonal, the left component of neutrino field can be written as

$$\nu_l(x)_L = \frac{1}{2}(1 - \gamma_5)\nu_l(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_l(x) \\ -\phi_l(x) \end{pmatrix}; \quad (4.15)$$

correspondingly the neutrino currents become

$$V_S^{lm\mu}(x) = 2\phi_l^\dagger(x) \sigma_\mu \phi_m(x), \quad (4.16)$$

where $\{\sigma_\mu, \mu = 0, \dots, 3\}$ is the set of Pauli matrices plus the 2×2 identity. The Weyl spinors $\phi_l(x)$ can be expanded only in the sector of interest over components of negative helicity and positive energy [116] so that at $t = 0$

$$\phi_l(\vec{x}, 0) = \frac{1}{\sqrt{V}} \sum_{\vec{k}} e^{i\vec{k} \cdot \vec{x}} \beta(\vec{k}) \phi_l(\vec{k}), \quad (4.17)$$

where V is the volume of the layer of matter crossed by the neutrino, and $\beta(\vec{k})$ is the negative elicity eigenstate,

$$\frac{\sigma \cdot \vec{k}}{k} \beta(\vec{k}) = -\frac{1}{2} \beta(\vec{k}). \quad (4.18)$$

$\phi_l(\vec{k})$ are the annihilation operators of neutrino states with flavor l and momentum $\vec{k}$ and, together with the creation operators $\phi_l^\dagger(\vec{k})$, they satisfy anti-commutation relations:

$$\phi_l^\dagger(\vec{k})|0\rangle = |l, \vec{k}\rangle; \quad \{\phi_l(\vec{k}), \phi_l^\dagger(\vec{k}')\} = \delta_{\vec{k}\vec{k}'}, \quad \{\phi_l(\vec{k}), \phi_l(\vec{k}')\} = \{\phi_l^\dagger(\vec{k}), \phi_l^\dagger(\vec{k}')\} = 0. \quad (4.19)$$

Using (4.17) the currents (4.12) read

$$V_S^\alpha(\vec{x}, 0) = \frac{1}{V} \sum_{\vec{k}, \vec{k}'} e^{i(\vec{k}' - \vec{k}) \cdot \vec{x}} S_\mu(\vec{k}, \vec{k}') \phi_l^\dagger(\vec{k}) \phi_m(\vec{k}'), \quad (4.20)$$

where $\alpha = (l, m, \mu)$ and $S_\mu(\vec{k}, \vec{k}') \equiv \beta^\dagger(\vec{k}) \sigma_\mu \beta(\vec{k}')$. The behavior of this quantity under complex conjugation is

$$[S_\mu(\vec{k}, \vec{k}')]^* = S_\mu(\vec{k}', \vec{k}). \quad (4.21)$$

These quantities shall be employed in the following sections.

The generic neutrino state ρ_S in the finite volume V can be written in terms of flavor and momentum eigenstates:

$$\rho_S(t) = \sum_{i,j} \sum_{\vec{r}, \vec{s}} \rho_{ij}(\vec{r}, \vec{s}; t) |i\vec{r}\rangle \langle j\vec{s}| = \sum_{i,j} \sum_{\vec{r}, \vec{s}} \rho_{ij}(\vec{r}, \vec{s}; t) \phi_i^\dagger(\vec{r}) |0\rangle \langle 0| \phi_j(\vec{s}), \quad (4.22)$$

where the creation and annihilation operators satisfy (4.19).

4.3 The Markovian approximation

Following the standard steps introduced in Chapter 1, in order to deal with the environment and to end up with a Markov time-evolution for the neutrino state ρ_S , we have to perform a suitable approximation.

In Section 1.4 we have described two different procedures to obtain a Markovian approximation for the evolution of a system in a thermal bath. Introduced the typical times τ_S and τ_B , characterizing system and environment respectively, a Markovian approximation amounts to $\tau_S/\tau_B \rightarrow +\infty$, that is either $\tau_S \rightarrow +\infty$ or $\tau_B \rightarrow 0$. In the physical context we are actually considering, the typical time of the system is of the order of the period of oscillations, $\tau_S \sim E/\Delta m^2$. On the other hand, the characteristic time of the environment, τ_B , is related to the decay time of its correlations. To get a Markovian approximation,

we consider the case $\tau_B \rightarrow 0$; that is, we consider environments whose correlation functions decay in sufficiently short times. As we have seen in Section 2.4, this procedure corresponds to the singular coupling limit.

Remark 4.6 We are considering a homogeneous thermal bath in a finite volume V , thus the Markovian approximation, that requires an infinite volume, is to be performed in the hypothesis of a volume V big enough so that the Poincaré recurrences can be neglected [1].

The singular coupling amounts to considering two-point correlation functions behaving like Dirac-deltas in time:

$$h_{\alpha\alpha'}(x, x') = \text{Tr}_B[\rho_B V_B^\alpha(\vec{x}, t) V_B^{\alpha'}(\vec{x}', t')] \rightarrow c_{\alpha\alpha'}(\vec{x}, \vec{x}') \delta(t - t'). \quad (4.23)$$

According to the discussion of the previous chapters, the resulting Markovian evolution equation is:

$$\begin{aligned} \dot{\rho}_S(t) = -i \left[H_S^{eff}, \rho_S(t) \right] + \frac{1}{2} \sum_{\alpha, \alpha'} \iint_V c_{\alpha\alpha'}(\vec{x}, \vec{x}') & \left([V_S^\alpha(\vec{x}, 0) \rho_S(t), V_S^{\alpha'}(\vec{x}', 0)] + \right. \\ & \left. + [V_S^\alpha(\vec{x}, 0), \rho_S(t) V_S^{\alpha'}(\vec{x}', 0)] \right) d\vec{x} d\vec{x}' \end{aligned} \quad (4.24)$$

and the statistical operator ρ_S lives in the neutrino Hilbert space $\mathfrak{H}_S$, containing flavor and space degrees of freedom. We shall later try to reduce further the description onto the flavor degrees of freedom. In (4.24), H_S^{eff} is the effective neutrino Hamiltonian obtained from H_S by taking into account the mean field term produced by the interaction with the environment: $H_S^{eff} = H_S + \text{Tr}_B(\rho_B H_I)$, with H_I defined in (4.10) and ρ_B the thermal equilibrium state of the bath; see the discussion in Section 1.3. It is the effective Hamiltonian usually introduced in the description of the interactions of neutrinos in matter leading to the MSW effect [91, 92, 93, 108]. In the two-dimensional flavor space, the representation of H_S^{eff} over the physical states is

$$H_S^{eff} = \frac{\Delta m^2}{4E} \begin{pmatrix} -\cos 2\theta & \sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{pmatrix} + \begin{pmatrix} V_{ee} & V_{e\mu} \\ V_{\mu e} & V_{\mu\mu} \end{pmatrix} \quad (4.25)$$

where

$$\begin{aligned} V_{ee} &\equiv \langle \nu_e | \text{Tr}_B(\rho_B H_I) | \nu_e \rangle, & V_{e\mu} &\equiv \langle \nu_e | \text{Tr}_B(\rho_B H_I) | \nu_\mu \rangle, \\ V_{\mu\mu} &\equiv \langle \nu_\mu | \text{Tr}_B(\rho_B H_I) | \nu_\mu \rangle, & V_{\mu e} &\equiv \langle \nu_\mu | \text{Tr}_B(\rho_B H_I) | \nu_e \rangle, \end{aligned} \quad (4.26)$$

are contributions of first order in H_I and $V_{ee}, V_{\mu\mu} \in \mathbf{R}$, $V_{\mu e} = V_{e\mu}^*$.

Remark 4.7 The off-diagonal entries vanish for flavor-preserving interactions. Indeed, in this case in (4.11) we have

$$V_B^{lm\mu}(x) = \delta_{lm} V_B^{ll\mu}(x), \quad (4.27)$$

where $l, m = e, \mu$. For instance, for neutrinos in solar matter, we find:

$$V_{ee} = \sqrt{2}G_F \left(n_e - \frac{1}{2}n_n \right), \quad V_{\mu\mu} = -\frac{1}{\sqrt{2}}G_F n_n, \quad V_{e\mu} = V_{\mu e} = 0, \quad (4.28)$$

where n_e and n_n are the density of electrons, neutrons respectively, and G_F is the Fermi constant.

The effective Hamiltonian is diagonalized by an orthogonal transformation U that connects the eigenstates of H_S^{eff} to the physical states:

$$\begin{cases} |\nu_e\rangle = U|\nu_{m1}\rangle, \\ |\nu_\mu\rangle = U|\nu_{m2}\rangle; \end{cases} \quad U = \begin{pmatrix} \cos \theta_m & \sin \theta_m \\ -\sin \theta_m & \cos \theta_m \end{pmatrix}, \quad (4.29)$$

where $\{|\nu_{m1}\rangle, |\nu_{m2}\rangle\}$ are the eigenstates of H_S^{eff} and θ_m is the mixing angle in matter. The Hamiltonian eigenvalues in the adopted relativistic approximation are $\epsilon_{m1} = -\Delta m_m^2/4E$ and $\epsilon_{m2} = \Delta m_m^2/4E$ where $\Delta m_m^2 = m_{m2}^2 - m_{m1}^2$ and m_{m1}, m_{m2} are the neutrino masses in the medium. They are related to the equivalent quantities in vacuum; if $V_{e\mu} = V_{\mu e} = 0$, as in the case of solar matter, we have [92, 93]

$$\tan 2\theta_m = \frac{\Delta m^2 \sin 2\theta}{\Delta m^2 \cos 2\theta - A}, \quad (4.30a)$$

$$\Delta m_m^2 = \sqrt{A^2 + (\Delta m^2)^2 - 2A\Delta m^2 \cos 2\theta}, \quad (4.30b)$$

where

$$A \equiv 2E(V_{ee} - V_{\mu\mu}). \quad (4.31)$$

Remark 4.8 The resonance in equation (4.30a) produces an enhancement of the oscillating behavior that, combined with a suitable variation of the density of the crossed medium, generates the so-called MSW effect, actually the best candidate to the explanation of the problem of solar neutrinos [92, 93].

To first order in the interaction (that is neglecting the dissipative contributions of equation (4.24)), the transition probability in the uniform medium considered reads

$$P_{\nu_e \rightarrow \nu_\mu}(t) = \sin^2 2\theta_m \sin^2 \frac{\Delta m_m^2}{4E} t. \quad (4.32)$$

The dissipative effects are of second order in H_I ; they result in Hamiltonian, damping and noise terms as results from rearranging equation (4.24):

$$\begin{aligned} \dot{\rho}_S(t) = & -i \left[H_S^{eff}, \rho_S(t) \right] - \frac{1}{2} \sum_{\alpha, \alpha'} \iint_V d\vec{x} d\vec{x}' b_{\alpha\alpha'}(\vec{x}, \vec{x}') [V_S^\alpha(\vec{x}, 0) V_S^{\alpha'}(\vec{x}', 0), \rho_S(t)] + \\ & - \frac{1}{2} \sum_{\alpha, \alpha'} \iint_V d\vec{x} d\vec{x}' a_{\alpha\alpha'}(\vec{x}, \vec{x}') \{V_S^\alpha(\vec{x}, 0) V_S^{\alpha'}(\vec{x}', 0), \rho_S(t)\} + \\ & + \sum_{\alpha, \alpha'} \iint_V d\vec{x} d\vec{x}' a_{\alpha\alpha'}(\vec{x}, \vec{x}') V_S^{\alpha'}(\vec{x}', 0) \rho_S(t) V_S^\alpha(\vec{x}, 0), \end{aligned} \quad (4.33)$$

where the new coefficients are defined as

$$a_{\alpha\alpha'}(\vec{x}, \vec{x}') \equiv \int_{-\infty}^{\infty} h_{\alpha\alpha'}(\vec{x}, \vec{x}'; u) du; \quad (4.34a)$$

$$b_{\alpha\alpha'}(\vec{x}, \vec{x}') \equiv \int_0^{\infty} h_{\alpha\alpha'}(\vec{x}, \vec{x}'; u) du - \int_{-\infty}^0 h_{\alpha\alpha'}(\vec{x}, \vec{x}'; u) du. \quad (4.34b)$$

According to Section 1.5, in equation (4.33) we recognize the typical form of the generator of a Markov semigroup.

Remark 4.9 Accordingly to Remark 1.17, in the first row of (4.33) we note that the vacuum Hamiltonian H_S is redefined by contributions of first order in the interaction, already accounted for in H_S^{eff} , and of second order, determined by the generalized matrix $[b_{\alpha\alpha'}(\vec{x}, \vec{x}')]$.

Remark 4.10 Since the time-evolution (4.33) is obtained from a singular coupling, it is completely positive and the generalized matrix $[a_{\alpha\alpha'}(\vec{x}, \vec{x}')] is positive definite.$

4.4 The generator of the dissipative evolution

We want to write the form of the generator of the dissipative evolution, eq. (4.33) in the flavor space $\mathfrak{H}_2$ only; we thus have to get rid of the momentum degrees of freedom. We start by considering the expression (4.22) for the statistical operator ρ_S . By rearranging

the summation indices, (4.33) results in the following equation for the entries of $\rho_S(t)$:

$$\begin{aligned} \dot{\rho}_{ij}(\vec{r}, \vec{s}; t) = & -i \sum_k \left[(H_S^{eff})_{ik} \rho_{kj}(\vec{r}, \vec{s}; t) - \rho_{ik}(\vec{r}, \vec{s}; t) (H_S^{eff})_{kj} \right] + \\ & - \frac{g^2}{V^2} \sum_{l,m} \sum_{\vec{k}, \vec{q}} \left\{ \left[A_{imlm}(\vec{r}, \vec{q}, \vec{k}, \vec{q}) + B_{imlm}(\vec{r}, \vec{q}, \vec{k}, \vec{q}) \right] \rho_{lj}(\vec{k}, \vec{s}; t) + \right. \\ & + \left[A_{lmjm}(\vec{q}, \vec{k}, \vec{s}, \vec{k}) - B_{lmjm}(\vec{q}, \vec{k}, \vec{s}, \vec{k}) \right] \rho_{il}(\vec{r}, \vec{q}; t) + \\ & \left. - 2A_{ljli}(\vec{k}, \vec{s}, \vec{q}, \vec{r}) \rho_{lm}(\vec{q}, \vec{k}; t) \right\}, \end{aligned} \quad (4.35)$$

where

$$A_{lm'l'm'}(\vec{k}, \vec{k}', \vec{q}, \vec{q}') \equiv \sum_{\mu, \mu'} \iint_V d\vec{x} d\vec{x}' a_{\alpha\alpha'}(\vec{x}, \vec{x}') e^{i(\vec{k}' - \vec{k}) \cdot \vec{x}} e^{-i(\vec{q}' - \vec{q}) \cdot \vec{x}'} S_\mu(\vec{k}, \vec{k}') S_{\mu'}(\vec{q}', \vec{q}), \quad (4.36a)$$

$$B_{lm'l'm'}(\vec{k}, \vec{k}', \vec{q}, \vec{q}') \equiv \sum_{\mu, \mu'} \iint_V d\vec{x} d\vec{x}' b_{\alpha\alpha'}(\vec{x}, \vec{x}') e^{i(\vec{k}' - \vec{k}) \cdot \vec{x}} e^{-i(\vec{q}' - \vec{q}) \cdot \vec{x}'} S_\mu(\vec{k}, \vec{k}') S_{\mu'}(\vec{q}', \vec{q}), \quad (4.36b)$$

and $\alpha = (l, m, \mu)$, $\alpha' = (l', m', \mu')$.

Equation (4.35) is rather complex; however, under suitable conditions, it is possible to simplify it, reducing the effective Hilbert space to a two-dimensional one. The Hilbert space describing our system is $\mathfrak{H}_S = \mathfrak{H}_2 \otimes \mathfrak{H}_p$, where $\mathfrak{H}_2$ is the two-dimensional flavor Hilbert space whereas $\mathfrak{H}_p$ is the momentum Hilbert space. The reduction to $\mathfrak{H}_2$ is performed by adapting the projector technique developed in Section 1.3. We shall see that such a reduction will produce a nicer equation than (4.35), under conditions that are not too restrictive.

Remark 4.11 Usually the formalism of projectors is applied to eliminate the environment degrees of freedom, leaving only those of the system of interest. In the present case, however, the trace over the environment has already been performed, and an adapted version of the projection technique allows us to discard the momentum degrees of freedom.

Let us start with the time-evolution (4.35); for later convenience we shall adopt the compact notation

$$\dot{\rho}_S(t) = L[\rho_S(t)] = (L_H + L_D)[\rho_S(t)], \quad (4.37)$$

in which we have separated the Hamiltonian and the dissipative part of the generator (4.33). In particular, $L_H[\rho_S(t)]$ represents the first row of (4.33) while $L_D[\rho_S(t)]$ the latter two.

The state ρ_S acts on $\mathfrak{H}_2 \otimes \mathfrak{H}_p$ and we want to extract the time-evolution of the reduced statistical operator acting on $\mathfrak{H}_2$ only. We will denote this latter by ρ ; it is obtained by tracing ρ_S over the momentum degrees of freedom: $\rho = \text{Tr}_p(\rho_S)$.

In order to apply the projector technique, we need a reference state for the momentum degrees of freedom: we choose $\rho_p = |\vec{p}\rangle\langle\vec{p}|$, where $\vec{p}$ is the momentum of the incoming neutrino; accordingly we define the projectors $P[\cdot] = \text{Tr}_p(\cdot) \otimes \rho_p$ and $Q = \mathbf{I} - P$. Adapting relations (1.29) and (1.30) to this case, from (4.37) we extract the following equation of motion:

$$\dot{\rho}(t) = P(L_H + L_D)\rho(t) + \int_0^t dt' P(L_H + L_D)e^{QL(t-t')}QL_D\rho(t'), \quad (4.38)$$

where we have used the property $QL_HP = 0$ and supposed the initial state to be factorized: $\rho_S(0) = \rho(0) \otimes \rho_p$.

This is not a manageable expression yet because of the convolution term in the right-hand side. In order to get a time-evolution having the structure of semigroup, for flavor degrees of freedom only, we have to make further assumptions.

After explicit computation, the dependence of $A(\vec{k}, \vec{k}', \vec{q}, \vec{q}')$ and $B(\vec{k}, \vec{k}', \vec{q}, \vec{q}')$ on the momenta, in the operators PL_D and QL_D , is rather different: in PL_D these coefficients present in any case two pairs of equal momentum indices, while this never happens for QL_D . To understand the physical meaning of this, observe that these coefficients are Fourier transforms, for example

$$A(\vec{k}, \vec{k}', \vec{q}, \vec{q}') \approx \iint_V d\vec{x}d\vec{x}' a(\vec{x}, \vec{x}') e^{i(\vec{k}' - \vec{k}) \cdot \vec{x}} e^{-i(\vec{q}' - \vec{q}) \cdot \vec{x}'}, \quad (4.39)$$

where $a(\vec{x}, \vec{x}')$ is a spatial correlation function of the bath; in general, it can be written as the sum of two functions: $f(\vec{x}, \vec{x}') = f(|\vec{x} - \vec{x}'|)$, traslationally invariant, and a contribution $g(\vec{x}, \vec{x}')$:

$$a(\vec{x}, \vec{x}') = f(|\vec{x} - \vec{x}'|) + g(\vec{x}, \vec{x}'). \quad (4.40)$$

Therefore (4.39) becomes

$$A(\vec{k}, \vec{k}', \vec{q}, \vec{q}') \approx \delta_{(\vec{k} - \vec{k}'), (\vec{q} - \vec{q}')} \int_V d\vec{z} f(|z|) e^{i(\vec{q}' - \vec{q}) \cdot \vec{z}} + \iint_V d\vec{x}d\vec{x}' g(\vec{x}, \vec{x}') e^{i(\vec{k}' - \vec{k}) \cdot \vec{x}} e^{-i(\vec{q}' - \vec{q}) \cdot \vec{x}'}. \quad (4.41)$$

The contributions to PL_D come from both the terms in the right-hand side of this equation whereas the contributions to QL_D are produced only from the Fourier transform of $g(\vec{x}, \vec{x}')$. Translational invariance means momentum conservation, therefore the term PL_D does not change momentum, QL_D instead does. Thus, if the correlation function $a(\vec{x}, \vec{x}')$ is traslationally invariant (i.e. $g(\vec{x}, \vec{x}') = 0$) automatically $QL_D = 0$ and the integral term in eq. (4.38) vanishes. In general it could be neglected if, in $a(\vec{x}, \vec{x}')$, the

translationally invariant part $f(|\vec{x} - \vec{x}'|)$ is dominant with respect to the non invariant one, $g(\vec{x}, \vec{x}')$.

In the following, we work within the hypothesis of almost translational invariance within V , so that (4.38) becomes:

$$\dot{\rho}(t) = P(L_H + L_D)\rho(t), \quad (4.42)$$

and the evolution equation for the entries of the 2×2 statistical operator ρ reads

$$\begin{aligned} \dot{\rho}_{ij}(t) = & -i \sum_k \left[\left(H_S^{eff} \right)_{ik} \rho_{kj}(t) - \rho_{ik}(t) \left(H_S^{eff} \right)_{kj} \right] + \\ & - \frac{g^2}{V^2} \sum_l \left\{ [A_{il}(\vec{p}) + B_{il}(\vec{p})] \rho_{lj}(t) + [A_{lj}(\vec{p}) - B_{lj}(\vec{p})] \rho_{il}(t) + 2 \sum_m A_{mjli}(\vec{p}) \rho_{lm}(t) \right\}, \end{aligned} \quad (4.43)$$

where

$$A_{imjl}(\vec{p}) = \sum_{\vec{k}} A_{imjl}(\vec{p}, \vec{k}, \vec{p}, \vec{k}), \quad A_{ij}(\vec{p}) = \sum_m A_{imjm}(\vec{p}); \quad (4.44a)$$

$$B_{ij}(\vec{p}) = \sum_{\vec{k}} \sum_m B_{imjm}(\vec{p}, \vec{k}, \vec{p}, \vec{k}). \quad (4.44b)$$

In vector representation, using the physical basis for $\mathfrak{H}_2$, this equation assumes the form

$$\frac{d}{dt} |\rho(t)\rangle = \mathcal{L} |\rho(t)\rangle = (\mathcal{H} + \mathcal{D}) |\rho(t)\rangle, \quad (4.45)$$

where

$$\mathcal{H} = -2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & h_3 & -h_2 \\ 0 & -h_3 & 0 & h_1 \\ 0 & h_2 & -h_1 & 0 \end{pmatrix}, \quad \mathcal{D} = -2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ u & a & b & c \\ v & b & \alpha & \beta \\ w & c & \beta & \gamma \end{pmatrix}. \quad (4.46)$$

The matrix $\mathcal{L} = \mathcal{H} + \mathcal{D}$ is the generator of the reduced time-evolution we are looking for. The parameters characterizing the Hamiltonian part are

$$\begin{aligned} h_1 &= \frac{\Delta m^2}{4E} \sin 2\theta + \text{Re } V_{e\mu} + \omega_1, \\ h_2 &= -\text{Im } V_{e\mu} + \omega_2, \\ h_3 &= -\frac{\Delta m^2}{4E} \cos 2\theta + \frac{1}{2}(V_{ee} - V_{\mu\mu}) + \omega_3, \end{aligned} \quad (4.47)$$

where ω_1 , ω_2 and ω_3 are the Hamiltonian contributions of second order in the interaction (see Remark 4.9), whose expressions are reported in Appendix D.

Following the discussion of Sections 1.8 and 2.3, the coefficients u , v and w in the dissipative part $\mathcal{D}$ are related to entropy behavior and to the existence of asymptotic states different from the maximally mixed state $|\rho_2\rangle = 1/2(1, 0, 0, 0)$. The remaining parameters, instead, produce decoherence and dissipation. The expressions of all of these parameters are listed in Appendix D.

These coefficients satisfy the inequalities (2.21) that guarantee the complete positivity of the evolution because they are obtained via a singular coupling limit [39].

In the neutrino's case the impact of complete positivity cannot be fully appreciated since this property is needed for the physical consistency of the time-evolution of bipartite entangled systems, and there are not phenomenological contexts of this kind in neutrino physics.

Remark 4.12 If the interactions are flavor-preserving most of these coefficients vanish since we have

$$A_{imjl} = \delta_{im}\delta_{jl}A_{iijj}, \quad A_{ij} = \delta_{ij}A_{ii}, \quad B_{ij} = \delta_{ij}B_{ii}. \quad (4.48)$$

Considering the expressions reported in Appendix D, we find $\omega_1 = \omega_2 = 0$, $u = v = w = 0$, $a = \alpha$ and $b = c = \gamma = 0$; the dissipative contribution is fully parameterized by only one coefficient, a , in agreement with [98].

4.5 An example of flavor changing dissipative evolution

As already mentioned (see Section 2.4), the exact integration of equation (4.45) is always possible, but the expression of $\mathcal{G}_t$ is not transparent enough to be amenable to physical interpretations. It is more convenient to expand $\mathcal{G}_t$ to first order in the dissipative parameters appearing in $\mathcal{D}$; this computation has already been motivated and developed in Section 2.4 and applied to neutrinos in [117, 118].

In this section, however, we wish to give the exact transition probability in a simplified case [117]: $h_2 = 0$, $u = v = w = 0$, $a = \alpha = \gamma$ and $c = 0$. As explained in Remark 4.12, this latter case amounts to flavor-changing interactions since b , β and γ are not vanishing and it is compatible with the inequalities expressing the complete positivity of the evolution provided that $a^2 \geq b^2 + \beta^2$.

Remark 4.13 In our vector representation the transition probability can be expressed as an inner product:

$$\mathcal{P}_{\nu_e \rightarrow \nu_\mu}(t) = 2\langle \rho_{\nu_\mu} | \rho_{\nu_e}(t) \rangle = 2\langle \rho_{\nu_\mu} | \mathcal{G}_t | \rho_{\nu_e} \rangle, \quad (4.49)$$

as can be seen by putting in (4.9) the expansions of both ρ_{ν_e} and ρ_{ν_μ} over the basis of Pauli matrices plus identity.

After exponentiation of the generator $\mathcal{L}$, the time-evolutor $\mathcal{G}_t = e^{\mathcal{L}t}$ amounts to

$$\mathcal{G}_t = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & & & \\ 0 & \tilde{\mathcal{G}}_t & & \\ 0 & & & \end{pmatrix} \quad (4.50)$$

where the 3×3 sub-matrix $\tilde{\mathcal{G}}_t$ is given by

$$\tilde{\mathcal{G}}_t = e^{-2at} \left[\mathbf{I}_3 + \frac{\sin 2\Theta_m t}{2\Theta_m} \mathcal{M} + \frac{\sin^2 \Theta_m t}{4\Theta_m^2} \mathcal{M}^2 \right], \quad (4.51)$$

with

$$\mathcal{M} = -2 \begin{pmatrix} 0 & b + h_3 & 0 \\ b - h_3 & 0 & \beta + h_1 \\ 0 & \beta - h_1 & 0 \end{pmatrix} \quad (4.52)$$

and $\mathbf{I}_3$ the 3×3 identity matrix. We have introduced $\Theta_m = \sqrt{h_1^2 + h_3^2 - b^2 - \beta^2}$, that can be either real or complex, depending on the relative size of the coefficients appearing in it. Since in the present context the dissipative parameters are small, we consider a real Θ_m ; the general case is presented in [117].

Remark 4.14 Note that, that, in absence of dissipation (i.e. $b = \beta = 0$) Θ_m reduces to $\Delta m_m^2/4E$, and then, in absence of matter, to $\Delta m^2/4E$.

In vectorial notation, the pure ν_e state (the initial state) and the pure ν_μ state are

$$\rho_{\nu_e} = |\nu_e\rangle\langle\nu_e|, \quad |\rho_{\nu_e}\rangle = \frac{1}{2}(1, 0, 0, 1), \quad (4.53a)$$

$$\rho_{\nu_\mu} = |\nu_\mu\rangle\langle\nu_\mu|, \quad |\rho_{\nu_\mu}\rangle = \frac{1}{2}(1, 0, 0, -1), \quad (4.53b)$$

whence

$$\mathcal{P}_{\nu_e \rightarrow \nu_\mu}(t) = \frac{1}{2} \left[1 - e^{-2at} - 2 \frac{\beta^2 - h_1^2}{\Theta_m^2} e^{-2at} \sin^2 \Theta_m t \right]. \quad (4.54)$$

Consistently, if there is not dissipation (4.54) gives the transition probability in matter to first order in the interaction, that is equation (4.32). In absence of matter, we get the vacuum transition probability (4.7).

The damping exponentials, typical of dissipative phenomena, flatten the oscillations in the transition probability so that it approaches the value $1/2$ for very long times, and this happens regardless of the value of the vacuum mixing angle θ . We remind that in vacuum the transition probability has the upper limit $\sin^2 2\theta$ and then tends to vanish for θ approaching 0 whereas the asymptotic mixing between the two flavors, in the dissipative case, is maximal, regardless of the value of θ .

4.6 Non homogeneous matter

Up to now we have considered a simplified environment with a constant mean density. However the physical appeal of neutrinos interacting with matter arises when the density of the considered medium is varying along the neutrino path. This is indeed the real situation: in the solar case the neutrinos are produced near the core of the Sun, the region with higher density, then they travel through regions with gradually decreasing densities. Actually the more plausible explanation of the missing neutrino solar flux depends strongly on the varying profile of the Sun density [92, 93].

Since any slab of matter with a not-constant density profile can be well approximated by a sequence of homogeneous layers with suitable thickness, the time evolver can be obtained by composing elementary evolutors computed using the formalism described before. That is, if the inhomogeneous medium consists of n homogeneous layers, one has

$$\mathcal{G}_t = \mathcal{G}_{t_n}^n \circ \dots \circ \mathcal{G}_{t_2}^2 \circ \mathcal{G}_{t_1}^1, \quad t = t_1 + t_2 + \dots + t_n \quad (4.55)$$

where $t_i, i = 1, \dots, n$ are the times needed to traverse the successive layers. The composition of maps in (4.55) is completely positive by construction.

Then in our formalism all we need to describe the neutrino evolution in the real matter is to divide the traversed medium into suitable slabs, possibly of different thickness, depending on the rapidity of the mean density change.

Traditionally, in the study of neutrino transitions in matter two approaches are possible to account for the change of the medium density: it can vary slowly, and the transitions are called adiabatic, or rapidly. In the adiabatic approximation, neutrino state evolves in time like eigenstates of the effective Hamiltonian that, in the neutrino frame, is varying in time; otherwise neutrinos do not necessarily remain in eigenstates of $H_S^{eff}(t)$ but can pass from an eigenstate to another. This standard formalism has been adapted to the case of a dissipative evolution for statistical operators in [117].

4.7 The solar matter case

In this section we apply our formalism to describe the dissipative time-evolution of a neutrino crossing a layer of solar matter with constant mean density and to evaluate the impact of this modified evolution on the flavors transition probability. We adopt the Hamiltonian density (4.14).

The autocorrelation functions reflect the decoupling of flavors:

$$\begin{aligned} a_{lm\mu, l'm'\mu'}(\vec{x}, \vec{x}') &= \delta_{lm} \delta_{l'm'} \delta_{ll'} a_{lm\mu, l'm'\mu'}(\vec{x}, \vec{x}'), \\ b_{lm\mu, l'm'\mu'}(\vec{x}, \vec{x}') &= \delta_{lm} \delta_{l'm'} \delta_{ll'} b_{lm\mu, l'm'\mu'}(\vec{x}, \vec{x}'); \end{aligned} \quad (4.56)$$

$\delta_{ll'}$ means statistical independence of the bath operators of different flavors. As a consequence of (4.56) most of the coefficients in Appendix D vanish and the evolution equation

in the flavor basis, eq. (4.45), takes the simple form

$$\frac{d}{dt}|\rho(t)\rangle = -2 \left[\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & h_3 & 0 \\ 0 & -h_3 & 0 & h_1 \\ 0 & 0 & -h_1 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & a & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \right] |\rho(t)\rangle, \quad (4.57)$$

where, following (4.47), (4.28) and Appendix D,

$$\begin{aligned} h_1 &= i \frac{\Delta m^2}{4E} \sin 2\theta, \\ h_3 &= i \left(\frac{G_F}{\sqrt{2}} n_e - \frac{\Delta m^2}{4E} \cos 2\theta \right) + \omega_3, \\ \omega_3 &= \frac{1}{2} \text{Im} [B_{\mu\mu}(\vec{p}) - B_{ee}(\vec{p})], \\ a &= \frac{1}{2} [A_{ee}(\vec{p}) + A_{\mu\mu}(\vec{p})]. \end{aligned} \quad (4.58)$$

In order to evaluate the time-evolution of this state, we use the formalism developed in Section 2.4, with $r_1 = -\cot \theta_m$ and $r_2 = -\tan \theta_m$. It is convenient to perform the computation in the basis $\{|\nu_{m1}\rangle, |\nu_{m2}\rangle\}$ that diagonalizes H_S^{eff} (introduced in (4.29)); indeed $P_{\nu_e \rightarrow \nu_\mu}(t)$ is basis independent by definition.

The resulting expression of the transition probability in the dissipative regime is thus

$$\begin{aligned} P_{\nu_e \rightarrow \nu_\mu}(t) &= \frac{1}{2} \left\{ 1 - e^{-2a \sin^2 2\theta_m t} \cos^2 2\theta_m + \right. \\ &\quad \left. - \sin^2 2\theta_m \left[e^{-a(1+\cos^2 2\theta_m)t} \cos \frac{\Delta m^2}{2E} t + a \frac{2E}{\Delta m^2} (1 + 3 \cos^2 2\theta_m) \sin \frac{\Delta m^2}{2E} t \right] \right\}, \end{aligned} \quad (4.59)$$

where the parameter ω_3 has been absorbed in a redefinition of the parameters appearing in the Hamiltonian part of the evolution. For $a = 0$ we obtain the transition probability at first order in the interaction for an homogeneous medium, eq. (4.32). It reduces to the vacuum oscillation probability in absence of matter.

In order to obtain the transition probability in the real, not uniform, solar matter, the considerations of Section 4.6 should be applied; however, using a magnitude estimate based on dimensional analysis, the impact of dissipation for the phenomena that can be described in our framework is negligible, as we shall shortly explain.

4.8 Discussion and conclusions

We have treated the topic of neutrino oscillations in matter from the rigorous point of view of open quantum systems, using rigorous Markovian limits that produce a fully consistent

dynamics for the described system and avoids problems of physical consistency.

The interaction between neutrinos and environment has been considered to be as general as possible, including some scenarios forbidden in the Standard Model, like flavor changing currents. The formalism of the singular coupling has been employed, producing a request on the bath correlation time: it must be much shorter than the characteristic time of the system.

The conditions to obtain a convolutionless dynamics in the flavor space have been discussed, showing the tight connection between the semigroup structure of the reduced dynamics and the isotropy of the bath correlation functions (i.e. their invariance under translations). No specific models of fluctuations have been introduced; we have rather assumed that the contributions breaking this isotropy be negligible. If they are not so, flavor and momentum degrees of freedom need to be treated together and an effective two-dimensional prescription is unavailable.

In this framework, several dissipative phenomena can be described. We have evaluated the transition probability between two neutrino flavors in two different contexts: a flavor-changing and a flavor-preserving interaction and explicit results for homogeneous matter have been obtained. To account for matter inhomogeneities, in general, numerical computations should be performed.

The typical terms introduced by dissipation are damping exponentials whose lifetimes are the reciprocals of the dissipative coefficients; this means that the effect of the irreversible dynamics should be relevant on a time scale of order $t_{diss} \sim 1/(gn)^2\tau_B$, the inverse of the typical magnitude of dissipative parameters in non-relativistic matter, where n is the density of scatterers and g the coupling constant in the interaction term.

For example, the thermal fluctuations in the Sun satisfy the limits imposed by our formalism; a satisfactory evaluation of the correlation time is given by the inverse of the thermal energy at temperature T : $\tau_B = \hbar/kT$, where $\hbar$ and k are the Planck and Boltzmann constants respectively. The explicit evaluation for the Sun core ($T \sim 10^7$ K) gives $\tau_B \sim 10^{-21}$ s, much shorter than $\tau_S \sim E/\Delta m^2 \sim 10^{-4}$ s (obtained considering the values $E \sim 1$ MeV and $\Delta m^2 \sim 10^{-5}$ eV²). However, neutrinos cross the Sun in a time so short that the thermal contributions are completely negligible: $t_{diss} \sim 10^{14}$ s (using the electron density $n_e \sim 10^{26}$ cm⁻³ and $G_F \sim 10^{-5}$ GeV⁻²), versus a crossing time of about 3 seconds.

Does it exist a physical instance in which the dissipative effects give a strong effect despite the weak interaction with the surroundings? It is difficult to imagine a real situation of this kind. However the gravitational thermal bath described in Section 3.2 could give such an environment: it should be everywhere present, and plausibly homogeneous and isotropic. In this case the bath energy is parametrized by the Plank mass, $M_P \sim 10^{19}$ GeV, and for a neutrino energy of order $E_S \sim 1$ MeV, we can roughly evaluate $t_{diss} \sim 10^7$ s, that is of order of the time needed for a neutrino to cross intergalactic distances. The asymptotical behavior for the transition probability, with and without dissipation, should be very different. In the first case it would approach the 1/2 value; in the second one it would oscillate with a maximal amplitude of $\sin^2 2\theta$ (observe that in this scenario it is impossible

to distinguish between θ and θ_m). The averaged transition probability would be certainly increased and its time dependence destroyed. In general, decoherence in neutrino physics has been studied in [117, 119, 120, 121].

The last remark is about the discrimination of the Dirac or Majorana nature of neutrino fields [122, 123]: although dissipative dynamics in general allow to distinguish between these two possibilities [117], this is not the case in this chapter, since the microscopic interactions just presented are diagonal in the flavor neutrino fields.

Chapter 5

Neutron interferometry

In this chapter we apply the formalism of open quantum systems to study neutron interferometry. There are two main motivations for this kind of approach. First of all, physical setups that are subjected to irreversible evolutions are actually available: they give the opportunity to study several models of Markovian dynamics, and to test their physical consistency. In the second place, since experiments based on quantum interferometry strongly rely upon the property of quantum correlations between different degrees of freedom, they can be suitably employed to investigate the notion of complete positivity (CP) of the reduced time-evolution Λ_t . As we have seen in Chapter 1, this property, tightly connected to quantum correlations expressed by entanglement, has a somewhat abstract and technical definition; it could seem at first sight a mathematical artifact rather than a physically motivated request. However in this context the relevance of CP becomes apparent since we are dealing here with reduced dynamics of the form $\mathbf{I} \otimes \Lambda_t$, precisely the one used in its definition (see Section 1.2). Indeed, as explicitly shown in the models discussed below, lack of CP necessarily leads to the appearance of unacceptable negative eigenvalues in the time evolving physical state describing entangled degrees of freedom.

After a brief introduction to the topic of neutron interferometry, and to the mathematical framework usually employed to describe it, we shall study non standard dissipative evolutions for neutrons inside the interferometer. These reduced dynamics are the result of the interaction with an external environment in which the interferometric device is immersed.

We shall be very general, allowing for evolutions that are not completely positive. Three different prototypes will be taken into account, all of them experimentally achievable by choosing appropriately the properties of the external environment (typically a thermal bath): they cover instances of completely positive, simply positive (but not CP) and even not-positive time-evolutions. Some of these dynamics are commonly used in some branches of physics (for example, in chemical physics [125, 126, 27]).

Neutron interferometry has been proposed to test the non-contextuality hypothesis in quantum mechanics [129], that is the possibility that all the members of any set of

commuting observables posses a value independent on the choice of the commuting observables measured at the same time (that is, independent on the context in which the measure of that observable is performed). This hypothesis can be tested by studying violations of the CHSH inequality [128] (a particular Bell-like inequality); in the final part of the chapter, using the previously developed results, we will study how this relation is modified when the interferometer is immersed in a stochastic background.

5.1 The physical apparatus

For our discussion, we shall refer to an idealized, but standard model of neutron interferometer¹ [129]. The apparatus, sketched in Figure 5.1, consists of three basic components: an initial beam splitter (bs_1), dividing an incoming, polarized, neutron beam in two components; a spin flipper (sf), acting on one of the two components emerging from the beam splitter; a final beam splitter (bs_2) that recombines the two components, equipped with a phase shifter (ps). The neutrons of the incoming beam are polarized along a fixed direction, say the z -axis; the beam splitter bs_1 is characterized by a transmission p and reflection q coefficients satisfying the condition $|p|^2 + |q|^2 = 1$ (in the ideal case where no neutrons are absorbed). The two components follow spatially separated (by distances of the order of centimeters) trajectories, denoted by u (for up) and d (for down). Then, the translational degree of freedom may be described by a two-dimensional Hilbert space $\mathfrak{H}_2^a$ generated by the orthogonal states $|\psi_u\rangle$ and $|\psi_d\rangle$, that are associated to the two macroscopically distinguished paths.

Remark 5.1 Since the neutrons are 1/2-spin particles, their spin degree of freedom is also described by a two-dimensional Hilbert space $\mathfrak{H}_2^b$, spanned by the states $|\uparrow_z\rangle$, spin-up, and $|\downarrow_z\rangle$, spin-down along the z -axis. On the whole, neutrons in the apparatus may be appropriately described by the 4-dimensional tensor Hilbert space $\mathfrak{H}_2^a \otimes \mathfrak{H}_2^b = \mathfrak{H}_4$.

Inside the interferometer, the component following the u trajectory is subjected to a spin-flip, $|\uparrow_z\rangle \rightarrow |\downarrow_z\rangle$; as a result, the state describing a neutron travelling in the inner region of the interferometer is entangled (where the entanglement relies upon a quantum correlation between internal and spatial degrees of freedom of the same system):

$$|\psi\rangle = p|\psi_u\rangle \otimes |\downarrow_z\rangle + q|\psi_d\rangle \otimes |\uparrow_z\rangle. \quad (5.1)$$

In the density matrix formalism, this state is represented by the 4×4 statistical operator

$$\rho^{ab} = |\psi\rangle\langle\psi| = |p|^2 P_1 \otimes Q_2 + |q|^2 P_2 \otimes Q_1 + pq^* P_3 \otimes Q_4 + p^* q P_4 \otimes Q_3, \quad (5.2)$$

¹similar set-ups can be constructed using polarized hotons see [130]




Figure 5.1: Typical model of neutron interferometer for non-contextuality tests. bs_1 and bs_2 are the two beam splitters, sf is the spin flipper, ps the phase shifter, c_1 and c_2 the position plus spin counters.

where

$$\begin{aligned} P_1 &\equiv |\psi_u\rangle\langle\psi_u|, & P_2 &\equiv |\psi_d\rangle\langle\psi_d|, & P_3 &\equiv |\psi_u\rangle\langle\psi_d|, & P_4 &\equiv |\psi_d\rangle\langle\psi_u|; \\ Q_1 &\equiv |\uparrow_z\rangle\langle\uparrow_z|, & Q_2 &\equiv |\downarrow_z\rangle\langle\downarrow_z|, & Q_3 &\equiv |\uparrow_z\rangle\langle\downarrow_z|, & Q_4 &\equiv |\downarrow_z\rangle\langle\uparrow_z|. \end{aligned} \quad (5.3)$$

Remark 5.2 We stress that the entanglement in the state (5.1) is different from the usual one, for instance the one described in the case of neutral **B** mesons, where quantum correlations connect the degrees of freedom of two spatially separated physical systems, a neutral **B** meson and its antiparticle, both produced in the decay of a resonance. In the present case, on the contrary, the entanglement is between different degrees of freedom of the same physical system: translational and spin degrees of freedom.

By denoting with Λ_t the dynamical map describing the time-evolution undergone inside the apparatus, at the second beam splitter, after a time t , the initial state ρ^{ab} in (5.2) is changed according to a law $\rho^{ab} \rightarrow \rho^{ab}(t) = \Lambda_t[\rho^{ab}]$.

At the second beam splitter, the u -component is further shifted by an angle φ , with respect to the d -component, by the phase shifter ps . The effects of the whole operation $bs_2 + ps$ are described by a unitary transformation

$$\begin{aligned} |\psi_u\rangle &\rightarrow |\psi_u(\theta, \varphi)\rangle = U(\theta, \varphi)|\psi_u\rangle, \\ |\psi_d\rangle &\rightarrow |\psi_d(\theta, \varphi)\rangle = U(\theta, \varphi)|\psi_d\rangle, \end{aligned} \quad ; \quad U(\theta, \varphi) = \begin{pmatrix} e^{-i\varphi} \sin \theta & e^{-i\varphi} \cos \theta \\ \cos \theta & -\sin \theta \end{pmatrix} \quad (5.4)$$

where $\cos^2 \theta$ and $\sin^2 \theta$ are the reflection and transmission probabilities for bs_2 (the analogue of q and p); their values can be modified by changing the angle θ , that is by changing

the physical properties of the beam splitter bs_2 (the parameters p and q , characterizing the first beam splitter, are instead kept fixed). The initial statistical operator thus evolves as follows:

$$\rho^{ab} \rightarrow (U(\theta, \varphi) \otimes \mathbf{I}_2) \rho^{ab}(t) (U^\dagger(\theta, \varphi) \otimes \mathbf{I}_2), \quad (5.5)$$

where $\mathbf{I}_2$ is the 2×2 identity in the translational Hilbert space $\mathfrak{H}_2^a$.

The exiting neutron beam in the state (5.5) then impinges on two counters (c_1 and c_2), equipped with spin-analyzers that can be set along any direction $\vec{n} = (n_x, n_y, n_z)$ in space. These counters are used to measure the relative frequencies with which neutrons are detected in one of the two exiting beams, with spin up or down along $\vec{n}$. These frequencies of counts, denoted by $\mathcal{O}_t^{j, \vec{n}}(\theta, \varphi)$ are given by mean values of the projectors $P_j(\theta, \varphi) \otimes Q_{\vec{n}}$ in $\mathfrak{H}_2^a \otimes \mathfrak{H}_2^b$ over $\rho^{ab}(t)$:

$$\mathcal{O}_t^{j, \vec{n}}(\theta, \varphi) \equiv \text{Tr}[\rho^{ab}(t) P_j(\theta, \varphi) \otimes Q_{\vec{n}}], \quad (5.6)$$

where

$$P_j(\theta, \varphi) \equiv U^\dagger(\theta, \varphi) P_j U(\theta, \varphi), \quad (5.7)$$

with $j = 1, 2$, are the spatial projectors in (5.3), whereas $Q_{\vec{n}} \equiv |\uparrow_{\vec{n}}\rangle\langle\uparrow_{\vec{n}}|$ is the projector over the eigenspace spanned by the state $|\uparrow_{\vec{n}}\rangle$ with spin-up along the $\vec{n}$ -direction.

Remark 5.3 The last beam splitter together with the counters plus spin analyzers constitutes the measuring device, since it sets the values of the angles θ and φ , that can be varied by the experimenter.

Neutron interferometry has proved to be an extremely powerful tool to investigate gravitational, inertial and phase shifting effects occurring inside the interferometer [131, 137]: they are described by a time-evolution for the spatial density matrix. Instead, in what follows, we shall assume a dissipative time-evolution to affect the spin degrees of freedom, because of the presence of a stochastic magnetic field inside the apparatus [39, 125]. The neutron beam thus behaves like an open system with time-evolution given by $\Lambda_t = \mathbf{I}_2 \otimes \Lambda_t^b$, Λ_t^b being the dynamical map describing the non-trivial evolution in the spin sector. Explicitly, while traversing the interferometer, the neutron beam state changes according to $\rho^{ab} \rightarrow \rho^{ab}(t) = \mathbf{I}_2 \otimes \Lambda_t^b[\rho^{ab}]$. Neutron interferometry can then be used to investigate the property of complete positivity. In particular, the form of Λ_t is the one appearing in the definition of complete positivity in Section 1.2: to guarantee the positivity of $\rho^{ab}(t) = \Lambda_t[\rho^{ab}(0)]$ for any time t we need necessarily a completely positive map Λ_t^b .

In Section 5.4 we shall show explicitly that lack of complete positivity generates inconsistencies in the physical interpretation of the statistical operator $\rho^{ab}(t)$, that is the appearance of negative eigenvalues. Noticeably, these eigenvalues can be directly accessed in experiments by setting appropriately the angles θ and φ and the direction $\vec{n}$. Indeed, the

entries of $\rho^{ab}(t)$ can be expressed by means of the frequencies measured at the counters. Writing $\rho^{ab}(t) = \sum_{i,j=1}^4 \rho_{ij}(t) P_i \otimes Q_j$, with P_i and Q_j as in (5.3), it turns out that

$$\begin{aligned} \rho_{11}(t) &= \mathcal{O}_t^{1,z}(0,0), & \rho_{12}(t) &= \mathcal{O}_t^{1,-z}(0,0), \\ \rho_{21}(t) &= \mathcal{O}_t^{2,z}(0,0), & \rho_{22}(t) &= \mathcal{O}_t^{2,-z}(0,0); \end{aligned} \quad (5.8)$$

the remaining 12 entries ρ_{ij} cannot be evaluated directly since in these cases the $P_i \otimes Q_j$ operators are not Hermitian; nevertheless they can be written by using suitable combinations of observables involving the following projectors, see (5.7):

$$\begin{aligned} P_{\pm} &\equiv P_{1,2} \left(\frac{\pi}{4}, 0 \right), & P_{\pm i} &\equiv P_{1,2} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right), \\ Q_{\pm x} &\equiv \frac{1}{2} (|\uparrow_z\rangle \pm |\downarrow_z\rangle)(\langle\uparrow_z| \pm \langle\downarrow_z|), & Q_{\pm y} &\equiv \frac{1}{2} (|\uparrow_z\rangle \pm i|\downarrow_z\rangle)(\langle\uparrow_z| \mp i\langle\downarrow_z|). \end{aligned} \quad (5.9)$$

Then, from

$$P_3 = \frac{1}{2} [(P_+ - P_-) + i(P_{+i} - P_{-i})] = P_4^*, \quad Q_3 = \frac{1}{2} [(Q_x - Q_{-x}) + i(Q_y - Q_{-y})] = Q_4^* \quad (5.10)$$

the expressions of the remaining entries can be explicitly evaluated (see Appendix E); for example

$$\begin{aligned} \rho_{43}(t) &= \frac{1}{4} \left\{ \left[\mathcal{O}_t^{1,x} \left(\frac{\pi}{4}, 0 \right) - \mathcal{O}_t^{1,-x} \left(\frac{\pi}{4}, 0 \right) \right] + i \left[\mathcal{O}_t^{1,y} \left(\frac{\pi}{4}, 0 \right) - \mathcal{O}_t^{1,-y} \left(\frac{\pi}{4}, 0 \right) \right] + \right. \\ &\quad - \left[\mathcal{O}_t^{2,x} \left(\frac{\pi}{4}, 0 \right) - \mathcal{O}_t^{2,-x} \left(\frac{\pi}{4}, 0 \right) \right] - i \left[\mathcal{O}_t^{2,y} \left(\frac{\pi}{4}, 0 \right) - \mathcal{O}_t^{2,-y} \left(\frac{\pi}{4}, 0 \right) \right] + \\ &\quad - \left[\mathcal{O}_t^{1,y} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) - \mathcal{O}_t^{1,-y} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) \right] + i \left[\mathcal{O}_t^{1,x} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) - \mathcal{O}_t^{1,-x} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) \right] + \\ &\quad \left. + \left[\mathcal{O}_t^{2,y} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) - \mathcal{O}_t^{2,-y} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) \right] - i \left[\mathcal{O}_t^{2,x} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) - \mathcal{O}_t^{2,-x} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) \right] \right\} \quad (5.11) \end{aligned}$$

Also these entries are thus, in principle, accessible to experimental measures.

5.2 The stochastic thermal bath

We assume that, inside the interferometer, the neutron beam is affected by a weak stochastic magnetic field; as a result, the dynamics is that of a dissipative open system.

We use the formalism introduced in Section 1.7; namely, the spin degree of freedom is described by a 2×2 stochastic density matrix $\Sigma(t)$, satisfying the time-dependent Liouville-Von Neumann equation

$$\begin{aligned} \frac{d}{dt}\Sigma(t) &= (L_0 + L_I(t))[\Sigma(t)], \\ L_0[\cdot] &\equiv -i[H_0, \cdot]; \quad L_I(t)[\cdot] \equiv -i[H_I(t), \cdot] \end{aligned} \quad (5.12)$$

where $H_0 = \omega_0 \sigma_z/2$; $H_I(t) = \sum_{k=1}^3 V_k(t) \sigma_k$ is the fluctuating part of the Hamiltonian and $V_k(t)$, $k = 1, \dots, 3$, are proportional to Gaussian stochastic magnetic field components. We assume $\langle V_k(t) \rangle = 0 \ \forall k$ and the two-point correlation functions to be real, stationary and symmetric, that is

$$h_{ij}(t-s) = \langle V_i(t) V_j(s) \rangle = h_{ij}^*(t-s) = h_{ji}(s-t). \quad (5.13)$$

The covariance matrix, defined as $\mathcal{W}(t) \equiv [h_{ij}(t)]$, is thus real, stationary and symmetric.

The effective 2×2 spin density matrix of the neutron beam when it impinges upon the second beam splitter is given by $\rho(t) = \langle \Sigma(t) \rangle$, where an average over the noise has been performed; its equation of motion has the form

$$\dot{\rho}(t) = -i[H_0, \rho(t)] + \sum_{k,j=1}^3 d_{kj}(t) [\sigma_k, [\sigma_j, \rho(t)]] \quad (5.14)$$

with

$$d_{kj}(t) = - \sum_{l=1}^3 \int_0^t h_{kl}(s) u_{lj}(-s) ds \quad (5.15)$$

where the U_{lj} are the entries of the orthogonal matrix

$$\mathcal{U}(t) = [u_{ij}(t)] = \begin{pmatrix} \cos \omega_0 t & -\sin \omega_0 t & 0 \\ \sin \omega_0 t & \cos \omega_0 t & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad (5.16)$$

such that $\sigma_l(-s) = \sum_{j=1}^3 u_{lj}(-s) \sigma_j$. Equation (5.14) is the form assumed by the general equation (1.70) in the two-dimensional case and we refer to Section 1.7 for details. The only assumptions required to extract the reduced dynamics (5.14) are the Gaussian nature of the stochastic magnetic field and the initial decoupling between spin and noise, that is $\rho(0) = \langle \Sigma(0) \rangle = \Sigma(0)$.

Our assumptions on the correlation functions make the 3×3 matrix $D(t) \equiv [d_{ij}(t)]$ real; thus it can be written as the sum of a symmetric and a skew-symmetric part. Defining the two 3×3 matrices

$$D^{S,A}(t) \equiv \frac{1}{2}[D(t) \pm D^T(t)] = [d_{ij}^{S,A}(t)], \quad (5.17)$$

where T means transposition, the dissipative part in equation (5.14) can be written as the sum of a Hamiltonian contribution plus a purely dissipative one. Then

$$\dot{\rho}(t) = -i \left[H_0 - \sum_{i,j,k=1}^3 d_{ij}^A(t) \epsilon_{ijk} \sigma_k, \rho(t) \right] + 2 \sum_{i,j=1}^3 d_{ij}^S(t) \left(\frac{1}{2} \{ \sigma_i \sigma_j, \rho(t) \} - \sigma_j \rho(t) \sigma_i \right). \quad (5.18)$$

In the limit of weak stochastic magnetic field, it is possible to perform a Markovian approximation for the reduced dynamics (5.18) as discussed in Chapter 1; in practice it is usually performed by extending to $+\infty$ of the upper limit of the integral in (5.15). The matrices D^S and D^A become then time-independent and we obtain dynamical equations of motion in Lindblad form:

$$\dot{\rho}(t) = -i \left[H_0 - \sum_{i,j,k=1}^3 d_{ij}^A \epsilon_{ijk} \sigma_k, \rho(t) \right] + \frac{1}{2} \sum_{i,j=1}^3 c_{ij} \left(\sigma_i \rho(t) \sigma_j - \frac{1}{2} \{ \sigma_j \sigma_i, \rho(t) \} \right), \quad (5.19)$$

where $C \equiv [c_{ij}] = -4D^S$.

Remark 5.4 Following the result of Theorem 1.5, the dynamics (5.19) is completely positive if and only if the matrix $C = [c_{ij}]$ is positive definite. In general, the brute force Markovian approximation adopted above may provide a Markovian equation (5.19) that generates time-evolutions completely positive, simply positive or even not-positive, depending on the behavior of the correlation functions, in (5.15).

Equation (5.19) can be rewritten using the vector representation $|\rho(t)\rangle$ for density matrices $\rho(t)$ presented in Section 2.1. The Schrödinger-like equation that follows is:

$$\frac{d}{dt} |\rho(t)\rangle = (\mathcal{H} + \mathcal{D}) |\rho(t)\rangle \quad (5.20)$$

where $\mathcal{H}$ and $\mathcal{D}$ are 4×4 real matrices representing the Hamiltonian, respectively dissipative contribution in (5.19):

$$\mathcal{H} = -2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & h_3 & -h_2 \\ 0 & -h_3 & 0 & h_1 \\ 0 & h_2 & -h_1 & 0 \end{pmatrix}, \quad \mathcal{D} = -2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & a & b & c \\ 0 & b & \alpha & \beta \\ 0 & c & \beta & \gamma \end{pmatrix} \quad (5.21)$$

where

$$h_1 = 2d_{32}^A, \quad h_2 = 2d_{13}^A, \quad h_3 = \frac{\omega_0}{2} + 2d_{21}^A \quad (5.22)$$

and

$$\begin{aligned}
a &= -2(d_{22}^S + d_{33}^S), & b &= 2d_{12}^S, \\
\alpha &= -2(d_{11}^S + d_{33}^S), & \beta &= 2d_{23}^S, \\
\gamma &= -2(d_{11}^S + d_{22}^S), & c &= 2d_{13}^S.
\end{aligned} \tag{5.23}$$

Notice that, in this parametrization

$$C = \begin{pmatrix} \alpha + \gamma - a & -2b & -2c \\ -2b & a + \gamma - \alpha & -2\beta \\ -2c & -2\beta & a + \alpha - \gamma \end{pmatrix}. \tag{5.24}$$

From Theorems 2.4 and 2.5, it immediately follows that

1. The reduced dynamics generated by (5.19) is positive if and only if the parameters entering in $\mathcal{D}$ satisfy the inequalities:

$$\begin{cases} a \geq 0 \\ \alpha \geq 0 \\ \gamma \geq 0 \end{cases} ; \begin{cases} \alpha\gamma \geq b^2 \\ a\gamma \geq c^2 \\ a\alpha \geq \beta^2 \end{cases} ; a\alpha\gamma + 2bc\beta - \alpha c^2 - \gamma b^2 - a\beta^2 \geq 0; \tag{5.25}$$

2. The reduced dynamics generated by (5.19) is completely positive if and only if the parameters entering in $\mathcal{D}$ satisfy the inequalities:

$$\begin{cases} 2R \equiv \alpha + \gamma - a \geq 0 \\ 2S \equiv a + \gamma - \alpha \geq 0 \\ 2T \equiv a + \alpha - \gamma \geq 0 \end{cases} ; \begin{cases} RS \geq b^2 \\ RT \geq c^2 \\ ST \geq \beta^2 \end{cases} ; RST - 2bc\beta - Sc^2 - R\beta^2 - Tb^2 \geq 0 \tag{5.26}$$

that is, if C is positive.

5.3 Three different stochastic backgrounds

In order to investigate the notion of complete positivity in more detail, we consider different decaying properties of the covariance matrix $\mathcal{W}(t)$. We focus our attention on three particular choices, reproducing equations of motion usually employed in the study of open systems [125, 126, 27]; the aim of this work is to give again direct evidence to the fact that the absence of complete positivity in the reduced dynamics generates necessarily physical inconsistencies in the interpretation of the formalism. In particular, these reduced dynamics may be experimentally realized by suitable stochastic magnetic fields and thus describe concrete physical contexts.

5.3.1 White noise

The first case is that with white noise correlations for the stochastic magnetic field:

$$h_{ij}(t) = h_{ij}\delta(t), \quad (5.27)$$

where h_{ij} ($i, j = 1, \dots, 3$) are time-independent, such that $h_{ij} = h_{ji} = h_{ij}^*$.

The covariance matrix $\mathcal{W}(t)$ is thus time-independent, positive and symmetric. Consequently $D^A = 0$ and $D^S = -\mathcal{W} \leq 0$. Then $C \geq 0$ and the evolution is automatically completely positive; the dissipative coefficients are related to the correlation functions by

$$\begin{aligned} a &\equiv -2(h_{22} + h_{33}), & b &\equiv 2h_{12}, \\ \alpha &\equiv -2(h_{11} + h_{33}), & \beta &\equiv 2h_{23}, \\ \gamma &\equiv -2(h_{11} + h_{22}), & c &\equiv 2h_{13}. \end{aligned} \quad (5.28)$$

5.3.2 Diagonal covariance matrix

In this case, we suppose $\mathcal{W}(t)$ diagonal with

$$h_{11}(t) = h_{22}(t) = g^2 B_1^2 e^{-\lambda|t|}, \quad h_{33}(t) = g^2 B_3^2 e^{-\mu|t|} \quad (5.29)$$

where the B_i 's are the time-independent components of the magnetic field, g is a constant proportional to the neutron magnetic moment and λ, μ are decay coefficients.

We find

$$D^A = -\frac{g^2 \omega_0 B_1^2}{\lambda^2 + \omega_0^2} \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} \gamma & 0 & 0 \\ 0 & \gamma & 0 \\ 0 & 0 & 2a - \gamma \end{pmatrix}, \quad (5.30)$$

where

$$a \equiv \frac{2g^2 B_3^2}{\mu} + \frac{2g^2 B_1^2 \lambda}{\lambda^2 + \omega_0^2}, \quad \gamma \equiv \frac{4g^2 B_1^2 \lambda}{\lambda^2 + \omega_0^2}. \quad (5.31)$$

The 4×4 matrices $\mathcal{H}$ and $\mathcal{D}$, in (5.20) take the form

$$\mathcal{H} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & -\omega & 0 \\ 0 & \omega & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \mathcal{D} = -2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & a & 0 & 0 \\ 0 & 0 & a & 0 \\ 0 & 0 & 0 & \gamma \end{pmatrix} \quad (5.32)$$

where $\omega \equiv \omega_0 + \frac{4g^2 \omega_0 B_1^2}{\lambda^2 + \omega_0^2}$.

Therefore, the entries of the spin density matrix $\rho(t)$ satisfy the Redfield-Bloch [4] equations of motion:

$$\begin{aligned}\dot{\rho}_1(t) &= -\gamma\rho_1(t) + \gamma\rho_2(t), & \dot{\rho}_2(t) &= -\dot{\rho}_1(t), \\ \dot{\rho}_3(t) &= -i\omega\rho_3(t) - 2a\rho_3(t), & \dot{\rho}_4(t) &= [\dot{\rho}_3(t)]^*,\end{aligned}\tag{5.33}$$

that can be explicitly solved yielding

$$\begin{aligned}\rho_1(t) &= \frac{1}{2} [(1 + e^{-2\gamma t}) \rho_1(0) + (1 - e^{-2\gamma t}) \rho_2(0)], \\ \rho_2(t) &= \frac{1}{2} [(1 - e^{-2\gamma t}) \rho_1(0) + (1 + e^{-2\gamma t}) \rho_2(0)], \\ \rho_3(t) &= e^{-(2a-i\omega)t} \rho_3(0), \\ \rho_4(t) &= e^{-(2a+i\omega)t} \rho_4(0).\end{aligned}\tag{5.34}$$

Since a and γ are real and positive, the positivity of the dynamics is assured; complete positivity requires $C \geq 0$ and then $2a - \gamma \geq 0$. From (5.34) we observe that diagonal and off-diagonal elements of ρ decay with different relaxing times, T_1 and T_2 respectively; they are the reciprocals of the coefficients 2γ and $2a$. Then the condition of complete positivity reads $1/T_2 \geq 1/2T_1$ [34, 125]. In [125, 126] it is shown that this typical order relation can be reversed by setting $B_3 = 0$ and keeping fourth-order terms in the general master equation (1.67) leading to (5.14). In such a case we have a positive, but not completely positive time-evolution.

5.3.3 Single component field correlation

In this case the magnetic field is supposed to have only the x -component: $V_1(t) \neq 0$, $V_2(t) = V_3(t) = 0$ with

$$h_{11}(t) = g^2 B^2 e^{-\lambda|t|};\tag{5.35}$$

consequently

$$D^A = -\frac{g^2 \omega_0 B^2}{2(\lambda^2 + \omega_0^2)} \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} 2\alpha & -2b & 0 \\ -2b & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}\tag{5.36}$$

with

$$\alpha \equiv \frac{2g^2 B^2 \lambda}{\lambda^2 + \omega_0^2}, \quad b \equiv -\frac{g^2 B^2 \omega_0}{\lambda^2 + \omega_0^2}.\tag{5.37}$$

In the vector representation (5.20), the dissipative matrix has the form

$$\mathcal{D} = -2 \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & b & 0 \\ 0 & b & \alpha & 0 \\ 0 & 0 & 0 & \alpha \end{pmatrix} \quad (5.38)$$

whereas the Hamiltonian contribution $\mathcal{H}$, expressed as in equation (5.32), is characterized by $\omega \equiv \omega_0 + \frac{2g^2\omega_0 B^2}{\lambda^2 + \omega_0^2}$. The corresponding Redfield-Bloch equations are

$$\begin{aligned} \dot{\rho}_1(t) &= -\alpha\rho_1(t) + \alpha\rho_2(t), & \dot{\rho}_2(t) &= -\dot{\rho}_1(t), \\ \dot{\rho}_3(t) &= -i\omega\rho_3(t) - \alpha\rho_3(t) + \alpha\rho_4(t) + 2ib\rho_4(t), & \dot{\rho}_4(t) &= [\dot{\rho}_3(t)]^*, \end{aligned} \quad (5.39)$$

with solutions

$$\begin{aligned} \rho_1(t) &= \frac{1}{2} \left[(1 + e^{-2\alpha t}) \rho_1(0) + (1 - e^{-2\alpha t}) \rho_2(0) \right], \\ \rho_2(t) &= \frac{1}{2} \left[(1 - e^{-2\alpha t}) \rho_1(0) + (1 + e^{-2\alpha t}) \rho_2(0) \right], \\ \rho_3(t) &= e^{-\alpha t} \left[\rho_4(0) \cosh \delta t + \frac{\alpha - 2ib}{\delta} \rho_4(0) \sinh \delta t + \frac{i\omega}{\delta} \rho_3(0) \sinh \delta t \right], \\ \rho_4(t) &= e^{-\alpha t} \left[\rho_3(0) \cosh \delta t + \frac{\alpha + 2ib}{\delta} \rho_3(0) \sinh \delta t - \frac{i\omega}{\delta} \rho_4(0) \sinh \delta t \right], \end{aligned} \quad (5.40)$$

where $\delta \equiv \sqrt{\alpha^2 + 4b^2 - \omega^2}$.

In this case, positivity and complete positivity asks for $b = 0$ (and thus $\omega = \omega_0$), while for $b \neq 0$ the time-evolution is not even positive.

5.4 Positivity and complete positivity

The covariance matrices in Subsections 5.3.2 and 5.3.3 do not guarantee the complete positivity of the time evolution. In this section we address the problems that arise in these cases.

In the diagonal covariance matrix approach, let us consider the maximally entangled state, $p = -q = 1/\sqrt{2}$ in (5.1); it evolves into

$$\rho^{ab}(t) = \frac{1}{2} [P_1 \otimes Q_2(t) + P_2 \otimes Q_1(t) - P_3 \otimes Q_4(t) - P_4 \otimes Q_3(t)] \quad (5.41)$$

where $Q_i(t)$ are computed from equations (5.34). The 4×4 statistical operator has the form

$$\rho^{ab}(t) = \begin{pmatrix} E_-(t) & 0 & 0 & 0 \\ 0 & E_+(t) & F(t) & 0 \\ 0 & F^*(t) & E_+(t) & 0 \\ 0 & 0 & 0 & E_-(t) \end{pmatrix}, \quad (5.42)$$

where

$$E_{\pm}(t) \equiv \frac{1}{4} (1 \pm e^{-2\gamma t}), \quad F(t) \equiv -\frac{1}{2} e^{-(2a-i\omega)t}, \quad (5.43)$$

the eigenvalues of $\rho^{ab}(t)$ at the exit of the interferometer are thus

$$\begin{aligned} \lambda_{1,2}(t) &= \frac{1}{4} (1 - e^{-2\gamma t}), \\ \lambda_{3,4}(t) &= \frac{1}{4} (1 + e^{-2\gamma t} \pm 2e^{-2at}). \end{aligned} \quad (5.44)$$

Observe that $\lambda_4(0) = 0$ and $d\lambda_4(0)/dt = (2a - \gamma)/2$; then, by continuity, if $a < \gamma/2$, there is whole range of t where $\lambda_4(t) < 0$ and $\rho^{ab}(t)$ loses its physical meaning. On the other hand, if the complete positivity condition is fulfilled, $\lambda_4(t) \geq 0$ for any time t and no problem arises.

Similar conclusions hold for the single component field correlation case.

Remark 5.5 It is necessary an entangled initial state to see the problems generated by positive dynamics that are not completely positive. Indeed, suppose to have a factorized initial state, i.e. in the form

$$\rho^{ab} = \sum_{i,j} c_{ij} \rho_i^a \otimes \rho_j^b > 0$$

with $c_{ij} \geq 0$, and let it evolve following the factorized dynamics $\mathbf{I}_2 \otimes \Lambda_b^s$; then the statistical operator at time t is positive for any t , because of the positivity of Λ_b^s :

$$\rho^{ab}(t) = (\mathbf{I}_2 \otimes \Lambda_b^s)[\rho^{ab}] = \sum_{i,j} c_{ij} \rho_i^a \otimes \Lambda_b^s[\rho_j^b] \geq 0$$

Remark 5.6 Besides the appearance of negative eigenvalues, we observe that, in the single component field correlation case, for $\delta > \alpha$, $\rho_3(t)$ and $\rho_4(t)$ diverge exponentially, see (5.40); the positivity (and complete positivity) condition, $b = 0$, avoids this unacceptable behavior.

Remark 5.7 Although apparently formal, these results are far from being academic. Indeed, as already stressed at the end of Section 5.1, the entries of ρ are in principle directly




Figure 5.2: Three different behaviors for $\lambda_4(t)$: for $a > \gamma/2$ (green line), for $a = \gamma/2$ (blue line) and for $a < \gamma/2$ (red line). We see that in the latter case there is a whole range of t where $\lambda_4(t) < 0$ and then the density matrix loses its physical meaning; indeed, the time-evolution is not completely positive.

accessible to the experiment. For example, in the diagonal covariance matrix case, it follows from eq. (5.8) that $E_-(t) = \mathcal{O}_t^{1,z}(0,0)$ and $E_+(t) = \mathcal{O}_t^{2,z}(0,0)$ while $F(t)$ coincides with the expression in equation (5.11).

By modulating a background magnetic field close to the stochastic properties investigated in the previous three cases, one might reproduce experimentally the conditions for three different reduced dynamics and check their consequences.

Then, one may conclude that reduced, Markovian time evolutions Λ_b^s must be not only positive, but also completely positive, since the lack of any of these constraints results in experimentally detectable inconsistencies.

The use of one reduced dynamics instead of another depends on the Markovian approximation used to derive it and whether, given the properties of the stochastic field, it was justified or not. It thus seems appropriate to conclude that, whenever a semigroup composition law is expected, the physically appropriate Markovian approximations are those which lead to reduced dynamics consisting of completely positive maps [35].

5.5 Noncontextuality: the Clauser-Horne-Shimony-Holt inequality

Quantum Mechanics has been shown to be a non-local and contextual theory, as it violates the so-called Bell inequalities that have to be necessarily satisfied by local, realistic and

noncontextual theories [127].

Neutron interferometry is an optimal tool for investigating noncontextuality [129]; indeed in this case the entanglement does not involve spatially separated parts, as in the usual tests exhibiting violations of Bell-like inequalities which focus on locality.

It is customary to use a particular form of Bell inequality, due to Clauser, Horne, Shimony and Holt [128]. Using the previously defined projectors in configuration and spin space, one constructs the observables

$$A(\theta, \varphi) \equiv P_1(\theta, \varphi) - P_2(\theta, \varphi), \quad B(\vec{n}) \equiv Q_{\vec{n}} - Q_{-\vec{n}}. \quad (5.45)$$

These observables commute as translational and spin observables do; they are called dichotomic since their eigenvalues are ± 1 . Choosing two different values for both θ and φ and two distinct directions for the vector $\vec{n}$, we obtain the set of 4 observables $A(\theta_i, \varphi_i)$ and $B(\vec{n}_j)$ with $i, j = 1, 2$.

From the mean values

$$\begin{aligned} C_t(\theta, \varphi; \vec{n}) &\equiv \text{Tr}[\rho^{ab}(t)A(\theta, \varphi) \otimes B(\vec{n})] = \\ &= \mathcal{O}_t^{1, \vec{n}}(\theta, \varphi) + \mathcal{O}_t^{2, -\vec{n}}(\theta, \varphi) - \mathcal{O}_t^{1, -\vec{n}}(\theta, \varphi) - \mathcal{O}_t^{2, \vec{n}}(\theta, \varphi), \end{aligned} \quad (5.46)$$

with four possible configurations of the control parameters θ , φ and $\vec{n}$, the Clauser-Horne-Shimony-Holt inequality is obtained [128]:

$$|C_t(\theta_1, \varphi_1; \vec{n}_1) + C_t(\theta_1, \varphi_1; \vec{n}_2) + C_t(\theta_2, \varphi_2; \vec{n}_1) - C_t(\theta_2, \varphi_2; \vec{n}_2)| \leq 2. \quad (5.47)$$

This inequality holds if we suppose the two possible outcomes ± 1 of a measurement of the observables $A(\theta_i, \varphi_i)$ and $B(\vec{n}_j)$ are predetermined by the state $\rho^{ab}(t)$ independently of which other commuting observable is simultaneously measured (for example, the value of $A(\theta_1, \varphi_1)$ is unchanged if we measure simultaneously either $B(\vec{n}_1)$ or $B(\vec{n}_2)$). A violation of this relation implies necessarily the contextuality of the underlying theory. Equation (5.46) shows that this analysis can be directly performed by means of the frequencies of counts in the interferometric apparatus.

Remark 5.8 In the configuration described in [129, 130] the state ρ^{ab} is not subject to any dynamics inside the interferometer, but in our context it does: this is the reason why we have considered an explicit time dependence in the $C_t(\theta, \varphi, \vec{n})$ mean values.

For the maximally entangled initial state ($p = -q = 1/\sqrt{2}$) these quantities take the form

$$\begin{aligned} C_t(\theta, \varphi; \vec{n}) &= \sin^2 \theta \text{Tr}_b[Q_2(t)B(\vec{n})] + \cos^2 \theta \text{Tr}_b[Q_1(t)B(\vec{n})] + \\ &\quad - \sin 2\theta \text{Re} (e^{-i\varphi} \text{Tr}_b[Q_4(t)B(\vec{n})]) \end{aligned} \quad (5.48)$$

where Tr_b is the trace over the spin degree of freedom.

In vectorial representation, the time-evolution in the spin sector reads $|\rho(t)\rangle = \mathcal{G}_t|\rho(0)\rangle$; in the following we shall consider the first order approximation of $\mathcal{G}_t$ in the dissipative parameters, supposed to be small with respect to the parameter characterizing the standard Hamiltonian evolution, ω_0 (for the expression of the entries of $\mathcal{G}_t$, see Section 2.4). We introduce two vectors, $\vec{G}(t)$ and $\vec{F}(t)$, whose components are defined by

$$G_i(t) \equiv (\mathcal{G}_t)_{i3}, \quad F_i(t) \equiv (\mathcal{G}_t)_{i1} - i(\mathcal{G}_t)_{i2}, \quad i = 1, 2, 3. \quad (5.49)$$

Then, the traces in (5.48) can be written as:

$$\text{Tr}_b[Q_{1,2}(t)B(\vec{n})] = \pm \vec{G}(t) \cdot \vec{n}, \quad \text{Tr}_b[Q_4(t)B(\vec{n})] = \vec{F}(t) \cdot \vec{n}; \quad (5.50)$$

whence the mean values (5.2) read

$$C_t(\theta, \varphi; \vec{n}) = \vec{n} \cdot \left[\vec{G}(t) \cos 2\theta - \text{Re} \left(e^{-i\varphi} \vec{F}(t) \right) \sin 2\theta \right]. \quad (5.51)$$

We will now discuss the explicit behaviour of $C_t(\theta, \varphi; \vec{n})$ in the three cases introduced in Section 5.3, to leading order in the dissipative parameters.

5.5.1 White noise

Though analytic expressions of $\mathcal{G}_t$ are obtainable in the general case of a stochastic magnetic field with white noise correlations, these are rather involved and scarcely illuminating. More conveniently, we consider the expression of the time evolver $\mathcal{G}_t$ to first order in the parameters, appearing in the dissipative part $\mathcal{D}$ of (5.21), assumed to be small. One calculates

$$\begin{aligned} G_1(t) &= -\frac{4|C|}{\omega_0} \sin \frac{\omega_0 t}{2} \cos \left(\frac{\omega_0 t}{2} + \phi_C \right), \\ G_2(t) &= \frac{4|C|}{\omega_0} \sin \frac{\omega_0 t}{2} \sin \left(\frac{\omega_0 t}{2} + \phi_C \right), \\ G_3(t) &= e^{-2\gamma t}, \end{aligned} \quad (5.52)$$

where $|C|^2 = c^2 + \beta^2$ and $\tan \phi_C = \beta/c$;

$$\begin{aligned} \text{Re} [e^{-i\varphi} F_1(t)] &= e^{-(a+\alpha)t} \cos(\omega_0 t - \varphi) + \frac{|B|}{\omega_0} \sin \omega_0 t \cos(\varphi + \phi_B), \\ \text{Re} [e^{-i\varphi} F_2(t)] &= e^{-(a+\alpha)t} \sin(\omega_0 t - \varphi) - \frac{|B|}{\omega_0} \sin \omega_0 t \sin(\varphi - \phi_B), \\ \text{Re} [e^{-i\varphi} F_3(t)] &= -\frac{4|C|}{\omega_0} \sin \frac{\omega_0 t}{2} \cos \left(\frac{\omega_0 t}{2} - \varphi - \phi_C \right), \end{aligned} \quad (5.53)$$

where $|B|^2 = (a - \alpha)^2 + 4b^2$ and $\tan \phi_B = 2b/(\alpha - a)$. It thus follows that, in leading order with respect to a, b, c, α, β and γ

$$\begin{aligned}
C_t(\theta, \varphi; \vec{n}) = & n_x \left\{ -\frac{4|C|}{\omega_0} \cos 2\theta \sin \frac{\omega_0 t}{2} \cos \left(\frac{\omega_0 t}{2} + \phi_C \right) + \right. \\
& \left. - \sin 2\theta \left[e^{-(a+\alpha)t} \cos(\omega_0 t - \varphi) + \frac{|B|}{\omega_0} \sin \omega_0 t \cos(\varphi + \phi_B) \right] \right\} + \\
& + n_y \left\{ \frac{4|C|}{\omega_0} \cos 2\theta \sin \frac{\omega_0 t}{2} \sin \left(\frac{\omega_0 t}{2} + \phi_C \right) + \right. \\
& \left. - \sin 2\theta \left[e^{-(a+\alpha)t} \sin(\omega_0 t - \varphi) - \frac{|B|}{\omega_0} \sin \omega_0 t \sin(\varphi - \phi_B) \right] \right\} + \\
& + n_z \left\{ e^{-2\gamma t} \cos 2\theta + \frac{4|C|}{\omega_0} \sin 2\theta \sin \frac{\omega_0 t}{2} \cos \left(\frac{\omega_0 t}{2} - \varphi - \phi_C \right) \right\}.
\end{aligned} \tag{5.54}$$

5.5.2 Diagonal covariance matrix

For the stochastic magnetic fields with correlation functions as in (5.29) we have

$$\vec{G}(t) = (0, 0, e^{-2\gamma t}), \quad \vec{F}(t) = e^{-(2a-i\omega)t}(1, -i, 0), \tag{5.55}$$

whence

$$C_t(\theta, \varphi; \vec{n}) = n_z e^{-2\gamma t} \cos 2\theta - e^{-2at} \sin 2\theta [n_x \cos(\omega t - \varphi) + n_y \sin(\omega t - \varphi)]. \tag{5.56}$$

5.5.3 Single component field correlation

In the case of magnetic field with correlation functions as in (5.35) the computation is similar. One finds:

$$\begin{aligned}
\vec{G}(t) &= (0, 0, e^{-2\alpha t}), \\
\vec{F}(t) &= e^{-\alpha t} \left[\left(\cosh \delta t + i \frac{\omega}{\delta} \sinh \delta t \right) (1, -i, 0) + \frac{\alpha + 2ib}{\delta} \sinh \delta t (1, i, 0) \right],
\end{aligned} \tag{5.57}$$

whence, assuming $\delta > 0$, one obtains:

$$\begin{aligned}
C_t(\theta, \varphi; \vec{n}) = & n_z e^{-2\alpha t} \cos 2\theta + \\
& - e^{-\alpha t} \sin 2\theta \left\{ n_x \left[\left(\cosh \delta t + \frac{\alpha}{\delta} \sinh \delta t \right) \cos \varphi + \frac{\omega + 2b}{\delta} \sinh \delta t \sin \varphi \right] + \right. \\
& \left. + n_y \left[\left(-\cosh \delta t + \frac{\alpha}{\delta} \sinh \delta t \right) \sin \varphi + \frac{\omega - 2b}{\delta} \sinh \delta t \cos \varphi \right] \right\}.
\end{aligned} \tag{5.58}$$

The lack of positivity preservation which characterizes the time-evolution leading to equation (5.58) manifests itself in that the quantities $C_t(\theta, \varphi; \vec{n})$ diverge in t when $\delta > \alpha$.

Expressions (5.54), (5.56) and (5.58) agree with those used when there is no dissipation, namely putting $a = b = c = \alpha = \beta = \gamma = 0$:

$$C_t(\theta, \varphi; \vec{n}) = -n_x \sin 2\theta \cos(\omega_0 t - \varphi) - n_y \sin 2\theta \sin(\omega_0 t - \varphi) + n_z \cos 2\theta. \quad (5.59)$$

Notice that the unitary time-evolution generated by the Hamiltonian H_0 contributes to a time-varying redefinition of the angle φ .

Concerning the issue of complete positivity versus simple positivity, in expressions (5.54) and (5.56) the two possibilities result in different relaxation properties due to whether inequalities (5.26) or (5.25) are fulfilled. No physical inconsistencies may affect the mean values $C_t(\theta, \varphi; \vec{n})$; indeed, negative probabilities may result in negative mean values of positive observables only if the latter are entangled. In the case of the quantities involved in inequality (5.46), the observables are factorized, $P_{1,2}(\theta, \varphi) \otimes Q_{\vec{n}}$ and the positivity of their mean values is preserved even when the time evolution is only positive and not completely positive.

This can be seen as follows. By definition of dual map Λ_t^* , to the Schrödinger time-evolution $\rho^{ab}(t) = (\mathbf{I}_2 \otimes \Lambda_t)[\rho^{ab}(0)]$, there corresponds the Heisenberg time-evolution of observables $A(t) = (\mathbf{I}_2 \otimes \Lambda_t^*)[A]$,

$$\text{Tr}[(\mathbf{I}_2 \otimes \Lambda_t)[\rho^{ab}(0)]A] = \text{Tr}[\rho^{ab}(\mathbf{I}_2 \otimes \Lambda_t^*)[A]]. \quad (5.60)$$

The maps Λ_t^* , dual to Λ_t , form a semigroup of dynamical maps that transform positive observables into positive observables, if the Λ_t 's preserve the positivity of states. Consequently, even when the initial state ρ^{ab} is entangled and the Λ_t 's positivity preserving, but not completely positive, it turns out that

$$\text{Tr}[\rho^{ab}(t)P_{1,2}(\theta, \varphi) \otimes Q_{\vec{n}}] = \text{Tr}[\rho^{ab}P_j(\theta, \varphi) \otimes \Lambda_t^*[Q_{\vec{n}}]] \geq 0. \quad (5.61)$$

5.6 Discussion and conclusions

Complete positivity is a property of quantum time-evolutions which is enjoyed by the standard dynamics of closed quantum systems generated by Hamiltonian operators, but not automatically by the more general reduced dynamics describing time-evolution of open quantum systems in interaction with suitable environments. Complete positivity is intimately related to the phenomenon of quantum entanglement, between two different systems, but also between two different degrees of freedom of a same physical system.

In this chapter we have considered the two entangled degrees of freedom, translational and rotational, of a beam of neutrons travelling through an interferometric apparatus. We have studied the consequences of placing the interferometer in a stochastic, gaussian

magnetic field weakly coupled to the spin degree of freedom, providing an experimentally controllable environment. As explained in the three cases of Section 5.3, the same Markov approximation naively yields a semigroup of dynamical maps $\mathbf{I}_2 \otimes \Lambda_t^b$, where only the spin degree of freedom evolves in time; by varying the decay properties of the external field correlations, these maps turn out to be alternatively completely positive, simply positive, not even positive.

The noncontextuality tests proposed in [129, 130] are based on the Clauser-Horne-Shimony-Holt inequality (5.47) without time-dependence. The presence of a fluctuating magnetic field induces relaxation on the spin degree of freedom with strength and properties depending on those of the field. Typically, the mean values in the inequality are damped and make it more difficult to be violated. However, in presence of stochastic fields yielding reduced dynamics that do not preserve positivity, the inequality might be dramatically violated because of possible mean values diverging in time.

This latter possibility is a manifestation of the fact that any physically consistent time-evolution Λ_t^b must preserve the positivity of spin states in order that the eigenvalues of the corresponding spin density matrices might at any time be used as probabilities, in agreement with the statistical interpretation of quantum mechanics. If Λ_t^b preserves the trace of spin density matrices, but not their positivity, spin states may evolve in time in such a way that some of their eigenvalues become negative, while others greater than 1, without upper bounds. It is this physically unacceptable phenomenon that leads to diverging mean values.

The request of positivity preservation by the maps Λ_t^b with respect to spin states is thus unexcapable, but it is not enough to avoid physical inconsistencies when the time-evolution maps $\mathbf{I}_2 \otimes \Lambda_t^b$ act on states ρ^{ab} with correlations between spin and translational degrees of freedom.

Inequality (5.47) does reveal the difference between completely positive and simply positivity preserving Λ_t^b , but only as long as the relaxation characteristic are concerned, without any further effect (as the divergence of some contributions to the inequality). In fact, the positive observables in (5.46) are factorized, that is they incorporate no entanglement between the translational and spin degree of freedom. Even if the initial state does incorporate entanglement, it nevertheless follows that the mean values of factorized observables remain positive and bounded.

However, the interferometric apparatus proposed in [129, 130] might also be used to measure the entries of the states of the neutron beam at the exit of the interferometer. In this way, one might have access to the spectrum of an initially entangled state after being subjected to the effects of the stochastic magnetic field.

In the case of fluctuating fields yielding reduced dynamics that preserve positivity, but are not completely positive, the theoretical predictions indicate the appearance of negative eigenvalues, that is of negative probabilities, in the spectrum of the entangled exiting state. The fact that they are, in line of principle, detectable experimentally, does not allow to dismiss such an occurrence as practically negligible. Rather, it forces to reconsider the Markov approximation used to derive the time-evolution and to select as

physically consistent only those providing completely positive reduced dynamics.

Appendix A

Dissipative parameters in the eigenstates basis

For an unstable system, the dissipative parameters entering into (2.46) are defined by:

$$\begin{aligned}\lambda_1 &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -w(1 + |r_1|^2)(1 - |r_2|^2) - 2(1 + |r_1|^2) \operatorname{Re}(Ur_2) \right\}, \\ \lambda_2 &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -w(1 + |r_2|^2)(1 - |r_1|^2) + 2(1 + |r_2|^2) \operatorname{Re}(Ur_1) \right\}, \\ \sigma &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -w(1 + |r_2|^2)(1 - |r_2|^2) - 2(1 + |r_2|^2) \operatorname{Re}(Ur_2) \right\}, \\ \xi &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -w(1 + |r_1|^2)(1 - |r_1|^2) + 2(1 + |r_1|^2) \operatorname{Re}(Ur_1) \right\}, \\ \delta_1 &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ w(1 - r_1 r_2^*)(1 - |r_2|^2) + 2(1 - r_1 r_2^*) \operatorname{Re}(Ur_2) \right\}, \\ \delta_2 &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -w(1 + |r_2|^2)(1 + r_2 r_1^*) + (1 + |r_2|^2)(U^* r_1^* - Ur_2) \right\}, \\ \phi_1 &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ w(1 - r_1 r_2^*)(1 - |r_1|^2) - 2(1 - r_1 r_2^*) \operatorname{Re}(Ur_1) \right\}, \\ \phi_2 &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -w(1 + |r_1|^2)(1 + r_2 r_1^*) + (1 + |r_1|^2)(U^* r_1^* - Ur_2) \right\}, \\ \omega &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ w(1 - r_1 r_2^*)(1 + r_2 r_1^*) - (1 - r_1 r_2^*)(U^* r_1^* - Ur_2) \right\}, \\ \theta &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ w(1 - r_2 r_1^*)(1 + r_2 r_1^*) - (1 - r_2 r_1^*)(U^* r_1^* - Ur_2) \right\},\end{aligned}$$

$$\begin{aligned}
\Lambda &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ \gamma(1 - |r_1|^2)(1 - |r_2|^2) - 2A \operatorname{Re}(r_2 r_1^*) + 2 \operatorname{Re}(B r_1^* r_2^*) + \right. \\
&\quad \left. - 2(1 - |r_2|^2) \operatorname{Re}(C r_1^*) + 2(1 - |r_1|^2) \operatorname{Re}(C r_2^*) \right\}, \\
\Xi &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ \gamma(1 - |r_1|^2)^2 + 2A|r_1|^2 - 2 \operatorname{Re}(B r_1^{*2}) - 4(1 - |r_1|^2) \operatorname{Re}(C r_1^*) \right\}, \\
\Sigma &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ \gamma(1 - |r_2|^2)^2 + 2A|r_2|^2 - 2 \operatorname{Re}(B r_2^{*2}) + 4(1 - |r_2|^2) \operatorname{Re}(C r_2^*) \right\}, \\
\Delta &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -\gamma(1 - |r_2|^2)(1 + r_1 r_2^*) - A(|r_2|^2 - r_1 r_2^*) + B r_2^{*2} - B^* r_1 r_2 + \right. \\
&\quad \left. + (C^* r_1 - C r_2^*)(1 - |r_2|^2) - 2(1 + r_1 r_2^*) \operatorname{Re}(C r_2^*) \right\}, \\
\Phi &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -\gamma(1 - |r_1|^2)(1 + r_1 r_2^*) - A(|r_1|^2 - r_1 r_2^*) + B^* r_1^2 - B r_1^* r_2^* + \right. \\
&\quad \left. + (C^* r_1 - C r_2^*)(1 - |r_1|^2) + 2(1 + r_1 r_2^*) \operatorname{Re}(C r_1^*) \right\}, \\
\Omega &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -\gamma|1 + r_1 r_2^*|^2 - A(|r_1|^2 + |r_2|^2) - 2 \operatorname{Re}(B r_1^* r_2^*) + \right. \\
&\quad \left. + 2 \operatorname{Re} [C (r_1^* (1 - |r_2|^2) - r_2^* (1 - |r_1|^2))] \right\}, \\
\Theta &\equiv \frac{1}{|r_1 + r_2|^2} \left\{ -\gamma(1 + r_2 r_1^*)^2 + 2A r_2 r_1^* + B r_1^{*2} + B^* r_2^2 - 2(1 + r_2 r_1^*)(C^* r_2 - C r_1^*) \right\}.
\end{aligned}$$

For a stable system, they are:

$$\begin{aligned}
\lambda_1 &= \left\{ w(1 - |r_1|^2) - 2 \operatorname{Re}(U r_1) \right\}, \\
\lambda_2 &= -\left\{ w(1 - |r_1|^2) - 2 \operatorname{Re}(U r_1) \right\}, \\
\sigma &= \frac{1}{|r_1|^2} \left\{ w(1 - |r_1|^2) - 2 \operatorname{Re}(U r_1) \right\}, \\
\xi &= -|r_1|^2 \left\{ w(1 - |r_1|^2) - 2 \operatorname{Re}(U r_1) \right\}, \\
\delta_2 &= -\left\{ 2w - U^* r_1^* + \frac{U}{r_1^*} \right\}, \\
\phi_2 &= -|r_1|^2 \left\{ 2w - U^* r_1^* + \frac{U}{r_1^*} \right\}, \\
\delta_1 &= \phi_1 = \omega = \theta = 0,
\end{aligned}$$

$$\begin{aligned}
\Lambda &= \frac{1}{(1 + |r_1|^2)^2} \left\{ -\gamma(1 - |r_1|^2)^2 - 2A|r_1|^2 + 2\operatorname{Re}(Br_1^{*2}) + 4(1 - |r_1|^2) \operatorname{Re}(Cr_1^*) \right\}, \\
\Xi &= -\frac{|r_1|^2}{(1 + |r_1|^2)^2} \left\{ -\gamma(1 - |r_1|^2)^2 - 2A|r_1|^2 + 2\operatorname{Re}(Br_1^{*2}) + 4(1 - |r_1|^2) \operatorname{Re}(Cr_1^*) \right\}, \\
\Sigma &= -\frac{1}{|r_1|^2(1 + |r_1|^2)^2} \left\{ -\gamma(1 - |r_1|^2)^2 - 2A|r_1|^2 + 2\operatorname{Re}(Br_1^{*2}) + 4(1 - |r_1|^2) \operatorname{Re}(Cr_1^*) \right\}, \\
\Delta &= \frac{1}{(1 + |r_1|^2)^2} \left\{ 2\gamma(1 - |r_1|^2)^2 - A(1 - |r_1|^2) + \frac{B}{r_1^2} - B^*r_1^2 + \right. \\
&\quad \left. - \left(C^*r_1 - \frac{C}{r_1} \right) (1 - |r_1|^2) - 4\operatorname{Re}(Cr_1^*) \right\}, \\
\Phi &= -\frac{|r_1|^2}{(1 + |r_1|^2)^2} \left\{ 2\gamma(1 - |r_1|^2)^2 - A(1 - |r_1|^2) + \frac{B}{r_1^2} - B^*r_1^2 + \right. \\
&\quad \left. - \left(C^*r_1 - \frac{C}{r_1} \right) (1 - |r_1|^2) - 4\operatorname{Re}(Cr_1^*) \right\}, \\
\Omega &= \frac{1}{(1 + |r_1|^2)^2} \left\{ -4\gamma|r_1|^2 - A(|r_1|^4 + 1) - 2\operatorname{Re}(Br_1^{*2}) - 4(1 - |r_1|^2) \operatorname{Re}(Cr_1^*) \right\}, \\
\Theta &= \frac{|r_1|^2}{(1 + |r_1|^2)^2} \left\{ -4\gamma + 2A + Br_1^{*2} + \frac{B^*}{r_1^{*2}} - 4\left(\frac{C^*}{r_1^*} - Cr_1^* \right) \right\}.
\end{aligned}$$

Note that, in this case, the following relations hold:

$$\begin{aligned}
\lambda_2 &= -\lambda_1, \quad \xi = -|r_1|^2\lambda_1, \quad \phi_2 = |r_1|^2\delta_2, \\
\Xi &= -|r_1|^2\Lambda, \quad \Sigma = -\frac{1}{|r_1|^2}\Lambda, \quad \Phi = -|r_1|^2\Delta.
\end{aligned}$$

Appendix B

Expressions of the time evolutors

The entries of the time evolver $\hat{\mathcal{G}}_t^d$ in (2.30) for an unstable system are¹:

$$\begin{aligned}
(\hat{\mathcal{G}}_t^d)_{11} &= e^{-\gamma_1 t}, \\
(\hat{\mathcal{G}}_t^d)_{12} &= \frac{1}{\Delta\Gamma}(\Sigma + \sigma)(e^{-\gamma_2 t} - e^{-\gamma_1 t}), \\
(\hat{\mathcal{G}}_t^d)_{13} &= \frac{2}{\Delta\Gamma_+}(\Delta + \delta_1)(e^{-\Gamma-t} - e^{-\gamma_1 t}), \\
(\hat{\mathcal{G}}_t^d)_{14} &= \frac{2}{\Delta\Gamma_-}(\Delta^* + \delta_1^*)(e^{-\Gamma+t} - e^{-\gamma_1 t}), \\
(\hat{\mathcal{G}}_t^d)_{21} &= \frac{1}{\Delta\Gamma}(\Xi + \xi)(e^{-\gamma_2 t} - e^{-\gamma_1 t}), \\
(\hat{\mathcal{G}}_t^d)_{22} &= e^{-\gamma_2 t}, \\
(\hat{\mathcal{G}}_t^d)_{23} &= \frac{2}{\Delta\Gamma_-}(\Phi + \phi_1)(e^{-\gamma_2 t} - e^{-\Gamma-t}), \\
(\hat{\mathcal{G}}_t^d)_{24} &= \frac{2}{\Delta\Gamma_+}(\Phi^* + \phi_1^*)(e^{-\gamma_2 t} - e^{-\Gamma+t}), \\
(\hat{\mathcal{G}}_t^d)_{31} &= \frac{2}{\Delta\Gamma_+}(\phi_2 - \Phi^*)(e^{-\Gamma-t} - e^{-\gamma_1 t}), \\
(\hat{\mathcal{G}}_t^d)_{32} &= \frac{2}{\Delta\Gamma_-}(\delta_2 - \Delta^*)(e^{-\gamma_2 t} - e^{-\Gamma-t}), \\
(\hat{\mathcal{G}}_t^d)_{33} &= e^{-(\Gamma_- + \Lambda - \Omega + \lambda - \omega)t}, \\
(\hat{\mathcal{G}}_t^d)_{34} &= \frac{1}{\Delta m}(\Theta + \theta)e^{-\Gamma t} \sin \Delta m t, \\
(\hat{\mathcal{G}}_t^d)_{41} &= \frac{2}{\Delta\Gamma_-}(\phi_2^* - \Phi)(e^{-\Gamma+t} - e^{-\gamma_1 t}),
\end{aligned}$$

¹By definition, $\lambda \equiv (\lambda_1 + \lambda_2)/2$

$$\begin{aligned}
(\hat{\mathcal{G}}_t^d)_{42} &= \frac{2}{\Delta\Gamma_+}(\delta_2^* - \Delta)(e^{-\gamma_2 t} - e^{-\Gamma_+ t}), \\
(\hat{\mathcal{G}}_t^d)_{43} &= \frac{1}{\Delta m}(\Theta^* + \theta^*)e^{-\Gamma t} \sin \Delta m t, \\
(\hat{\mathcal{G}}_t^d)_{44} &= e^{-(\Gamma_+ + \Lambda - \Omega + \lambda - \omega^*)t}.
\end{aligned}$$

For a stable system they take the form:

$$\begin{aligned}
(\hat{\mathcal{G}}_t^d)_{11} &= \frac{1}{2}(e^{2\Lambda t} + 1) + \frac{\lambda_1}{2\Lambda}(e^{2\Lambda t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{12} &= \frac{\Sigma + \sigma}{2\Lambda}(e^{2\Lambda t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{13} &= -\frac{i}{\Delta m}\Delta(e^{i\Delta m t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{14} &= \frac{i}{\Delta m}\Delta^*(e^{-i\Delta m t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{21} &= \frac{\Xi + \xi}{2\Lambda}(e^{2\Lambda t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{22} &= \frac{1}{2}(e^{2\Lambda t} + 1) - \frac{\lambda_1}{2\Lambda}(e^{2\Lambda t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{23} &= -\frac{i}{\Delta m}\Phi(e^{i\Delta m t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{24} &= \frac{i}{\Delta m}\Phi^*(e^{-i\Delta m t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{31} &= -\frac{i}{\Delta m}(\phi_2 - \Phi^*)(e^{i\Delta m t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{32} &= -\frac{i}{\Delta m}(\delta_2 - \Delta^*)(e^{i\Delta m t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{33} &= e^{(\Omega + i\Delta m)t} \\
(\hat{\mathcal{G}}_t^d)_{34} &= \frac{1}{\Delta m}\Theta \sin \Delta m t \\
(\hat{\mathcal{G}}_t^d)_{41} &= \frac{i}{\Delta m}(\phi_2^* - \Phi)(e^{-i\Delta m t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{42} &= \frac{i}{\Delta m}(\delta_2^* - \Delta)(e^{-i\Delta m t} - 1) \\
(\hat{\mathcal{G}}_t^d)_{43} &= \frac{1}{\Delta m}\Theta^* \sin \Delta m t \\
(\hat{\mathcal{G}}_t^d)_{44} &= e^{(\Omega - i\Delta m)t}.
\end{aligned}$$

Appendix C

Neutral B mesons decays

C.1 Semileptonic decays

$$\begin{aligned}
\mathcal{P}_{h^-}(\mathbf{B}^0; \tau) = |\mathcal{M}_h|^2 e^{-\tau} \Big\{ & \sinh \delta \tau \left[\frac{|r_H|^2 - |r_L|^2}{|r_H + r_L|^2} - \text{Re } \lambda_h - \frac{\Lambda'}{\Delta \Gamma} \right] + \\
& + \cosh \delta \tau \left[\frac{|r_H|^2 + |r_L|^2}{|r_H + r_L|^2} - \text{Re } y_h + \frac{2}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Re } \Delta' \right] + \\
& + \sin \omega \tau \left[2 \frac{\text{Im}(r_H r_L^*)}{|r_H + r_L|^2} + \text{Im } \lambda_h - \frac{1}{2\Delta m} \text{Re } \Theta' \right] + \\
& + \cos \omega \tau \left[2 \frac{\text{Re}(r_L r_H^*)}{|r_H + r_L|^2} e^{-(\Omega' + \Lambda') \frac{\tau}{\Gamma}} - \text{Re } y_h - \frac{2}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Re } \Delta' \right] \Big\};
\end{aligned}$$

$$\begin{aligned}
\mathcal{P}_{h^+}(\mathbf{B}^0; \tau) = |\mathcal{M}_h|^2 e^{-\tau} \Big\{ & \sinh \delta \tau \left[-\text{Re } \mu_h - \frac{2}{\Gamma} \frac{\omega}{\delta^2 + \omega^2} \text{Im } \Delta' - \frac{\Lambda'}{\Delta \Gamma} \right] + \\
& + \cosh \delta \tau \left[\frac{2}{|r_H + r_L|^2} + \text{Re } y_h \right] + \\
& + \sin \omega \tau \left[\text{Im } \mu_h + \frac{2}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Im } \Delta' + \frac{1}{2\Delta m} \text{Re } \Theta' \right] + \\
& + \cos \omega \tau \left[-\frac{2}{|r_H + r_L|^2} e^{-(\Omega' + \Lambda') \frac{\tau}{\Gamma}} - \text{Re } y_h \right] \Big\};
\end{aligned}$$

$$\begin{aligned}
\mathcal{P}_{h^+}(\bar{\mathbf{B}}^0; \tau) = |\mathcal{M}_h|^2 e^{-\tau} \Big\{ & \sinh \delta\tau \left[\frac{|r_L|^2 - |r_H|^2}{|r_H + r_L|^2} - \text{Re } \mu_h - \frac{\Lambda'}{\Delta\Gamma} \right] + \\
& + \cosh \delta\tau \left[\frac{|r_H|^2 + |r_L|^2}{|r_H + r_L|^2} + \text{Re } y_h - \frac{2}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Re } \Delta' \right] + \\
& + \sin \omega\tau \left[2 \frac{\text{Im}(r_L r_H^*)}{|r_H + r_L|^2} - \text{Im } \mu_h - \frac{1}{2\Delta m} \text{Re } \Theta' \right] + \\
& + \cos \omega\tau \left[2 \frac{\text{Re}(r_H r_L^*)}{|r_H + r_L|^2} e^{-(\Omega' + \Lambda') \frac{\tau}{\Gamma}} + \text{Re } y_h + \frac{2}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Re } \Delta' \right] \Big\};
\end{aligned}$$

$$\begin{aligned}
\mathcal{P}_{h^-}(\bar{\mathbf{B}}^0; \tau) = |\mathcal{M}_h|^2 e^{-\tau} \Big\{ & \sinh \delta\tau \left[-\text{Re } \lambda_h + \frac{2}{\Gamma} \frac{\omega}{\delta^2 + \omega^2} \text{Im } \Delta' - \frac{\Lambda'}{\Delta\Gamma} \right] + \\
& + \cosh \delta\tau \left[2 \frac{|r_L r_H|^2}{|r_H + r_L|^2} - \text{Re } y_h \right] + \\
& + \sin \omega\tau \left[-\text{Im } \lambda_h - \frac{2}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Im } \Delta' + \frac{1}{2\Delta m} \text{Re } \Theta' \right] + \\
& + \cos \omega\tau \left[-2 \frac{|r_L r_H|^2}{|r_H + r_L|^2} e^{-(\Omega' + \Lambda') \frac{\tau}{\Gamma}} + \text{Re } y_h \right] \Big\}.
\end{aligned}$$

C.2 Decay in a CP eigenstate f

$$\begin{aligned}
\mathcal{P}_f(\mathbf{B}^0; \tau) = \frac{1}{2} |\mathcal{M}_h|^2 e^{-\tau} \Big\{ & \sinh \delta\tau \left[2 \frac{|1 - \lambda_H^f|^2 |r_H|^2 - |1 + \lambda_L^f|^2 |r_L|^2}{|r_H + r_L|^2} + \right. \\
& \left. - \frac{4}{\Gamma} \zeta_f \frac{\delta}{\delta^2 + \omega^2} \text{Re } \Delta' - \frac{4\Lambda'}{\Delta\Gamma} - \frac{4}{\Gamma} \frac{\omega}{\delta^2 + \omega^2} \text{Im } \Delta' \right] + \\
& + \cosh \delta\tau \left[2 \frac{|1 - \lambda_H^f|^2 |r_H|^2 + |1 + \lambda_L^f|^2 |r_L|^2}{|r_H + r_L|^2} \right] + \\
& + \frac{4}{\Gamma} \zeta_f \frac{\omega}{\delta^2 + \omega^2} \text{Im } \Delta' + \frac{4}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Re } \Delta' \Big] + \\
& + \sin \omega\tau \left[-4 \frac{\text{Im}(r_L r_H^* (1 + \lambda_L^f)(1 - \lambda_H^{f*}))}{|r_H + r_L|^2} e^{-(\Omega' + \Lambda') \frac{\tau}{\Gamma}} + \right. \\
& \left. - \frac{4}{\Gamma} \zeta_f \frac{\omega}{\delta^2 + \omega^2} \text{Re } \Delta' + \frac{4}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Im } \Delta' \right] + \\
& + \cos \omega\tau \left[4 \frac{\text{Re}(r_L r_H^* (1 + \lambda_L^f)(1 - \lambda_H^{f*}))}{|r_H + r_L|^2} e^{-(\Omega' + \Lambda') \frac{\tau}{\Gamma}} + \right. \\
& \left. - \frac{4}{\Gamma} \zeta_f \frac{\omega}{\delta^2 + \omega^2} \text{Im } \Delta' - \frac{4}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \text{Re } \Delta' \right] \Big\};
\end{aligned}$$

$$\begin{aligned}
\mathcal{P}_f(\overline{\mathbf{B}}^0; \tau) = & \frac{1}{2} |\mathcal{M}_h|^2 e^{-\tau} \left\{ \sinh \delta \tau \left[2 |r_L r_H|^2 \frac{|1 - \lambda_H^f|^2 - |1 + \lambda_L^f|^2}{|r_H + r_L|^2} + \right. \right. \\
& + \frac{4}{\Gamma} \zeta_f \frac{\delta}{\delta^2 + \omega^2} \operatorname{Re} \Delta' - \frac{4\Lambda'}{\Delta\Gamma} + \frac{4}{\Gamma} \frac{\omega}{\delta^2 + \omega^2} \operatorname{Im} \Delta' \Big] + \\
& + \cosh \delta \tau \left[2 |r_L r_H|^2 \frac{|1 - \lambda_H^f|^2 + |1 + \lambda_L^f|^2}{|r_H + r_L|^2} \right] + \\
& - \frac{4}{\Gamma} \zeta_f \frac{\omega}{\delta^2 + \omega^2} \operatorname{Im} \Delta' - \frac{4}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \operatorname{Re} \Delta' \Big] + \\
& + \sin \omega \tau \left[-4 |r_L r_H|^2 \frac{\operatorname{Im}((1 - \lambda_H^f)(1 + \lambda_L^{f*}))}{|r_H + r_L|^2} e^{-(\Omega' + \Lambda') \frac{\tau}{\Gamma}} + \right. \\
& + \frac{4}{\Gamma} \zeta_f \frac{\omega}{\delta^2 + \omega^2} \operatorname{Re} \Delta' - \frac{4}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \operatorname{Im} \Delta' \Big] + \\
& + \cos \omega \tau \left[4 |r_L r_H|^2 \frac{\operatorname{Re}((1 - \lambda_H^f)(1 + \lambda_L^{f*}))}{|r_H + r_L|^2} e^{-(\Omega' + \Lambda') \frac{\tau}{\Gamma}} + \right. \\
& + \frac{4}{\Gamma} \zeta_f \frac{\omega}{\delta^2 + \omega^2} \operatorname{Im} \Delta' + \frac{4}{\Gamma} \frac{\delta}{\delta^2 + \omega^2} \operatorname{Re} \Delta' \Big] \Big\}.
\end{aligned}$$

Appendix D

Dissipative parameters in the neutrino case

$$\begin{aligned}
\omega_1 &= \frac{g^2}{2V^2} \text{Im} [2B_{e\mu}(\vec{p}) + A_{e\mu\mu\mu}(\vec{p}) - A_{ee\mu e}(\vec{p}) - A_{eee\mu}(\vec{p}) + A_{\mu e\mu\mu}(\vec{p})], \\
\omega_2 &= -\frac{g^2}{2V^2} \text{Im} [2B_{\mu e}(\vec{p}) + A_{ee\mu e}(\vec{p}) - A_{e\mu\mu\mu}(\vec{p}) - A_{eee\mu}(\vec{p}) + A_{\mu e\mu\mu}(\vec{p})], \\
\omega_3 &= -\frac{g^2}{2V^2} \text{Im} [B_{ee}(\vec{p}) - B_{\mu\mu}(\vec{p}) - 2A_{ee\mu\mu}(\vec{p})], \\
u &= -\frac{g^2}{V^2} \text{Re} [A_{e\mu ee}(\vec{p}) + A_{\mu\mu\mu e}(\vec{p}) - A_{e\mu}(\vec{p})], \\
v &= \frac{g^2}{V^2} \text{Im} [A_{e\mu ee}(\vec{p}) + A_{\mu\mu\mu e}(\vec{p}) - A_{e\mu}(\vec{p})], \\
w &= \frac{g^2}{V^2} [A_{e\mu e\mu}(\vec{p}) - A_{\mu e\mu e}(\vec{p})], \\
a &= \frac{g^2}{2V^2} \{A_{ee}(\vec{p}) + A_{\mu\mu}(\vec{p}) - 2 \text{Re} [A_{ee\mu\mu}(\vec{p}) + A_{e\mu\mu e}(\vec{p})]\}, \\
b &= \frac{g^2}{V^2} \text{Im} [A_{e\mu\mu e}(\vec{p})], \\
c &= -\frac{g^2}{2V^2} \text{Re} [A_{ee\mu e}(\vec{p}) - A_{e\mu\mu\mu}(\vec{p}) + A_{eee\mu} - A_{\mu e\mu\mu}], \\
\alpha &= \frac{g^2}{2V^2} \{A_{ee}(\vec{p}) + A_{\mu\mu}(\vec{p}) - 2 \text{Re} [A_{ee\mu\mu}(\vec{p}) - A_{e\mu\mu e}(\vec{p})]\}, \\
\beta &= -\frac{g^2}{2V^2} \text{Im} [A_{eee\mu}(\vec{p}) - A_{\mu e\mu\mu}(\vec{p}) - A_{ee\mu e}(\vec{p}) + A_{e\mu\mu\mu}(\vec{p})], \\
\gamma &= \frac{g^2}{V^2} [A_{e\mu e\mu}(\vec{p}) + A_{\mu e\mu e}(\vec{p})].
\end{aligned}$$

Appendix E

Frequencies of counts

$$\rho_{13}(t) = \frac{1}{2} \left\{ \left[\mathcal{O}_t^{1,x}(0,0) - \mathcal{O}_t^{1,-x}(0,0) \right] + i \left[\mathcal{O}_t^{1,-y}(0,0) - \mathcal{O}_t^{1,y}(0,0) \right] \right\}$$

$$\rho_{14}(t) = \frac{1}{2} \left\{ \left[\mathcal{O}_t^{1,x}(0,0) - \mathcal{O}_t^{1,-x}(0,0) \right] - i \left[\mathcal{O}_t^{1,-y}(0,0) - \mathcal{O}_t^{1,y}(0,0) \right] \right\}$$

$$\rho_{23}(t) = \frac{1}{2} \left\{ \left[\mathcal{O}_t^{2,x}(0,0) - \mathcal{O}_t^{2,-x}(0,0) \right] + i \left[\mathcal{O}_t^{2,-y}(0,0) - \mathcal{O}_t^{2,y}(0,0) \right] \right\}$$

$$\rho_{24}(t) = \frac{1}{2} \left\{ \left[\mathcal{O}_t^{2,x}(0,0) - \mathcal{O}_t^{2,-x}(0,0) \right] - i \left[\mathcal{O}_t^{2,-y}(0,0) - \mathcal{O}_t^{2,y}(0,0) \right] \right\}$$

$$\rho_{31}(t) = \frac{1}{2} \left\{ \left[\mathcal{O}_t^{1,z}\left(\frac{\pi}{4}, 0\right) - \mathcal{O}_t^{2,z}\left(\frac{\pi}{4}, 0\right) \right] - i \left[\mathcal{O}_t^{1,z}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) - \mathcal{O}_t^{2,z}\left(\frac{\pi}{4}, \frac{\pi}{2}\right) \right] \right\}$$

$$\rho_{32}(t) = \frac{1}{2} \left\{ \left[\mathcal{O}_t^{1,-z}\left(\frac{\pi}{4}, 0\right) - \mathcal{O}_t^{2,-z}\left(\frac{\pi}{4}, 0\right) \right] - i \left[\mathcal{O}_t^{1,-z}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) - \mathcal{O}_t^{2,-z}\left(\frac{\pi}{4}, \frac{\pi}{2}\right) \right] \right\}$$

$$\rho_{41}(t) = \frac{1}{2} \left\{ \left[\mathcal{O}_t^{1,z}\left(\frac{\pi}{4}, 0\right) - \mathcal{O}_t^{2,z}\left(\frac{\pi}{4}, 0\right) \right] + i \left[\mathcal{O}_t^{1,z}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) - \mathcal{O}_t^{2,z}\left(\frac{\pi}{4}, \frac{\pi}{2}\right) \right] \right\}$$

$$\rho_{42}(t) = \frac{1}{2} \left\{ \left[\mathcal{O}_t^{1,-z}\left(\frac{\pi}{4}, 0\right) - \mathcal{O}_t^{2,-z}\left(\frac{\pi}{4}, 0\right) \right] + i \left[\mathcal{O}_t^{1,-z}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) - \mathcal{O}_t^{2,-z}\left(\frac{\pi}{4}, \frac{\pi}{2}\right) \right] \right\}$$

$$\begin{aligned} \rho_{33}(t) = & \frac{1}{4} \left\{ \left[\mathcal{O}_t^{1,x}\left(\frac{\pi}{4}, 0\right) - \mathcal{O}_t^{1,-x}\left(\frac{\pi}{4}, 0\right) \right] - i \left[\mathcal{O}_t^{1,y}\left(\frac{\pi}{4}, 0\right) - \mathcal{O}_t^{1,-y}\left(\frac{\pi}{4}, 0\right) \right] + \right. \\ & - \left[\mathcal{O}_t^{2,x}\left(\frac{\pi}{4}, 0\right) - \mathcal{O}_t^{2,-x}\left(\frac{\pi}{4}, 0\right) \right] + i \left[\mathcal{O}_t^{2,y}\left(\frac{\pi}{4}, 0\right) - \mathcal{O}_t^{2,-y}\left(\frac{\pi}{4}, 0\right) \right] + \\ & - \left[\mathcal{O}_t^{1,y}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) - \mathcal{O}_t^{1,-y}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) \right] - i \left[\mathcal{O}_t^{1,x}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) - \mathcal{O}_t^{1,-x}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) \right] \\ & \left. + \left[\mathcal{O}_t^{2,y}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) - \mathcal{O}_t^{2,-y}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) \right] + i \left[\mathcal{O}_t^{2,x}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) - \mathcal{O}_t^{2,-x}\left(\frac{\pi}{4}, -\frac{\pi}{2}\right) \right] \right\} \end{aligned}$$

$$\begin{aligned} \rho_{44}(t) = & \frac{1}{4} \left\{ \left[\mathcal{O}_t^{1,x} \left(\frac{\pi}{4}, 0 \right) - \mathcal{O}_t^{1,-x} \left(\frac{\pi}{4}, 0 \right) \right] - i \left[\mathcal{O}_t^{1,y} \left(\frac{\pi}{4}, 0 \right) - \mathcal{O}_t^{1,-y} \left(\frac{\pi}{4}, 0 \right) \right] + \right. \\ & - \left[\mathcal{O}_t^{2,x} \left(\frac{\pi}{4}, 0 \right) - \mathcal{O}_t^{2,-x} \left(\frac{\pi}{4}, 0 \right) \right] + i \left[\mathcal{O}_t^{2,y} \left(\frac{\pi}{4}, 0 \right) - \mathcal{O}_t^{2,-y} \left(\frac{\pi}{4}, 0 \right) \right] + \\ & + \left[\mathcal{O}_t^{1,y} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) - \mathcal{O}_t^{1,-y} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) \right] + i \left[\mathcal{O}_t^{1,x} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) - \mathcal{O}_t^{1,-x} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) \right] + \\ & \left. - \left[\mathcal{O}_t^{2,y} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) - \mathcal{O}_t^{2,-y} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) \right] - i \left[\mathcal{O}_t^{2,x} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) - \mathcal{O}_t^{2,-x} \left(\frac{\pi}{4}, -\frac{\pi}{2} \right) \right] \right\} \end{aligned}$$

Bibliography

- [1] R. Alicki and K. Lendi, *Quantum Dynamical Semigroups and Applications*, Lecture Notes in Physics, vol. 286, Springer-Verlag, 1987
- [2] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems*, Oxford University Press, 2002
- [3] E.B. Davies, *Quantum Theory of Open Systems*, Academic Press, 1976
- [4] C.P. Slichter, *Principles of magnetic resonance, with examples from solid state physics*, Springer-Verlag, 1990
- [5] W.H. Louisell, *Quantum Statistical Properties of Radiation*, Wiley, 1973
- [6] R.R. Puri, *Mathematical Methods of Quantum Optics*, Springer, 2001
- [7] C.W. Gardiner and P. Zoller, *Quantum Noise*, Springer, 2000
- [8] M.O. Scully and M.S. Zubairy, *Quantum Optics*, Cambridge University Press, 1997
- [9] L. Fonda, G.C. Ghirardi and A. Rimini, Rep. Prog. Phys. **41** (1978), 587
- [10] F. Benatti and R. Floreanini, Phys. Rev. **D 62** (2000), 125009 (hep-ph/0009283)
- [11] F. Benatti and R. Floreanini, Phys. Lett. **B 451** (1999), 422 (quant-ph/9902026)
- [12] F. Benatti, R. Floreanini and R. Romano, J. Phys. **A 35** (2002), 4955 (quant-ph/0205007)
- [13] H. Nakazato, M. Namiki and S. Pascazio, Int. J. Mod. Phys. **B 10** (1996), 247
- [14] M. Takesaki, *Theory of Operator Algebras*, vol. 1, Springer, 1979
- [15] M. Horodecki, P. Horodecki and R. Horodecki, Phys. Lett. **A 223** (1996), 1 (quant-ph/9605038)
- [16] K. Kraus, Ann. Phys. **64** (1971), 311
- [17] W. F. Stinespring, Proc. Amer. Math. Soc. **6** (1955), 211
- [18] M. D. Choi, Canad. J. Math. **24** (1972), 520
- [19] M. D. Choi, Linear Alg. Appl. **10** (1975), 285
- [20] O. Nachtmann, *Elementary particle physics: concepts and phenomena*, Springer-Verlag, 1990
- [21] S. Nakajima, Progr. Theor. Phys. **20** (1958), 948
- [22] R. Zwanzig, Lect. Theor. Phys. **3** (1960), 106

- [23] F. Haake, *Statistical treatment of open systems by generalized master equations*, Springer Tracts in Modern Physics, vol. 66, 1973
- [24] P. Pechukas, Phys. Rev. Lett. **73** (1994), 1060
- [25] A. Royer, Phys. Rev. Lett. **77** (1996), 3272
- [26] V. Romero-Rochin and I. Oppenheim, Physica **155 A** (1989), 52
- [27] A. Suarez, R. Silbey and I. Oppenheim, J. Chem. Phys. **97** (1992), 5101
- [28] S. Gnutzmann and F. Haake, Z. Phys. **B 101** (1996), 263
- [29] T. Yu, L. Diosi, N. Gisin and W.T. Strunz, Phys. Rev. **B 60** (1999), 91
- [30] P. Gaspard and M. Nagaoka, J. Chem. Phys. **111** (1999), 5668
- [31] P. Gaspard and M. Nagaoka, J. Chem. Phys. **111** (1999), 5676
- [32] R. Alicki, Phys. Rev. Lett. **75** (1995), 3020
- [33] H. Spohn, Rev. Mod. Phys. **52** (1980), 569
- [34] V. Gorini, A. Frigerio, M. Verri, A. Kossakowski and E.C.G. Sudarshan, Rep. Math. Phys. **13** (1978), 149
- [35] R. Dümcke and H. Spohn, Z. Phys. **B 34** (1979), 419
- [36] E. B. Davies, Math. Ann. **219** (1976), 147
- [37] K. Hepp and E.H. Lieb, Helv. Phys. Acta **46** (1973), 573
- [38] P. F. Palmer, J. Math. Phys. **18** (1977), 527
- [39] V. Gorini and A. Kossakowski, J. Math. Phys. **17** (1976), 1298
- [40] A. Frigerio, C. Novellone and M. Verri, Rep. Math. Phys. **12** (1977), 279
- [41] G. Lindblad, Commun. Math. Phys. **48** (1976), 119
- [42] V. Gorini, A. Kossakowski and E.C.G. Sudarshan, J. Math. Phys. **17** (1976), 821
- [43] F. Benatti, R. Floreanini and R. Romano, J. Phys. **A: Math Gen.** **35** (2002), L551 (quant-ph/0205007)
- [44] F. Benatti and R. Floreanini, Phys. Lett. **B 389** (1996), 100 (hep-th/9607059)
- [45] F. Benatti and R. Floreanini, Phys. Lett. **B 401** (1997), 337 (hep-ph/9704283)
- [46] F. Benatti and R. Floreanini, Nucl. Phys. **B 488** (1997), 335

- [47] F. Benatti and R. Floreanini, Nucl. Phys. **B 511** (1998), 550 ([hep-ph/9711240](#))
- [48] F. Benatti and R. Floreanini, Phys. Lett. **B 468** (1999), 287 ([hep-ph/9910508](#))
- [49] F. Benatti, R. Floreanini and R. Romano, Nucl. Phys. **B 602** (2001), 541 ([hep-ph/0103239](#))
- [50] I.M. Gel'fand *Lectures on Linear Algebra*, Interscience Tracts in Pure and Applied Mathematics, vol. 9, 1961
- [51] F. Benatti, R. Floreanini and M. Piani, "Curing non-positive evolutions in open system dynamics", submitted to EPJ **B**
- [52] K. Lendi, J. Phys. **A 20** (1987), 15
- [53] H. Spohn, Rep. Math. Phys. **10** (1976), 189
- [54] A. Frigerio, Comm. Math. Phys. **63** (1978), 269
- [55] F. Benatti and R. Floreanini, Banach Center Publications **43** (1998), 71
- [56] A. Wehrl, Rev. Mod. Phys. **50** (1978), 221
- [57] G.C. Branco, L. Lavoura and J.P. Silva, *CP Violation*, Clarendon Press, 1999
- [58] I.I. Bigi and A.I. Silva, *CP Violation*, Cambridge University Press, 2000
- [59] S. Hawking, Commun. Math. Phys. **87** (1983), 395
- [60] S. Hawking, Phys. Rev. **D 37** (1988), 904
- [61] S. Hawking, Phys. Rev. **D 53** (1996), 3099
- [62] S. Hawking and C. Hunter, Phys. Rev. **D 59** (1999), 044025
- [63] J. Ellis, N.E. Mavromatos and D.V. Nanopoulos, Phys. Lett. **B 293** (1992), 37
- [64] J. Ellis, N.E. Mavromatos and D.V. Nanopoulos, Int. J. Mod. Phys. **A 11** (1996), 1489
- [65] F. Benatti and R. Floreanini, Ann. Phys. **273** (1999), 58
- [66] M.S. Marinov, JETP Lett. **15** (1972), 479
- [67] M.S. Marinov, Sov. J. Nucl. Phys. **19** (1974), 173
- [68] M.S. Marinov, Nucl. Phys. **B 253** (1985), 609
- [69] J. Ellis, J.S. Hagelin, D.V. Nanopoulos and M. Srednicki, Nucl. Phys. **B 241** (1984), 381

- [70] R.A. Bertlmann and W. Grimus, Phys. Rev. **D 64** (2001), 056004 ([hep-ph/0101160](#))
- [71] S. Coleman, Nucl. Phys. **B 307** (1988), 867
- [72] S.B. Giddings and A. Strominger, Nucl. Phys. **B 307** (1988), 854
- [73] M. Srednicki, Nucl. Phys. **B 410** (1993), 143
- [74] W.G. Unruh and R.M. Wald, Phys. Rev. **D 52** (1995), 2176
- [75] L.J. Garay, Phys. Rev. Lett. **80** (1998), 2508
- [76] L.J. Garay, Phys. Rev. **D 58** (1998), 124015
- [77] P. Huet and M.E. Peskin, Nucl. Phys. **B 434** (1995), 3
- [78] J. Ellis, J.L. Lopez, N.E. Mavromatos and D.V. Nanopoulos, Phys. Rev. **D 53** (1996), 3846
- [79] P. Ball *et al.*, **B** decays at the LHC, CERN-TH-2000-101 ([hep-ph/0003238](#))
- [80] BTeV proposal, 2000, <http://www-btev.fnal.gov>
- [81] M.C. Banuls and J. Bernabeu, Phys. Lett. **B 423** (1998), 151 ([hep-ph/9710348](#))
- [82] M.C. Banuls and J. Bernabeu, Phys. Lett. **B 464** (1999), 117 ([hep-ph/9908353](#))
- [83] M.C. Banuls and J. Bernabeu, Nucl. Phys. **B 590** (2000), 19 ([hep-ph/0005323](#))
- [84] M.C. Banuls, Nucl. Phys. Proc. Suppl. **93** (2001), 268 ([hep-ph/0009317](#))
- [85] L. Lavoura, Phys. Lett. **B** (1998), 390 ([hep-ph/9806470](#))
- [86] L. Lavoura and J.P. Silva, Phys. Rev. **D 60** (1999), 066003 ([hep-ph/9902348](#))
- [87] BaBar Collaboration, The Babar Physics Book, SLAC-R-504, 1998
- [88] G. De Domenico and C. Yèche, BaBar-Note-409, 1998
- [89] C. Leonidopoulos, Ph.D. Thesis, Princeton University, 2000
- [90] F. Benatti and R. Floreanini, Mod. Phys. Lett. **A 14** (1999), 1519
- [91] L. Wolfenstein, Phys. Rev. **D 17** (1978), 2369
- [92] S.P. Mikheyev and A.Y. Smirnov, Sov. J. Nucl. Phys. **42** (1985), 913
- [93] S.P. Mikheyev and A.Y. Smirnov, Nuovo Cimento **9C** (1986), 17
- [94] A. Halprin, Phys. Rev. **D 34** (1986), 3462

- [95] L. Stodolsky, Phys. Rev. **D 36** (1987), 2273
- [96] R.F. Sawyer, Phys. Rev. **D 42** (1990), 3908
- [97] S. Samuel, Phys. Rev. **D 48** (1993), 1462
- [98] C.P. Burgess and D. Michaud, Annals Phys. **256** (1997), 1 ([hep-ph/9606295](#))
- [99] P. Bamert, C.P. Burgess and D. Michaud, Nucl. Phys. **B 513** (1998), 319 ([hep-ph/9707542](#))
- [100] A.B. Balantekin, Phys. Rept. **315** (1999), 123 ([hep-ph/9808281](#))
- [101] A.B. Balantekin, J.M. Fetter and F.N. Loreti, Phys. Rev. **D 54** (1996), 3941 ([astro-ph/9604061](#))
- [102] F.N. Loreti and A.B. Balantekin, Phys. Rev. **D 50** (1994), 4762 ([nucl-th/9406003](#))
- [103] E. Torrente-Lujan, Phys. Rev. **D 59** (1999), 073001 ([hep-ph/9807361](#))
- [104] H. Nunokawa, A. Rossi, V.B. Semikoz and J.W.F. Valle, Nucl. Phys. **B 472** (1996), 495 ([hep-ph/9602307](#))
- [105] G. Raffelt, G. Sigl and L. Stodolsky, Phys. Rev. Lett. **70** (1993), 2363 ([hep-ph/9209276](#))
- [106] C.W. Kim and A. Pevsner, *Neutrinos in physics and astrophysics*, Contemporary Concepts in Physics, vol. 8
- [107] R.N. Mohapatra and P.B. Pal, *Massive Neutrinos in Physics and Astrophysics*, 2nd ed., World Scientific, 1999
- [108] S.M. Bilenky, C. Giunti and W. Grimus, Prog. Part. Nucl. Phys. **43** (1999), 1 ([hep-ph/9812360](#))
- [109] S.M. Bilenky, “On the status of neutrino mixing and oscillations”, Report at the XVI Recontres de Physique de La Vallee d’Aoste, La Thuile, Aosta Valley (Italy) March 3-9, 2002 ([hep-ph/0205047](#))
- [110] W. Grimus, P. Stockinger and S. Mohanty, Phys. Rev. **D 59** (1999), 013011 ([hep-ph/9807442](#))
- [111] C. Giunti, JHEP **0211** (2002), 017 ([hep-ph/0205014](#))
- [112] D. McKay and H. Munczek, Phys. Rev. **D 19** (1979), 985
- [113] A. Zee, Phys. Lett. **B 93** (1980), 389
- [114] K.S. Babu, Phys. Lett. **B 203** (1988), 132

- [115] L.M. Johnson and D.W. McKay, Phys. Lett. **B 508** (2001), 290 (hep-ph/9909355)
- [116] P.D. Mannheim, Phys. Rev. **D 37** (1988), 1935
- [117] F. Benatti and R. Floreanini, Phys. Rev. **D 64** (2001), 085015 (hep-ph/0105303)
- [118] F. Benatti and R. Floreanini, JHEP **0002** (2000), 032, (hep-ph/0002221)
- [119] E. Lisi, A. Marrone and D. Montanino, Phys. Rev. Lett. **85** (2000), 1166
- [120] Y. Liu, L. Hu and M.-L. Ge, Phys. Rev. **D 56** (1997), 6648
- [121] C.-H. Chang, W.-S. Dai, X.-Q. Li, Y. Liu, F.-C. Ma and Z. Tao, Phys. Rev. **D 60** (1999), 033006
- [122] J. Schechter and J.W.F. Valle, Phys. Rev. **D 22** (1980), 2227
- [123] L.F. Li and F. Wilczek, Phys. Rev. **D 25** (1982), 143
- [124] J.K. Koh, C. Giunti and C.W. Kim, J. Korean Phys. Soc. **24** (1991), 275
- [125] J. Budimir and J.L. Skinner, J. Stat. Phys. **49** (1987), 1029
- [126] B.B. Laird, J.L. Skinner, J. Chem. Phys. **94** (1991), 4405
- [127] A. Peres, *Quantum Theory: Concepts and Methods*, Fundamental theories of physics, vol.57, Kluwer, 1993.
- [128] J.F. Clauser, M.A. Horne, A. Shimony and R.A. Holt, Phys. Rev. Lett. **23** (1969), 880
- [129] S. Basu, S. Bandyopadhyay, G. Kar and D. Home, Phys. Lett. **A 279** (2001), 281 (quant-ph/9907030)
- [130] M. Michler, H. Weinfurter and M. Zukowski, Phys. Rev. Lett. **84** (2000), 5457 (quant-ph/0009061)
- [131] J.L. Staudenmann, S.A. Werner, R. Colella and A.W. Overhauser, Phys. Rev. **A 21** (1980), 1419
- [132] S.A. Werner and A.G. Klein, Meth. Exp. Phys. **A 23** (1986), 259
- [133] V.F. Sears, *Neutron Optics: an introduction to the theory of neutron optical phenomena and their applications*, Oxford University Press, 1989
- [134] M. Utsuro, S. Kawano, T. Kawai and A. Kawaguchi, *Advance in neutron optics and related research facilities*, The Physical Society of Japan, 1996
- [135] K.C. Littrell, B.E. Allmann and S.A. Werner, Phys. Rev. **A56** (1997), 1767

- [136] B.E. Allmann, H. Kaiser, S.A. Werner, A.G. Wagh, V.C. Rakhecha and J. Summhammer, Phys. Rev. **A 56** (1997), 4420
- [137] H. Rauch and S.A. Werner, *Neutron interferometry: lessons in experimental quantum mechanics*, Oxford University Press, 2000